ELSEVIER

Clinica Chimica Acta 250 (1996) 121–265

Cumulative Contents

Vol. 151–249

Vol. 151/1, 1985

Changes in serum osteocalcin associated with parathyroid hormone infusion in X-linked hypophosphatemic rickets
Cole D.E.C., Gundberg C.M. 1

Comparison of ^{15}N analysis by optical emission spectrometry and mass spectrometry for clinical studies during total parenteral nutrition
Ragon A., Reynier J.P., Guiraud G., Francois G. 9

Improved method of thyroid peroxidase extraction from the human thyroid gland
Sugawara M., Thayer C.L., Kita T., Kuma K. 17

Comparison of enzymatic characteristics of creatine kinase BB from human healthy and tumor stomach tissues
Kanemitsu F., Okigaki T. 23

Salivary cortisol for the evaluation of Cushing's syndrome
Luthold W.W., Marcondes J.A.M., Wajchenberg B.L. 33

The role of urea in albumin dimerisation in nephrotic urines
Birtwistle R.J., Hardwicke J. 41

Serum and interstitial fluid apolipoprotein E levels in the healthy and in hyperlipoproteinemia type III as studied by radioimmunoassay
Gustafson S., Vessby B., Ostlund-Lindqvist A.M. 49

Enzyme linked immunosorbent assay for lactoferrin. Plasma and tissue measurements
Bezwoda W.R., Baynes R.D., Khan Q., Mansoor N. 61

Determination of choline-containing phospholipids in human bile and serum by a new enzyme sensor
Campanella L., Mascini M., Palleschi G., Tomassetti M. 71

Colorimetric measurement of serum magnesium with an IL micro centrifugal analyzer
Datta P., Graham G.A. 85

A quantitative IgE radioimmunoassay in dried blood spots suitable for neonatal screening of atopy
Adriaenssens K.M., Philips E.S., Colfs B. 91

Vol. 151/2, 1985

Serum bile acid composition in patients with cystic fibrosis
Setchell K.D.R., Smethurst P., Giunta A.M., Colombo C. 101

Release pattern of CK-MB and mitochondrial CK following myocardial ischaemia
Sylven C., Kallner A., Henze A. et al. 111

Heterogeneity of ^{125}I-labeled human thyroglobulin preparations
Spencer C.A., Platler B.W., Guttler R.B., Nicoloff J.T. 121

Urinary kallikrein: Assay validation and physiological variability
Ash K.O., Smith J.B., Lynch M. et al. 133

Glycosaminoglycan excretion in random samples of urine
Huang K.-C., Sukegawa K., Orii T. 141
Screening test for urinary glycosaminoglycans and differentiation of various mucopolysaccharidoses
Huang K.-C., Sukegawa K., Orii T. 147
Immunological study of tissue polypeptide antigen (TPA) — demonstration of keratin-like sites and blood group antigen-like sites on TPA molecules
Ochi Y., Ura Y., Hamazu M. et al. 157
Erythrocyte malondialdehyde release in vitro: A functional measure of vitamin E status
Cynamon H.A., Isenberg J.N., Nguyen C.H. 169
Direct automated assay method for serum or urine levels of ketone bodies
Harano Y., Ohtsuki M., Ida M. et al. 177
Basal level of human growth hormone (hGH) in normal serum
Hashida S., Nakagawa K., Ishikawa E., Ohtaki S. 185
A simple method for quantitative determination of lipoprotein-X (LP-X) in human serum
Panteghini M., Malchiodi A., Bonora R. 187

Vol. 151/3, 1985

Free drug measurements: Methodology and clinical significance
Kwong T.C. 193
Deficiency of acyl-CoA: Dihydroxyacetone phosphate acyltransferase in thrombocytes of Zellweger patients: A simple postnatal diagnostic test
Wanders R.J.A., Van Weringh G., Schrakamp G. et al. 217
Circulating prostatic acid phosphatase-immunoglobulin complexes in Sjögren's syndrome
Virji M.A. 223
Determination of zinc and copper in urine using Zeeman effect flame atomic absorption spectroscopy
Liska S.K., Kerkay J., Pearson K.H. 231
Determination of zinc in whole blood, plasma and serum using Zeeman effect flame atomic absorption spectroscopy
Liska S.K., Kerkay J., Pearson K.H. 237
A study of the microheterogeneity of transferrin in cirrhotic patients
Poupon R.E., Papoz L., Sarmini H., Elinck R. 245
Thermal transition studies of a mature xanthelasma by differential scanning calorimetry
Winder A.F., Bruckdorfer K.R., Fielder A.R. 253
Urinary steroid profiles in pregnancy by capillary column gas chromatography
Hahnel E., Wilkinson S.P., Hahnel R. 259
Age and sex dependency of the biochemical indices of bone remodelling
Stepan J.J., Tesarova A., Havranek T. et al. 273
Estimation of VLDL cholesterol in hyperlipidemia
Wilson P.W.F., Zech L.A., Gregg R.E. et al. 285
Benzoylalanine: Detection and identification of an alanine conjugate with benzoic acid in hyperammonemic patients treated with sodium benzoate
Shinka T., Inoue Y., Kuhara T. et al. 293
Interference by sulfonylurea drugs with the Jaffe method for creatinine
Roach N.A., Kroll M.H., Elin R.J. 301
Determination of seminal fructose using D-fructose dehydrogenase
Nakashima K., Takei H., Adachi O. et al. 307
A rapid, simple method for the micro-estimation of purine nucleoside phosphorylase activity in peripheral blood lymphocytes
Renouf J.A., Thong Y.H., Chalmers A.H. 311
A non-competitive enzyme-linked immunosorbent assay for measuring human plasma apolipoprotein B levels
James R.W., Ferrer M., Pometta D. 317

Vol. 152/1–2, 1985

A simple spectrophotometric method for quantitative fecal carbohydrate measurement
Ameen V.Z., Powell G.K. 3

Failure of ultracentrifugation as a means of separating plasma free insulin from immunoglobulin fraction prior to radioimmunoassay
Escobar L.J., White N.H., Santiago J.V., Gingerich R.L. 11

Comparison of the peptide and saccharide moieties of γ-glutamyltransferase isolated from neoplastic and non-neoplastic human liver tissue
Tsuchida S., Yamazaki T., Camba E.M. et al. 17

Immunoextraction of lipoamide dehydrogenase from cultured skin fibroblasts in patients with combined α-ketoacid dehydrogenase deficiency
Otulakowski G., Nyhan W., Sweetman L., Robinson B.H. 27

Platelet-associated proteins in human breast cyst fluids
Miller W.R., Dawes J. 37

Effect of nifedipine in hypothermic cardioplegia: A phosphorus-31 nuclear magnetic resonance study
Bernard M., Menasche P., Fontanarava E. et al. 43

Quantitative determination of phosphatidylglycerol in amniotic fluid by enzymatic assay
Farquharson J., Jamieson E.C., Paton R.D. et al. 55

Liquid chromatographic determination of inhibition of monoamine oxidase activity in platelet rich plasma of depressed patients treated with phenelzine
Gupta R.N., Steiner M. 63

Changes in human plasma α-d-mannosidase activity during pregnancy
Hirani S., Winchester B. 71

Level of plasma fibronectin in patients with peripheral vascular disease
Dionyssiou-Asteriou A., Karavassilis J. 79

An enzyme-amplified monoclonal immunoenzymometric assay for prostatic acid phosphatase
Moss D.W., Self C.H., Whitaker K.B. et al. 85

Copper dependent control of the enzymic and phagocyte induced degradation of some biopolymers, a possible link to ischemic inflammation
Hartmann H.J., Gartner A., Weser U. 95

Characterisation of high affinity binding sites of androgens in primary hepatocellular carcinoma
Wilkinson M.L., Iqbal M.J., Williams R. 105

Bile acid therapies applied to patients suffering from cerebrotendinous xanthomatosis
Koopman B.J., Wolthers B.G., Van der Molen J.C., Waterreus R.J. 115

Semi-automated determination of chromium in whole blood and serum by Zeeman electrothermal atomic absorption spectrophotometry
Schermaier A.J., O'Connor L.H., Pearson K.H. 123

Mass screening for glucose-6-phosphate dehydrogenase deficiency: Improved fluorescent spot test
Solem E., Pirzer C., Siege M. et al. 135

Comparison of a cytosol radioreceptor assay with a radioimmunoassay for 1,25-dihydroxyvitamin D in serum or plasma
De Leenheer A.P., Bauwens R.M. 143

The concentration of blood components related to fuel metabolism during prolonged fasting in children
Lamers K.J.B., Doesburg W.H., Gabreels F.J.M. et al. 155

Serum ferritin in type I diabetes
Carenini A., Ronchi D., Garbetta F. et al. 165

Decreased serum β-d-mannosidase activity in diabetic patients, in comparison with other glycosidases
Bernard M., Brochet C., Percheron F. 171

A qualitative agar gel immunoprecipitin (IP) test for detection of fecal occult human hemoglobin
Kim Y.D., Nolan J.M., Malkin A. et al. 175

Determination of free and total 3-methoxy-4-hydroxyphenylethylene glycol in human plasma by high-performance liquid chromatography with electrochemical detection
Semba J.I., Watanabe A., Takahashi R. 185

Age- and sex-related differences in urinary excretion of 6β-hydroxycortisol in humans
Nakamura J., Yakata M. 193
Effect of differences in saturation sensitivity of phospholipid stains on clinical predictivity of L/S ratios
Spillman T., Cotton D.B., Gonik B. 199
Determination of glutamine and glutamate in plasma of men and women by ion exchange chromatography
Bos K.D., Slump P. 205
Evaluation of a commercial reagent for precipitating human serum low-density lipoprotein
Shaikh M., Miller N.E. 213
Mevalonic aciduria: An inborn error of cholesterol biosynthesis?
Berger R., Smit G.P.A., Schierbeek H. et al. 219
A radiochemical microassay of S-adenosylhomocysteine hydrolase in lymphoid cells
Dinjens W.N.M., Chiba P., Van Laarhoven J.P.R.M. et al. 223
Determination of individual human faecal bile acids by gas-liquid chromatography after enzymatic deconjugation and simultaneous solvolysis and methylation using dimethoxypropane
Van Faassen A., Nagengast F.M., Hectors M. et al. 231

Vol. 152/3, 1985

Sympathoadrenal activity and plasma glucose effects on plasma dopamine-beta-hydroxylase levels in rats
Munoz J.A., Garcia-Estan J., Salom M.G. et al. 243
Octanoylglucuronide excretion in patients with a defective oxidation of medium-chain fatty acids
Duran M., Ketting D., Van Vossen R. et al. 253
Plasma α_1-antichymotrypsin in liver disease
Lindmark B.E., Eriksson S.G. 261
Concentrations of succinylacetone after homogentisate and tyrosine loading in healthy individuals with low fumarylacetoacetase activity
Kvittingen E.A., Leonard J.V., Pettit B.R., King G.S. 271
Effect of alcohol intake on high and low density lipoprotein metabolism in healthy volunteers
Malmendier C.L., Delcroix C. 281
Platelet monoamine oxidase: Specific activity and turnover number in schizophrenics and their families
Summers K.M., Andrew B., Gillespie C. et al. 289
Serum unconjugated bile acids: Qualitative and quantitative profiles in ileal resection and bacterial overgrowth
Setchell K.D.R., Harrison D.L., Gilbert J.M., Mupthy G.M. 297
A microassay for acid β-galactosidase activity toward asialofetuin
Mutoh T., Naoi M., Sobue I. et al. 307
Kinetic studies on neurotoxin-binding inhibitory antibodies in myasthenia gravis
Ito A., Furukawa S., Satoyoshi E., Sakurai H. 315
Limitations of new thyroid function tests in pregnancy
Gow S.M., Kellett H.A., Seth J. et al. 325
The determination of plasma oxalate concentrations using an enzyme/bioluminescent assay
Parkinson I.S., Kealey T., Laker M.F. 335
The determination of osteopetrotic phenotypes by selective inactivation of red cell carbonic anhydrase isoenzymes
Conroy C.W., Maren T.H. 347
An improved method for the simultaneous analysis of normal and modified urinary nucleosides and nucleobases by high performance liquid chromatography
Sander G., Wieland J., Topp H. et al. 355
The effect of laser deterioration of nephelometer performance
Salkie M.L. 363

Vol. 153/1, 1985

Study of a myoglobin test in patients hospitalized for suspected myocardial infarction
Mainard F., Massoubre B., Le Marec H., Madec Y. 1
Raised C-reactive protein levels in serum from smokers
Das I. 9
Crude spectrin extraction from reticulocyte-rich blood samples
Marik T., Kodicek M., Brabec V. 15
Differential investigation of the capacity of succinate oxidation in human skeletal muscle
Fischer J.C., Ruitenbeek W., Berden J.A. et al. 23
Simple enzymatic method for plasma branched-chain amino acids
Burrin J.M., Paterson J.L., Hall G.M. 37
Improved high performance liquid chromatographic determination of thiamine diphosphate in erythrocytes
Baines M. 43
Liquid chromatographic analysis of 4-hydroxy-3-methoxymandelic acid (VMA) and 4-hydroxy-3-methoxyphenylacetic acid (HVA) in urine
Fenech C., Guyon F., Dauchy P. 49
Determination of urinary lysozyme during aminoglycoside therapy — Appraisal of lysozyme inhibition caused by gentamicin and tobramycin
Morsky P., Ylitalo P., Koivula T. 55
Sensitive determination of cerebrospinal fluid pyruvate, lactate and glucose concentrations
Bozimowski D., Artiss J.D., Zak B. 63

Vol. 153/2, 1985

Induction of cytochromes P-450 in pancreatic disease: Consequence, coincidence or cause?
Acheson D.W.K., Rose P., Houston J.B., Braganza J.M. 73
Rapid method for measuring physiological concentrations of 25-hydroxyvitamin D levels in blood serum
Proszynska K., Lukaszkiewicz J., Jarocewicz N., Lorenc R.S. 85
A sensitive sandwich enzyme immunoassay for human myoglobin using Fab'-horseradish peroxidase conjugate: Methods and results in normal subjects and patients with various diseases
Nishida Y., Kawai H., Nishino H. 93
The effect of [^{125}I]thyroglobulin tracer heterogeneity on serum Tg RIA measurement
Spencer C.A., Platler B.W., Nicoloff J.T. 105
Vitamin E sufficiency in children with cholestasis: A comparison between erythrocyte peroxide hemolysis and serum α-tocopherol
Clark J.H., Nagamori K.E., Ellett M.L., Fitzgerald J.F. 117
Fast separation of the three main plasma lipoprotein classes by ultracentrifugation using vertical rotor and multiple discontinuous density gradient
Sclavons M.M., Cordonnier C.M., Mailleux P.M. et al. 125
Discrimination of phenylketonurics from persistent hyperphenylalaninemia patients using a simple phenylalanine loading test
Blitzer M.G., Bailey-Wilson J.E., Shapira E. 137
Serum angiotensin converting enzyme activity in patients with psoriasis
Ryder K.W., Epinette W.W., Jay S.J. et al. 143

Vol. 153/3, 1985

Detection of Bence-Jones protein by isoelectricfocussing of unconcentrated urine followed by nitrocellulose blotting and immunoperoxidase staining
Norden A.G.W., Fulcher L.M., Flynn F.V. 149

Microviscosity of tracheal aspirates in newborns with respiratory distress
Motta C., Touillon A.M., Labbe A., Dastugue B. 157
Some aspects of the glycoprotein and glycopolypeptide content of human gastric mucus
Ene M.D., Clamp J.R. 165
The effect of proteolytic degradation of plasma fibronectin on the responses of functional and immunometric assays for intact fibronectin
Cotton G., Brown R.A. 173
Creatine kinase: Purification, characterization, and tissue distribution of the MM subtypes
Mickelson J.K., Carlson C.J., Kaysen G.A., Rapaport E. 181
An automated determination of urinary 4-hydroxy-3-methoxymandelic acid
Kodama K., Nakata T., Yamanaka K., Aoyama M. 191
Assay of platelet monoamine oxidase in whole blood
Van Kempen G.M.J., Van Brussel J.L., Pennings Ed. J.M. 197
Increased plasma glycosidase and protease activity in uraemia: Possible role in the aetiology of the anaemia of chronic renal failure
Shannon J.S., Lappin T.R.J., Elder G.E. et al. 203
Pancreatic secretory trypsin inhibitor-like immunoreactivity in pancreatectomized patients
Halila H., Huhtala M.-L., Schroder T. et al. 209
Purification and determination of the modifying protein responsible for the post-synthetic modification of creatine kinase (EC 2.7.3.2) and enolase (EC 4.2.1.11)
Van Landeghem A.A.J., Soons J.B.J., Wever R.A. et al. 217
A reproducible procedure for measuring sodium transport in cultured human fibroblasts from normal and obese donors
Spurlock G., Morgan K., Mir M.A. 225
Normal pyruvate oxidation in Friedreich ataxia and Charcot-Marie-Tooth disease fibroblasts
Wilson W.G. 233
A simple procedure for the routine determination of aspartate aminotransferase and alanine aminotransferase with pyridoxal phosphate
Javier Gella F., Olivella T., Cruz Pastor M. et al. 241
Plasma vitamin A analysis by liquid chromatography: Probucol interference
Shephard G.S., Hough B.J., Van Stuijvenberg M.E., Labadarios D. 249
The determination of methaemoglobin levels by first derivative spectroscopy
Heales S.J.R., Bennett J. 253

Vol. 154/1, 1986

Distribution of serum zinc between albumin and α_2-macroglobulin in patients with different zinc metabolic disorders
Kiilerich S., Christiansen C. 1
A simple quantitative mecromethod of arginase assay in blood spots dried on filter paper
Konarska L., Tomaszewski L. 7
Ribonuclease from cytosolic fraction of human erythrocytes
Czajkowska B., Naskalski J.W., Sznajd J. 19
Quantitation of apolipoprotein B by polyclonal and monoclonal antibodies
Riesen W.F., Sturzenegger E., Imhof C., Mordasini R. 29
Gilbert's syndrome': Diagnosis by typical serum bilirubin pattern
Sieg A., Stiehl A., Raedsch R. et al. 41
Measurement of plasma oxalate in healthy subjects and in patients with chronic renal failure using immobilised oxalate oxidase
Kasidas G.P., Rose G.A. 49
Inhibitory activity of whole urine: A comparison of urines from stone formers and healthy subjects
Ryall R.L., Hibberd C.M., Mazzachi B., Marshall V.R. 59
Serum angiotensin converting enzyme activity in cigarette smokers
Haboubi N.A.A., Bignell A.H.C., Haboubi N.Y. 69

Vol. 154/2, 1986

Reduced trypsin binding capacity of α_2-macroglobulin in diabetes
Roberts R.C., Hall P.K., Nikolai T.F., McKenzie A.K. 85
Impaired amylase activities caused by binding of abnormal immunoglobulin A in patients with macroamylasemia
Adachi K., Suzuki K., Ohno Y., Sato B. 103
Development of a radioimmunoassay for human deoxyribonuclease I
Miyauchi K., Ogawa M., Shibata T. et al. 115
Urinary pseudouridine as a tumor marker in patients with small cell lung cancer
Tamura S., Fujii J., Nakano T. et al. 125
Discrimination of gamma-glutamyltranspeptidase from normal and carcinomatous pancreas
Yamaguchi N., Kawai K., Ashihara T. 133
Clarifying lipemic samples with an air-driven ultracentrifuge for determination of high-density lipoprotein cholesterol
Narahari Reddy M., Fitzsimons E.J. 141
Prenatal diagnosis of GM_2 gangliosidoses using a fluorogenic sulfated substrate
Inui K., Wenger D.A., Furukawa M. et al. 145

Vol. 154/3, 1986

Diagnosis and characterization of GM 2 gangliosidosis type II (Sandhoff disease) by analysis of the accumulating N-acetyl-glucosaminyl oligosaccharides with high performance liquid chromatography
Warner T.G., De Kramer R.D., Applegarth D., Mock A.K. 151
Aminoterminal procollagen III peptide elevation in alcoholics who are selenium and vitamin E deficient
Lu W., Bantok I., Desai S. et al. 165
The analysis of data from breast cancer estrogen and progesterone receptor assays: Scatchard plots are inferior to direct fitting by computer
Cooke R.R., McIntosh J.E.A. 171
Source of increased serum aspartate and alanine aminotransferase: cycloheximide effect on carbon tetrachloride hepatotoxicity
Pappas Jr. N.J. 181
Lipid peroxides in blood plasma and enzymatic antioxidative defence of erythrocytes in Down's syndrome
Kedziora J., Bartosz G., Gromadzinska J. et al. 191
Use of selective media for distinguishing variant forms of hypoxanthine phosphoribosyl transferase
Page T.M., Broock R.L., Nyhan W.L., Hernandez N.L. 195
Highly sensitive enzyme immunoassay for human brain aldolase C
Haimoto H., Kato K. 203
Reexamination of the conditions for processing and storing blood for plasma renin assay
Matsunaga M., Suzuki Y., Nakagawa K. et al. 213
Transglutaminase in erythrocytes of cystic fibrosis patients
Kamoun P.P., Droin V. 219
Determination of lithium in blood serum — A comparison between molecular absorption spectrophotometry and emission flame spectrometry
Popov P., Otruba V., Sommer L. 223
Detection of lactyl lactate in the urine of patients receiving Hicaliq infusion
Maeda K., Ohki T., Hosoi M. et al. 227
Interference by monoclonal proteins in the o-phthalaldehyde method for blood urea nitrogen
Pierce G.F., Garrett N.C., Koenig J. et al. 233
Solvolysis of mono- and disulfated bile acids by ethereal HCl, prepared by a simple technique
Dommes P., Breuer N.F., Goebell H. 237

The effect of storage conditions on the stability of carbamazepine and carbamazepine-10,11-epoxide in plasma
Mendez-Alvarez E., Soto-Otero R., Sierra-Marcuno G. 243
Bilirubin interference in the colorimetric assay of plasma vitamin E
Desai I.D., Martinez F.E. 247

Vol. 155/1, 1986

Isoforms of creatine kinase MM and MB in acute myocardial infarction: A clinical evaluation
Panteghini M., Cuccia C., Malchiodi A. et al. 1
Urinary neopterin, a marker of clinical activity in patients with Crohn's disease
Prior Ch., Bollbach R., Fuchs D. et al. 11
Differential measurement of monomeric and polymeric IgA by one-directional immunoelectrodiffusion
Meillet D., Tallet F., Gobert J.G. et al. 23
Purification of vitamin D binding protein from human plasma using high performance liquid chromatography
Taylor G.A., Mazhindu H.N., Findlay J.B.C., Peacock M. 31
Changes in renal enzyme activities following the administration of gentamicin to unilateral nephrectomized rats
Avidan G., Aladjem M., Israeli B., Bogin E. 43
Pregnancy-associated alpha$_2$-glycoprotein (α_2-PAG): Development of a sensitive enzyme-linked immunoassay and comparison of serum concentrations in adults and children
Gerrie L.M., Armstrong S.S., Horne C.H.W. 51
Preparation of ^{125}I-labelled arginine vasopressin for radioimmunoassay
Sadler W.A., Wright C.P., Livesey J.H. 61
Measurement of beta-2-microglobulin in serum and plasma by an enzyme-linked immunosorbent assay (ELISA)
Bjerrum O.W., Birgens H.S. 69
Zinc in hair and urine of paediatric patients
Van Wouwe J.P., De Wolff F.A., Van Gelderen H.H. 77
Bromcresol purple method for serum albumin gives falsely low values in patients with renal insufficiency
Maguire G.A., Price C.P. 83
Reliability of immunoglobulin determination by two methods
Mencke M.W., Weicker H., Wolf G.K. et al. 89
Assay of free and total fatty acids (as 2-nitrophenylhydrazides) by high performance liquid chromatography
Miwa H., Yamamoto M., Nishida T. 95

Vol. 155/2, 1986

Involvement of thiol proteases in galactosialidosis
Takeda E., Kuroda Y., Toshima K. et al. 109
Occurrence of short aliphatic diols in human blood: Identification by gas chromatography-mass spectrometry
Sisfontes L., Nyborg G., Jones A.W., Blomstrand R. 117
Patients with coeliac disease have a characteristic gliadin antibody pattern
Friis S.U., Noren O., Sjostrom H., Gudmand-Hoyer E. 133
Determination in human cerebrospinal fluid of glial fibrillary acidic protein, S-100 and myelin basic protein as indices of non-specific or specific central nervous tissue pathology
Noppe M., Crols R., Andries D., Lowenthal A. 143
A turbidimetric immunochemical method for determination of serum β_2-microglobulin using a centrifugal analyzer
Evrin P.E., Jansson N.O., Bergdahl A. 151

Pulsatile pattern of thyrotropin-release in normal men
Brabant G., Ranft U., Ocran K. et al. 159
CA 125 (ovarian tumour-associated antigen) in ascitic liver diseases
Bergmann J.F., Beaugrand M., Labadie H. et al. 163
Analysis of volatile free fatty acids in human urine by capillary column gas chromatography/mass spectrometry
Maltby D.A., Millington D.S. 167
Rapid detection of argininosuccinic aciduria and citrullinuria by fast atom bombardment and tandem mass spectrometry
Millington D.S., Maltby D.A., Roe C.R. 173
Zinc concentrations in mononuclear and polymorphonuclear leukocytes
Purcell S.K., Hambidge K.M., Jacobs M.A. 179
Comparison of zinc contents in human serum and plasma
Chen Wei-jie, Zhao Cheng-yi, Zheng Tian-li 185
Deficient beta-ketothiolase activity in leukocytes from a patient with 2-methylacetoacetic aciduria
Hiyama K., Sakura N., Matsumoto T., Kuhara T. 189

Vol. 155/3, 1986

Increase of plasma and red cell cholesterol sulfate levels in induced hypothyroidism in man
Van Doormaal J.J., Muskiet F.A.J., Jansen G. et al. 195
Peroxidase activity and estradiol receptors in human breast cancer
Gilabert B.P., Zamora V.A., Oriola A.P. et al. 201
An enzyme mediated, colorimetric method for the measurement of salicylate
Chubb S.A.P., Campbell R.S., Ramsay J.R. et al. 209
A double antibody radioimmunoassay for human α_1-microglobulin
Itoh Y., Nishino H., Enomoto H., Kawai T. 221
Glycosaminoglycans of skin and urine in pseudoxanthoma elasticum: Evidence for chondroitin 6-sulfate alteration
Longas M.O., Wisch P., Lebwohl M.G., Fleischmajer R. 227
Sweat sulfate concentrations are decreased in cystic fibrosis
Cole D.E.C., Landry D.A., Boucher M.J., Gillespie C.T. 237
Rapid diagnosis of vitamin D-dependent rickets type II by use of phytohemagglutinin-stimulated lymphocytes
Takeda E., Kuroda Y., Saijo T. et al. 245
Determination of mitochondrial aspartate aminotransferase in serum
Hirano K., Matsuda K., Adachi T. et al. 251
Estimation of NADH oxidation in human skeletal muscle mitochondria
Fischer J.C., Ruitenbeek W., Trijbels J.M.F. et al. 263
The relationship between plasma triglycerides, cholesterol, total lipids and lipid peroxidation products during human atherosclerosis
Ledwozyw A., Michalak J., Stepien A., Kadziolka A. 275
Affinity electrophoresis of human serum alkaline phosphatase isoenzymes in agarose gel containing lectin
Onica D., Sundblad L., Waldenlind L. 285
An O_2-based enzyme electrode for whole blood lactate measurement under continuous flow conditions
Weaver M.R., Vadgama P.M. 295
Fatty acyl composition of the major lipid classes of HDL_2 and HDL_3 of human serum
Devitt M.M., Headon D.R. 309
The effects of arginine, dextran and Haemaccel infusions on urinary albumin, β_2-microglobulin and N-acetyl-β-D-glucosaminidase
O'Reilly St. D.J., Parry E.S., Whicher J.T. 319
Tissue origin of serum placental-like alkaline phosphatase in cigarette smokers
Williams G.H., McLaughlin P.J., Johnson P.M. 329

Effect of ultraviolet light on creatinine measurement in jaundiced specimens
Nisbet J.A., D'Souza R. 335

Vol. 156/1, 1986

Oncogenes
Garrett C.T. 1
Ganglioside loading of cultured fibroblasts: A provocative method for the diagnosis of the G(M2) gangliosidoses
Charrow J., Binns H.J. 41
Development of a sensitive enzyme immunoassay for human epidermal growth factor (Urogastrone)
Kurobe M., Tokida N., Furukawa S. et al. 51
Energy metabolism in adenosine deaminase-inhibited human erythrocytes
Buc H.A., Thuillier L., Hamet M. et al. 61
Determination of oxalate in urine by flow injection analysis
Gaetani E., Laureri C.F., Vitto M. et al. 71
Diuresis-dependent excretion of multiple forms of renal brush-border enzymes in urine
Jung K., Schulze G. 77
The analysis of thiodiglycollic acid by selected ion monitoring
Pettit B.R. 85
HPLC assay of phenylalanine and tyrosine in blood spots on filter paper
Roesel R.A., Blankenship P.R., Hommes F.A. 91
A simple enzymatic fluorimetric method for the determination of triglycerides in 10 μl of serum
Gleeson M., Maughan R.J. 97
Blood plasma fluoride in haemodialysed patients
Chaleil D., Simon P., Tessier B. et al. 105
Determination of silicon, calcium, magnesium and phosphorus in urine using inductively-coupled plasma emission spectrometry and a matrix-matching technique
Tanaka T., Hayashi Y. 109
Measurement of radioactivity of pyruvate and other 2-oxo-acids labelled at the 1-position
Van Laack H.L.J.M., Janssen A.J.M., Ruitenbeek W. et al. 115

Vol. 156/2, 1986

Pancreatic secretory trypsin inhibitor from human amniotic fluid and fetal and neonatal urine: Concentrations and physicochemical characterization
Kolho K.-L., Huhtala M.-L., Jalanko H. et al. 123
Quantification of human serum apolipoprotein A-I by zone immunoelectrophoresis assay and a procedure for the preparation of an A-I standard
Holmquist L., Vesterberg O. 131
Quantitation of CK-MB release: Diagnostic utility in coronary artery bypass grafting
Devine J.E., Wiens R.D., Halstead J.M., Codd J.E. 145
On-line clarification for the measurement of serum glucose in hyperlipidemic specimens
Artiss J.D., Strandbergh D.R., Sharma A., Zak B. 151
Myocardial chondroitin sulfates D and E in a case of acute carbon monoxide poisoning
Ohishi H., Binette J.P., Schmid K. 157
Probe of the $(Ca^{2+} + Mg^{2+})$-ATPase in erythrocyte membranes of cystic fibrosis patients
Hunsinger R.N., Cheung H.C. 165
Sulfogalactocerebroside and bis-(monoacylglyceryl)-phosphate as activators of spleen glucocerebrosidase
Prence E.M., Garrett K.O., Panitch H. et al. 179
Deficient activities and proteins of peroxisomal β-oxidation enzymes in infants with Zellweger syndrome
Suzuki Y., Orii T., Mori M. et al. 191

The measurement of glycosyiated albumin by reduction of alkaline nitro-blue tetrazolium
Walker S.W., Howie A.F., Smith A.F. 197
Determination of volatile fatty acids by gas chromatography on a capillary and a megabore column
Duisterwinkel F.J., Wolthers B.G., Van Der Slik W., Dankert J. 207
Use of pooled human serum in the standardization process of the serum fructosamine determination for the estimation of glycosylated serum proteins
Smid E., Ferencz A., Fodor M. 215
Reduced imprecision of the radioenzymatic assay of plasma catecholamines by improving the stability of the internal standards
Hoffmann J.J.M.L., Willemsen J.J., Lenders J.W.M., Benraad Th. J. 221
What information should be provided in the package inserts of commercial reference materials in clinical chemistry?
Broughton P.M.G. 227

Vol. 156/3, 1986

Enzymatic assessment of myocardial injury after infarction or cardiac surgery. Is isoenzyme analysis required?
Hermens Th. W., Willems G.M., Davids H.A. et al. 235
Quality control of azure B preparations by liquid chromatography and standardization with azure B tetrafluoroborate
Nelis H.J.C.F., De Leenheer A.P. 247
Sensitive quantitation of carcinoembryonic antigen in cerebrospinal fluid and its barrier-dependent differentiation
Reiber H., Jacobi C., Felgenhauer K. 259
Assessment of the AminoStat-FLM immunoagglutination test for phosphatidylglycerol in amniotic fluid
Farquharson J., Jamieson E.C., Berry E. et al. 271
Diagnosis of inherited adenylosuccinase deficiency by thin-layer chromatography of urinary imidazoles and by automated cation exchange column chromatography of purines
De Bree P.K., Wadman S.K., Duran M., Fabery de Jonge H. 279
ELISA for human proinsulin
Hartling S.G., Dinesen B., Kappelgard A.-M. et al. 289
Changes in fatty acid profiles of plasma, erythrocytes and polymorphonuclear leukocytes in induced hypothyroidism in man: Indirect evidence for altered Δ6 desaturase activity
Van Doormaal J.J., Muskiet F.A.J., Martini I.A., Doorenbos H. 299
Increase in blood plasma levels of boron and strontium in hemodialyzed patients
Mauras Y., Ang K.S., Simon P. et al. 315
Plasma and red cell lipids in alcoholics with macrocytosis
Clemens M.R., Kessler W., Schied H.W. et al. 321
Glycated haemoglobin and glycated plasma protein, a comparative study
Ross I.S., Mitchell A.F., Stroud R.E. 329
An accurate, simple and rapid test for detecting elevated levels of C-reactive protein in serum by agglutination of fat emulsion
Hulman G., Fuller M., Pearson H.J., Bell P.R.F. 337

Vol. 157/1, 1986

Clinical applications of gene probes in human genetic disease, malignancy, and infectious disease
Lowe J.B. 1
Erythrocyte cation transport activities as a function of cell age
Hentschel W.M., Wu L.L., Tobin G.O. et al. 33

A case of rheumatoid arthritis with various enzyme-immunoglobulin complexes
Maekawa M., Sudo K., Kanno T. 45
Serum creatine kinase isoenzyme MB concentration after endomyocardial biopsy
Chapelle J.P., El Allaf D., El Allaf M. et al. 55
Oligomerization of pregnancy-specific β_1-glycoprotein (SP_1) at physiologic pH and ionic strength
Rosen S.W., Calvert I., Lee N. et al. 65
Fluorometric determination of nanogram quantities of protein in small samples: Application to calcium-transport adenosine triphosphatase
Bridges M.A., McErlane K.M., Kwong E. et al. 73
The influence of serum and serum proteins on calcium oxalate crystal growth and aggregation
Edyvane K.A., Ryall R.L., Marshall V.R. 81
Isolation and characterisation of lactoferrin separated from human whey by adsorption chromatography using Cibacron Blue F3G-A linked affinity adsorbent
Bezwoda W.R., Mansoor N. 89
A procedure for large scale purification of human plasma amyloid P component
Ohkubo I., Sahashi W., Namikawa C. et al. 95
Colorimetric assay for carboxypeptidase N in serum
Hendriks D., Van Sande M., Scharpe S. 103
Lysosomal enzymes in chorionic villi, cultured amniocytes, and cultured skin fibroblasts
Evans M.I., Moore C., Kolodny E.H. et al. 109
Estimation of aluminoxamine and ferrioxamine in plasma by high performance liquid chromatography
Hughes H., Hagen L.E., Cameron E.C., Sutton R.A.L. 115
An improved spectrophotometric assay of urinary metanephrines
Crawford G.A., Gallery E.D.M., Gyory A.Z. 121

Vol. 157/2, 1986

IgD plasma cell leukemia associated with pyroglobulinemia and pyroglobulinuria
Gastearena J., Fernandez F.J., Orue M.T. et al. 133
Guanidino compounds in uraemic dialysed patients
De Deyn P., Marescau B., Lornoy W. et al. 143
Evaluation and comparison of cathodic trypsin-like immunoreactivity, pancreatic lipase and pancreatic isoamylase in the diagnosis of acute pancreatitis in 849 consecutive patients with acute abdominal pain
Moller-Petersen J., Klaerke M., Dati F. 151
Gold staining in cellulose acetate membranes
Righetti P.G., Casero P., Del Campo G.B. 167
Estimation of vitamin D_3 and 25-hydroxyvitamin D_3 in muscleand adipose tissue of rats and man
Lawson D.E.M., Douglas J., Lean M., Sedrani S. 175
Internalization of exogenous gangliosides in cultured skin fibroblasts for the diagnosis of mucolipidosis IV
Zeigler M., Bach G. 183
Enzyme electrode for the measurement of lactate in undiluted blood
Mullen W.H., Churchouse S.J., Keedy F.H., Vadgama P.M. 191
Serum calcium status in health and disease: A comparison of measured and derived parameters
White T.F., Farndon J.R., Conceicao S.C. et al. 199
Comparison of sampling methods for the analysis of methane in expired air
Peled Y., Gilat T. 215

Vol. 157/3, 1986

Quantitation of trans-vitamin K1 in small serum samples by off-line multidimensional liquid chromatography
Van Haard P.M.M., Engel R., Pietersma-de Bruyn A.L.J.M. 221

Enzyme immunoassay of serum cortisol using a new transferable needle lid technique
Dhar T.K., Voss E., Schoneshofer M. 231
Preliminary report on a case of apolipoproteins CI and CII deficiency
Dumon M.-F., Clerc M. 239
Separation of apolipoprotein B subfractions by high performance gel permeation chromatography
Schwandt P., Richter W.O., Weisweiler P. 249
Metabolic conversion of L-[U-^{14}C]phenylalanine to respiratory $^{14}CO_2$ in healthy subjects, phenylketonuria heterozygotes and classic phenylketonurics
Lehmann W.D., Fischer R., Heinrich H.C. et al. 253
ELISA of 6-beta-hydroxycortisol in human urine: Diurnal variations and effects of antiepileptic therapy
Zhiri A., Wellman-Bednawska M., Siest G. 267
Capillary gas chromatographic profiling of urinary, plasma and erythrocyte sugars and polyols as their trimethylsilyl derivatives, preceded by a simple and rapid prepurification method
Jansen G., Muskiet F.A.J., Schierbeek H. et al. 277
Diagnostic value of microalbuminuria in pre-eclampsia
Irgens-Moller L., Hemmingsen L., Holm J. 295
A non-random distribution of transferrin iron in fresh human sera
Van Eijk H.G., Van Noort W.L. 299
The reliability of the use of para toluene sulfonic acid for simultaneous hydrolysis and quantitation of both N-acetyl-glucosamine and amino acids in human transferrins
Van Eijk H.G., Van Noort W.L. 305
Multiwavelength analyses of derivative spectra: An application for the determination of urinary porphyrins
Walters M., Thornton D.J. 311

Vol. 158/1, 1986

Normal human serum contains two forms of acetylcholinesterase
Sorensen K., Brodbeck U., Rasmussen A.G., Norgaard-Pedersen B. 1
High molecular weight angiotensinogen: A pregnancy associated protein
Tewksbury D.A., Tryon E.S., Burrill R.E., Dart R.A. 7
A map of urine proteins based on one-dimensional SDS-polyacrylamide gel electrophoresis and Western blotting using one microliter of unconcentrated urine
Kshirsagar B., Wiggins R.C. 13
A modified radioimmunoassay for arylsulfatase A in human serum and urine
Laidler P.M., Silberring J., Van Etten R.L. 23
Measurement of total HDL, HDL_2 and HDL_3 by dextran-sulfate-$MgCl_2$ precipitation technique in human serum
Talameh Y., Wei R., Naito H. 33
Changes in plasma osteocalcin concentration following treatment with stanozolol
Couch M., Preston F.E., Malia R.G. et al. 43
An improved approach to thyroid function testing in patients with non-thyroidal illness
Gow S.M., Elder A., Caldwell G. et al. 49
Development and evaluation of a simple, direct, solid-phase radioimmunoassay of serum cortisol from readily available reagents
McConway M.G., Chapman R.S. 59
Lipid associated tissue ferritin
Cham B.E., Roeser P., Nikles A.C., Ridgway K.M. 71
Physiological concentrations of human epidermal growth factor in biological fluids: Use of a sensitive enzyme immunoassay
Joh R., Itoh M., Katsumi K. et al. 81
Acid stable trypsin inhibitor in bile
Yamamoto T., Sumi H., Maruyama M. et al. 91
Highly sensitive enzyme immunoassay for human creatine kinase MM and MB isozymes
Kato K., Shimizu A. 99

Excretion of octopamine metabolites in neuroblastoma
Couch M.W., Greer D.M., Williams C.M. 109
Improved assay of the enzymes of glutathione synthesis: γ-glutamylcysteine synthetase and glutathione synthetase
Beutler E., Gelbart T. 115

Vol. 158/2, 1986

Determination of ionised calcium by ion selective electrode is not independent of albumin concentration
Freaney R., Egan T., McKenna M.J. et al. 129
Determination of urinary estradiol using an enzymatic method during the menstrual cycle. Comparison with a method using isotope dilution-mass spectrometry (ID-MS)
Patricot M.C., Mathian B., Serpentie S., Revol A. 139
Identification of lignans and phytoestrogens in urine of chimpanzees
Adlercreutz H., Musey P.I., Fotsis T. et al. 147
Deficient activity of glucocerebrosidase in urine from patients with type 1 Gaucher disease
Aerts J.M.F.G., Donker-Koopman W.E., Koot M. et al. 155
Comparison of a tumour-derived form of intestinal alkaline phosphatase with foetal and adult intestinal alkaline phosphatases
Moss D.W., Parmar C.R., Whitaker K.B. 165
Assay of urinary 5-hydroxyindole-3-acetic acid: Two methodologies compared
Baars J.D., Van Haard P.M.M., Lombarts A.J.P.F. 173
Detection of carriers of cerebrotendinous xanthomatosis
Koopman B.J., Waterreus R.J., Van Den Brekel H.W.C., Wolthers B.G. 179
The effect of sotalol on plasma lipoproteins and apolipoproteins
Northcote R.J., Packard C.J., Ballantyne D. 187
Tissue homogenization using a micro-dismembrator for the measurement of enzyme activities
Peters G.J., Laurensse E.J., Leyva A., Pinedo H.M. 193
An improved source receptor for 1,25-dihydroxyvitamin D3 assay
Blayau M., Leray G., Prodhomme C. et al. 199

Vol. 158/3, 1986

Chemotherapeutic agent cisplatin monitoring in biological fluids by means of inductively-coupled plasma emission spectrometry (ICP-AES)
Dominici C., Alimonti A., Caroli S. et al. 207
N-Acetylaspartic aciduria in a child with a progressive cerebral atrophy
Kvittingen E.A., Guldal G., Borsting S. et al. 217
Crystal inhibition: The effects of polyanions on calcium oxalate crystal growth
Fellstrom B., Danielson B.G., Ljunghall S., Wikstrom B. 229
Errors in measurements of CO_2 with the use of drying agents
Elia M., McDonald T., Crisp A. 237
Sarcoplasmic reticulum Ca^{2+}-ATPase and acylphosphatase activities in muscle biopsies from patients with Duchenne muscular dystrophy
Landi N., Nassi P., Liguri G. et al. 245
Mitochondrial myopathy: Tissue-specific expression of a defect in ubiquinol-cytochrome c reductase
Darley-Usmar V.M., Watanabe M., Uchiyama Y. et al. 253
Cholesterol efflux from cultured skin fibroblasts in 12 patients clinically heterozygous for familial hypercholesterolemia
Esteva O., Baudet M.-F., Jacotot B. 263
Hepatic acyl-CoA:cholesterol acyltransferase. Development of a standard assay and determination in patients with cholesterol gallstones
Smith J.L., De Jersey J., Pillary S.P., Hardie I.R. 271

Rapid assay of pyruvate with a centrifugal analyser using a commercially available test kit
Van Erven A., Huisman R., Vermunt K. 283
Enzyme immunoassay for ceruloplasmin: Application to cancer patient serum
DiSilvestro R.A., David E.A. 287
Radioimmunoassay for nonenzymatically glycated protein in human serum
Nakayama H., Taneda S., Manda N. et al. 293

Vol. 159/1, 1986

Peroxisomal β-oxidation of palmitoyl-CoA in human liver homogenates and its deficiency in the cerebro-hepato-renal (Zellweger) syndrome
Wanders R.J.A., Van Roermund C.W.T., De Vries C.T. et al. 1
1,6-Anhydro-β-D-glucopyranose (β-glucosan), a constitutent of human urine
Dorland L., Wadman S.K., Fabery de Jonge H., Ketting D. 11
An immunoglobulin G inhibiting lactate dehydrogenase activity
Hirano T., Matsuzaki H., Miura M. et al. 17
Purification and characterization of pancreatic secretory trypsin inhibitor in human gastric mucosa
Shibata T., Ogawa M., Matsuda K. et al. 27
Microheterogeneity of Kasahara isozyme
Hada T., Imanishi H., Yamamoto T. et al. 37
Radioimmunoassay for atrial natriuretic peptide: Method and results in normal subjects and patients with various diseases
Nishiuchi T., Saito H., Yamasaki Y., Saito S. 45
Absence of argininosuccinate lyase protein in the liver of two patients with argininosuccinic aciduria
Kobayashi K., Itakura Y., Saheki T. et al. 59
The nature of gastric secretions in a patient with Zollinger-Ellison syndrome without peptic ulceration
Clamp J.R., Fairclough P.D. 69
Glutathione-linked enzyme activities in red cell aging
Imanishi H., Nakai T., Abe T., Takino T. 73
A simple method to diagnose adrenoleukodystrophy using a dried blood spot on filter paper
Nishio H., Kodama S., Yokoyama S. et al. 77
Placental alkaline phosphatase in an omental tumor
Miura M., Matsuzaki H., Sakagishi Y., Sekine T. 83

Vol. 159/2, 1986

Characteristics of antisera to antigenic forms of 1,25-dihydroxycholecalciferol
Brown W.B., Peacock M. 111
Analysis of conjugated and unconjugated bile acids in serum and jejunal fluid of normal subjects
Tangerman A., Van Schaik A., Van der Hoek E.W. 123
The subcellular distribution and levels of calmodulin in jejunal biopsies from control subjects and patients with coeliac disease
Macpherson A.J.S., Bjarnason I., Peters T.J. 133
Measurement of human growth hormone (hGH) using a rapid immunochemiluminometric assay
Weeks I., Woodhead J.S. 139
Lipoprotein and apoprotein levels in postmenopausal women during treatment with norethisterone
Farish E., Fletcher C.D., Hart D.M. et al. 147
Spin-echo and 2-dimensional ^1H nuclear magnetic resonance studies on urinary metabolites from patients with 2-methylacetocetyl CoA thiolase deficiency
Williams S.R., Iles R.A., Chalmers R.A. 153

Direct demonstration of an inhibitor of sodium, potassium dependent adenosine triphosphatase (Na^+,K^+-ATPase) in plasma from normotensive and hypertensive subjects
Roulston J.E., Traill K., Wathen C.G. 163
Determination of aromatic-L-amino acid decarboxylase in human plasma
Boomsma F., Van der Hoorn F.A.J., Schalekamp M.A.D.H. 173
A new solid-phase assay for biotin and biocytin and its application to the study of patients with biotinidase deficiency
Ping Wai Chan, Bartlett K. 185
Measurement of magnesium in serum and urine with a random access analyzer by use of a modified xylidyl blue-I procedure
Ratge D., Kohse K.P., Wisser H. 197
An enzyme-linked immunosorbent assay (ELISA) for urinary free cortisol
Lewis J.G., Manley L., Townsend J.C., Elder P.A. 205
Isolation of platelets from human blood by free-flow electrophoresis
Wilson R.B.J., Graham J.M. 211

Vol. 159/3, 1986

Creatine kinase and lactate dehydrogenase isoenzymes in stage D prostatic carcinoma
Strom S., Bendz R. 219
Determination of pipecolic acid in urine and plasma by isotope dilution mass fragmentography
Van Den Berg G.A., Breukelman H., Elzinga H. et al. 229
Enzyme-linked immunoadsorbent assay (ELISA) for calcium-binding protein of human small intestine. A quantitative method for measuring calcium-binding protein in small intestinal biopsies
Staun M. 239
Carnitine deficiency in haemodialysed patients
Rodriguez-Segade S., Del la Pena C.A., Paz et al. J.M. 249
Enzyme-linked immunosorbent assay for amitriptyline and other antidepressants using a monoclonal antibody
Denis H., Marullo S., Hoebeke J., Strosberg A.D. 257
Quantitation of human serum apolipoprotein B by enzyme immunoassay in excess antigen
Fievet C., Dedieu A., Kandoussi A. et al. 269
β-N-Acetyl-D-glucosaminidase isoenzymes from human amnionic membranes
Orlacchio A., Emiliani C., Di Renzo G.C., Cosmi E.V. 279
Citrate interference with the determination of acetylcarnitine: A method for its elimination
Cooper M.B., Forte C.A., Jones D.A. 291

Vol. 160/1, 1986

Analysis of plasma lipoproteins by ultracentrifugation in a new fixed angle rotor: Evaluation of a phosphotungstic acid/$MgCl_2$ precipitation and a quantitative lipoprotein electrophoresis assay
Marz W., Gross W. 1
Clinical significance and characteristics of creatine kinase-immunoglobulin complexes in sera from patients with malignant tumors
Kanemitsu F. 19
Sandwich electroimmunofixation (SEIF) for the assessment of isoenzyme specificities of immunoglobulins isolated from enzyme-immunoglobulin complexes
Kanemitsu F., Okigaki T. 27
A sandwich immunoassay for human prolyl 4-hydroxylase using monoclonal antibody
Yoshida S., Bai Y., Muragaki Y. et al. 37
Unusual trihydroxy bile acids in the urine of healthy humans
Nakashima T., Sano A., Seto Y. et al. 47

Functional activities and nonenzymatic glycosylation of plasma proteinase inhibitors in diabetes
Hall P., Tryon E., Nikolai T.F., Roberts R.C. 55
5-Amino-2-nitrobenzoic acid as a reference material for the determination of γ-GT activity
Yamadate S., Sekiguchi M., Kawano K. 63

Vol. 160/2, 1986

Dependende of urine composition on the age and sex of healthy subjects
Hesse A., Classen A., Knoll M. et al. 79
Trends in urolithiasis in various ethnic groups and by age, in Israel
Zaidman J.L., Eidelman A., Pinto N. et al. 87
Strategies and methods for the analytical investigation of urinary calculi
Uldall A. 93
2,8-Dihydroxyadenine lithiasis
Simmonds H.A. 103
Medical investigation and treatment of urinary stones: A search for new ideas
Rose G.A. 109
A stable liquid control material for urinary enzyme assays
Maruhn D., Wehling K., Metz U. 117
Fluorometric measurement of urinary N-acetyl-β-D-glucosaminidase and its correlation to uremia
Linko-Lopponen S. 119
Identification of inhibitors of urinary alanine aminopeptidase
Mattenheimer H., Frolke W., Grotsch H., Simane Z. 129
Urinary enzyme analysis in renal allograft transplantation
Kotanko P., Keiler R., Knabl L. et al. 137
Serum and urinary enzyme activities in renal artery embolism
Donadio C., Auner I., Giordani R. et al. 145
Urinary enzyme excretion after a single dose of phenacetin and paracetamol (acetaminophen) during antidiuresis and during water diuresis
Metz U., Graben N., Maruhn D., Bock K.D. 151
Increased levels of urinary adenosine deaminase binding protein in children treated with cisplatin or methotrexate
Goren M.P., Wright R.K., Horowitz M.E. 157
Value of enzyme determinations in urine for the diagnosis of nephrotoxicity in rats
Hofmeister R., Bhargava A.S., Gunzel P. 163
Kidney-derived urinary antigens assayed with monoclonal antibodies for the detection of renal damage
Falkenberg F.W., Mai U., Puppe C. et al. 171
Biological monitoring of aluminium in renal patients
De Wolff F.A., Van der Voet G.B. 183
Allopurinol in renal failure and the tumour lysis syndrome
Simmonds H.A., Cameron J.S., Morris G.S., Davies P.M. 189
α(2U)-Globulin: Measurement in rat kidney and relationship to hyaline droplets
Stonard M.D., Phillips P.G.N., Foster J.R. et al. 197
NADH-dependent prostaglandin E_2-9-ketoreductase activity and prostaglandin synthesis in the Brattleboro rat kidney: Effects of the antidiuretic hormone
Rathaus M., Werber M.M., Podjarny E. et al. 205
Biochemical changes in liver, kidney and blood associated with common bile duct ligation
Israeli B.-A., Bogin E. 211

Vol. 160/3, 1986

In vitro studies upon the release of γ-glutamyltransferase from human liver
Wenham P.R., Horn D.B., Smith A.F. 223

The physicochemical properties and the process of pyrogel formation of IgD pyroglobulin
Ohno T., Furukawa H., Kanoh T., Uchino H. 235
Hydrolysis of conjugated steroids by β-glucuronidase from Ampullaria and application to the determination of urinary 17-hydroxycorticosteroids
Tsukada T., Isoe M., Yoshino M. 245
Quantitative analysis of the mitochondrial cytochrome P-450-linked monooxygenase system: NADPH-hepatoredoxin reductase, hepatoredoxin, and cytochrome P-450(S27) in livers of patients with cerebrotendinous xanthomatosis
Miki H., Takeuchi H., Yamada A. et al. 255
Renal effects of dichloroacetate in vivo
Kodama H., Yamaguchi S., Okabe I. et al. 265
A rapid quantitative capillary tube enzyme immunoassay for human chorionic gonadotropin in urine
Nagainis P.A., Nakagawa C.H., Baron S.L. et al. 273
Calcium activated neutral proteases (milli- and micro-CANP) and endogenous CANP inhibitor of muscle in Duchenne muscular dystrophy (DMD)
Reddy P.A., Anandavalli T.E., Anandaraj M.P.J.S. 281
Plasma fibronectin in liver cirrhosis and its diagnostic value
Gabrielli G.B., Casaril M., Bonazzi L. et al. 289
Mitochondrial aspartate aminotransferase linked to immunoglobulin G of the κ-λ type: report of a case
Moriyama T., Ashiie T., Kikuiri K. et al. 297
Degradation of vitamin B_6 standard solutions
Shephard G.S., Labadarios D. 307
Diagnostic value of an immunoenzymetric assay compared with an enzymic immunoinhibition procedure for creatine kinase isoenzyme MB
Moerman J., Nellens P., Beckers F., Sierens L. 313
Stability of creatinine, urea and uric acid in urine stored under various conditions
Soliman S.A., Abdel-Hay M.H., Sulaiman M.I., Tayeb O.S. 319

Vol. 161/1, 1986

Determination of serum neuron-specific enolase by differential immunocapture
Viallard J.-L., Ven Murthy M.R., Betail G., Dastugue B. 1
Measurement of 5-aminolaevulinic acid by reversed phase HPLC and fluorescence detection
Minder E.I. 11
Impaired hepatocellular integrity during general anaesthesia, as assessed by measurement of plasma glutathione S-transferase
Hussey A.J., Howie J., Allan L.G. et al. 19
Assay of blood acetaldehyde by HPLC with fluorescence detection of its 2-diphenylacetyl-1,3-indandione-1-azine derivative
Rideout J.M., Lim C.K., Peters T.J. 29
Serum levels of immunoreactive PSTI in acute abdominal disorders, with special reference to a possible extrapancreatic PSTI production
Lasson A., Borgstrom A., Ohlsson K. 37
Comparison of radioimmunoassay and immunoradiometric assay for serum prostatic acid phosphatase
Sturgeon C.M., Hussey A.J., Beynon L. et al. 47
Human inter-α trypsin inhibitor and immunologically related inhibitors investigated by quantitative immunoelectrophoresis. I. Method and reference material
Byrjalsen I., Hansen-Nord G., Odum L. 59
Hepatic damage in the rat following administration of thyroxine or triiodothyronine, assessed by measurement of plasma glutathione S-transferase YaYa concentrations
Beckett G.J., Hunter J.E., Hayes J.D. 69

Very long chain fatty acids in genetic peroxisomal disease fibroblasts: Differences between the cerebro-hepato-renal (Zellweger) syndrome and adrenoleukodystrophy variants
Molzer B., Korschinsky M., Bernheimer H. et al. 81
Effect of ileal exclusion on kinetics of very low density lipoprotein triglycerides in familial hypercholesterolemia
Koivisto P.V.I., Miettinen T.A. 91
The effect of lead on tetrahydrobiopterin metabolism. A possible mechanism for neurotoxicity
Eggar C., Hamon C.G.B., Morar C. et al. 103
Determination of L-asparaginase activity in serum by thin-layer chromatography: Application to the treatment of acute lymphoblastic leukemia
Clausen N., Christensen E. 111

Vol. 161/2, 1986

Multiple forms of acid and alkaline phosphatases: Genetics, expression and tissue-specific modification
Moss D.W. 123
Serum type III procollagen peptide as a non-invasive marker of liver damage during infancy and childhood in extrahepatic biliary atresia, idiopathic hepatitis of infancy and α1 antitrypsin deficiency
Trivedi P., Cheeseman P., Portmann B., Mowat A.P. 137
Non-specific binding of insulin in an equilibrium binding assay of insulin antibodies
Storms F.E.M.G., Lutterman J.A., Ross A.H. et al. 147
Epoxydicarboxylic aciduria resulting from the ingestion of castor oil
Hagenfeldt L., Blomquist L., Midtvedt T. 157
Apolipoprotein profile in healthy males and its relation to maximum aerobic capacity (MAC)
Berg A., Frey I., Keul J. 165
Methylmalonic aciduria and propionic acidaemia studied by proton nuclear magnetic resonance spectroscopy
Iles R.A., Chalmers R.A., Hind A.J. 173
Precise measurement of glycated serum albumin by column affinity chromatography and immunoturbidimetry
Reed P., Bhatnagar D., Dhar H., Winocour P.H. 191
Characteristics of ferritins in human milk secretions: Similarities to serum and tissue isoferritins
Arosio P., Ponzone A., Ferrero R. et al. 201
Assay of human pancreatic lipase in biological fluids using a non-competitive enzyme immunoassay
Carrere J., Galabert C., Thouvenot J.P., Figarella C. 209
Rapid diagnosis of Zellweger syndrome and infantile Refsum's disease by fast atom bombardment-mass spectrometry of urine bile salts
Lawson A.M., Madigan M.J., Shortland D., Clayton P.T. 221
Effect of fenoprofen on thyroid function tests
Tillman J., Leask J.T.S., Campbell J. et al. 233
Autoantibodies to thyroid hormones: Association with falsely low hormone levels measured by 'Amerlex' assay
Heyma P., Harrison L.C. 239
The influence of freezing and thawing on platelet-associated vasopressin during osmotic stimulation of the neurohypophysis
Baylis P.H., Thompson C.J., Burd J.M. 243

Vol. 161/3, 1986

Serum concentrations of laminin and aminoterminal propeptide of type III procollagen in relation to the portal venous pressure of fibrotic liver diseases
Gressner A.M., Tittor W., Negwer A., Pick-Kober K.-H. 249

Cerebrospinal fluid lactate dehydrogenase activities in patients with central nervous system metastases
Van Zanten A.P., Twijnstra A., Hart A.A.M., Ongerboer De Visser B.W. 259
Substrate specificity of alkaline phosphatase from human polymorphonuclear leukocytes
Stinson R.A., Gainer A.L., Chai J., Chan J.A. 283
Effect of parathyroid hormone and uremia on erythrocyte deformability
Bogin E., Earon Y., Blum M. 293
Membrane lipid peroxidation in erythrocytes of the newborn
Jain S.K. 301
Lysosomal enzyme activities in fresh frozen chorionic villi and in cultured trophoblasts
Ben-Yoseph Y., Evans M.I., Bottoms S.F. et al. 307
Quantitation of serum lactate dehydrogenase-5 with monoclonal antibodies
Vaidya H.C., Dietzler D.N., Ladenson J.H. 315
Fatty acyl-CoA dehydrogenase enzymes in human skeletal muscle
Carroll J.E., McGuire B.S., Hall C.L. 327
A preliminary study of urinary transferrin as a marker for prostatic cancer
Fernandez C., Rifai N., Wenger A.S. et al. 335
Serum adenosine deaminase activity in the differentiation of respiratory diseases in children
Yasuhara A., Nakamura M., Shuto H., Kobayashi Y. 341
Discrimination of heterozygotes for phenylketonuria, persistent hyperphenylalaninemia and controls by phenylalanine loading
Blitzer M.G., Bailey-Wilson J.E., Shapira E. 347

Vol. 162/1, 1987

Carrier detection in Duchenne muscular dystrophy: A review of current issues and approaches
Gruemer H.-D., Prior T. 1
A novel alkaline phosphatase isozyme in human adipose tissue
Okochi T., Seike H., Saeki K. et al. 19
Spectrophotometric assay of xanthine oxidase with 2,2′-azino-di(3-ethylbenzthiazoline-6-sulphonate) (ABTS) as chromogen
Majkic-Singh N., Bogavac L., Kalimanovska V. et al. 29
Atypical LD isoenzymes with five diffuse bands and evidence for two types of binding
Chappell D.J., Buttery J.E., Pannall P.R. 37
Quantitation of serum bile acids by isotope dilution with ^{13}C-labelled homologs
Stellaard F., Paumgartner G. 45
Competitive inhibitor binding assay (CIBA) of captopril and other ACE inhibitors
Gronhagen-Riska C., Tikkanen I., Fyhrquist F. 53
Purification of porphobilinogen deaminase from human erythrocytes by fast protein liquid chromatography
De Rooij F.W.M., Hamer C.M., Wilson J.H.P. 61
Development of carboxypeptidase K (creatine kinase conversion factor) in man
Edwards R.J., Watts D.C. 69
Extended application of RP-HPLC for characterization of human hemoglobin variants
Baudin V., Wajcman H. 75
Enzyme-linked immunosorbent assay of serum pepsinogen I
Huang S.C., Miki K., Hirano K. et al. 85
Effect of the partial beta-agonist prenalterol on plasma renin activity in patients with left ventricular failure
Wathen C.G., Mackay I.G., Glover D.R., Roulston J.E. 97
Determination of neuron specific enolase in amniotic fluid and maternal serum for the prenatal diagnosis of fetal neural tube defects
Sorensen K., Brodbeck U., Norgaard-Pedersen B. 101
Identification and elimination of an ethylene glycol determination artifact
Smith N.B. 105

Vol. 162/2, 1987

Biosensor for lactate determination in biological fluids. I. Construction and properties of the biosensor
Racek J., Musil J. 129
Are 'precipitated LDL' really low density lipoproteins?
Mainard F., Madec Y. 141
Diagnostic sensitivity of fasting and postprandial serum bile acids determined by different methods
Mannes G.A., Stellaard F., Paumgartner G. 147
Serum lipids in hyper- and hypothyroidism before and after treatment
Friis T., Pedersen L.R. 155
The effect of assay conditions on the measurement of anti-elastase function in lung secretions
Morrison H.M., Kramps J.A., Afford S.C. et al. 165
Development and validation of an improved radioimmunoassay for serotonin in platelet-rich plasma
Gow I.F., Corrie J.E.T., Williams B.C., Edwards C.R.W. 175
Human inter-α-trypsin inhibitor and immunologically related inhibitors investigated by quantitative immunoelectrophoresis. II. Pathological conditions
Odum L., Hansen-Nord G., Byrjalsen I. 189
An enzyme-linked immunosorbent assay (ELISA) for plasma progesterone: Immobilised antigen approach
Elder P.A., Yeo K.H.J., Lewis J.G., Clifford J.K. 199
Interference in the measurement of red cell glycogen by glycogen from white cells
Collins W., Brown I., Taylor W.H. 207
Human erythrocyte transketolase: No evidence for variants
Kaufmann A., Uhlhaas S., Friedl W., Propping P. 215
The effect of combined fenofibrate and cholestyramine therapy on low-density lipoprotein kinetics in familial hypercholesterolemia patients
Malmendier C.L., Delcroix C., Lontie J.-F. 221
Human growth hormone (hGH) in urine and its correlation to serum hGH examined by a highly sensitive sandwich enzyme immunoassay
Hashida S., Ishikawa E., Kato Y. et al. 229
Inhibition by 5-sulphosalicylic acid of the glutathione reductase recycling assay for glutathione analysis
Allen K.G.D., Arthur J.R. 237

Vol. 162/3, 1987

An evaluation of the low-pH enzymatic assay of urinary D-glucaric acid, and its use as a marker of enzyme induction in exocrine pancreatic disease
Sandle L.N., Braganza J.M. 245
On the homogeneity of stools with respect to bile acid composition and normal day-to-day variations: A detailed qualitative and quantitative study using capillary column gas chromatography-mass spectrometry
Setchell K.D.R., Ives J.A., Cashmore G.C., Lawson A.M. 257
A method for measuring testosterone-estradiol-binding globulin - new analysis of experimental data and comparison with two commercially available kits
Lecureuil N., Trochet J., Crouzat-Reynes G. et al. 277
The fat meal test of malabsorption: Some further experience
Steen G., Holthuis N. 291
Deficient cholesterol side chain oxidation in patients without peroxisomes (Zellweger syndrome): Evidence for the involvement of peroxisomes in bile acid synthesis in man
Wanders R.J.A., Schutgens R.B.H., Heymans H.S.A. 295
Development of a method to monitor low molecular mass hydrocarbons in exhaled breath of man: Preliminary evaluation of its interest for detecting a lipoperoxidation process in vivo
Hotz P., Hoet P., Lauwerys R., Buchet J.-P. 303

Lectin-affinity chromatography of serum gamma-glutamyltransferase in liver disease
Delanghe J.R., De Buyzere M.L., De Scheerder I.K. et al. 311
Human erythrocyte γ-glutamyltransferase in liver diseases
Daniel D.S., Ramakrishna B.S., Balasubramanian K.A. 319
Uromucoid (Tamm-Horsfall glycoprotein) forms different polymeric arrangements on a filter surface under diffferent physicochemical conditions
Wiggins R.C. 329
Radioimmunoassay of glandular kallikrein in human plasma after partial purification by immunoaffinity column
Nishimura K., Inoue Y., Kokubu T. 341

Vol. 163/1, 1987

Coeliac disease: A critical review of aetiology and pathogenesis
Davidson A.G.F., Bridges M.A. 1
Enzyme-linked immunosorbent assay (ELISA) for IgG antibodies to thyroglobulin
Shimojo N., Kohno Y., Tarutani O. et al. 41
Urinary protein/creatinine ratio as an indicator of allograft function following live related donor renal transplantation
Krishna K.S., Pandey A.P., Kirubakaran M.G., Kanagasabapathy A.S. 51
Enzyme linked immunosorbent assay for the measurement of nonenzymatically glucosylated proteins in serum and in tissues
Go R., Quinn T.J., Gonen B. 63
Total and free disopyramide by fluorescence polarization immunoassay and relationship between free fraction and alpha-1 acid glycoprotein
Chen B.H., Taylor E.H., Pappas A.A. 75
Determination of trimethoprim in pediatric samples by high-performance liquid chromatography
Hoppu K., Arjomaa P. 81
Binding to cellulose of IgG from patients with interstitial lung diseases
Dewair M., Baur X. 87
Diagnostic significance of urinary enzymes: Development of a high performance liquid chromatographic method for the measurement of urinary lysozyme
Severini G., Aliberti L.M. 97
Enzyme immunoassay of total insulin in human serum containing anti-insulin antibodies
Kohno T., Ishikawa E., Sugiyama S. et al. 105
Apolipoprotein and lipoprotein cholesterol concentrations in trained and sedentary males
Rifai N., King M.E., De Meersman R. et al. 113

Vol. 163/2, 1987

The relative extent of glycation of haemoglobin and albumin
Olufemi S., Talwar D., Robb D.A. 125
The effect of 2,4-dihydroxypyridine-N-oxide, a new orally active iron chelator, on iron excretion in mice
Kontoghiorghes G.J. 137
Recognition of cervical neoplasia by the estimation of a free-radical reaction product (octadeca-9,11-dienoic acid) in biopsy material
Griffin J.F.A., Wickens D.G., Tay S.K. et al. 143
Recognition of cervical neoplasia by the estimation of a free-radical reaction product (octadeca-9,11-dienoic acid) in exfoliated cells
Tay S.K., Singer A., Griffin J.F.A. et al. 149
Evaluation of gravimetric assays of the H_2O concentration in human serum and urine
De Jong Th. G.M., Huizenga J.R., Gips C.H. 153

The increase in zincaemia provoked by starvation does not influence the intestinal absorption of zinc
Couturier E., Praet J.P., Vis H.L., Devroede P. 165
Effect of temperature and sample preparation on performance of ion-moderated partition chromatography of organic acids in biological fluids
Rumsby G., Belloque J., Ersser R.S., Seakins J.W.T. 171
Microheterogeneity forms of alpha$_1$-acid glycoprotein as indicators of rheumatoid arthritis activity
Mackiewicz A., Pawlowski T., Mackiewicz-Pawlowska A. et al. 185
Serum pepsinogens I and II and stomach cancer
Stemmermann G.N., Samloff I.M., Nomura A.M.Y., Heilbrun L.K. 191
Preparative fractionation of amyloid proteins on a microgram scale by high-performance liquid chromatography and polyacrylamide gel electrophoresis
Kaplan B., Pras M. 199
A simple immunotechnique for the determination of serum concentration of apolipoprotein E
Rifai N., Silverman L.M. 207
Stability of enzymes of lysosomal origin in human cerebrospinal fluid
Goi G., Fabi A., Lombardo A. et al. 215
m-Aminophenylboronate affinity ligands distinguish between nonenzymically glycosylated proteins and glycoproteins
Gould B.J., Hall P.M. 225
Electrophoretic differentiation of acquired angioedema from hereditary angioedema
Papadopoulos N.M., Frank M.M. 231

Vol. 163/3, 1987

ELISA for determination of albumin in the nanogram range: Assay in cerebrospinal fluid and comparison with radial immunodiffusion
Schwerer B., Bach M., Bernheimer H. 237
Quantification of human apolipoprotein E in plasma and lipoprotein subfractions by a non-competitive enzyme immunoassay
Koffigan M., Kora I., Clavey V. et al. 245
A non-competitive enzyme immunoassay of human α-lactalbumin in biological fluids and tissue extracts
Simickova M., Lang B.A., Semotan K., Kocent A. 257
Phosphatidylethanolamine methyltransferase and cAMP, cGMP phosphodiesterases in lymphocytes and monocytes in sarcoidosis
Pacheco Y., Dubois M., Prigent A.F. et al. 267
Partial characterization of human ileal alkaline phosphatase: Differences between human ileal and duodenal enzymes
Miura M., Matsuzaki H., Sakagishi Y., Komoda T. 279
Evidence for endotoxin binding capacity of human Gc-globulin and transferrin
Berger D., Beger H.G. 289
Human hypoxanthine-guanine phosphoribosyltransferase. Development of a spectrophotometric assay and its use in detection and characterization of mutant forms
Keough D.T., McConachie L.A., Gordon R.B. et al. 301
Direct assay for testosterone in saliva: Relationship with a direct serum free testosterone assay
Johnson S.G., Joplin G.F., Burrin J.M. 309
Protein transfer at the blood cerebrospinal fluid barrier and the quantitation of the humoral immune response within the central nervous system
Reiber H., Felgenhauer K. 319
Measurement by bioluminescence technique of erythrocyte membrane Na^+,K^+-ATPase activity in hypertensive patients
Lechi C., Corradini P., Polignano R. et al. 329
Detection of desialylated forms of human chorionic gonadotropin
Imamura S., Armstrong E.G., Birken S. et al. 339

Vol. 164/1, 1987

Evaluation of iron-deficiency anemia by an iron supplementation trial in children living at a 2800-m altitude
Estrella R., Hercberg S., Maggy G. et al. 1
Urinary thiosulphate as an indicator of exposure to hydrogen sulphide vapour
Kangas J., Savolainen H. 7
A circulating complex between ASAT and IgG in serum in an apparently healthy woman
Fex G., Berntorp K. 11
24-h Catecholamine excretion: Relationships with age and weight
Jenner D.A., Harrison G.A., Prior I.A.M. et al. 17
Immunological comparison of biotinidase in serum from normal and biotinidase-deficient individuals
Wolf B., Miller J.B., Hymes J. et al. 27
Systemic membrane defect and the inhibition of lymphocyte capping in Duchenne Muscular Dystrophy
Goldsmith B.M., Gruemer H.-D. 33
Determination of creatine kinase isozymes in sera and tissues of patients with various lung carcinomas
Usui A., Fujita K., Imaizumi M. et al. 47
Human pheochromocytoma phenol sulfotransferase: Biochemical properties and activities of thermolabile and thermostable forms
Sinsheimer E.G., Anderson R.J. 55
Enzyme-linked immunosorbent assay (ELISA) for steroid hormones with polyclonal and monoclonal antibodies: An assay for urinary aldosterone
Hanquez C., Urios P., Desfosses B. et al. 71
Physical state abnormality of erythrocyte membrane in myotonic dystrophy: A spin label study
Thomas M.A., Lamas E., Daveloose D. et al. 83
Free and Apo B-associated Lpa-specific protein in human serum
Gries A., Nimpf J., Nimpf M. et al. 93
Effects of exogenous β-galactosidase on cultured fibroblasts with β-galactosidase deficiency
Nakao Y., Kozutsumi Y., Fukui S. et al. 101

Vol. 164/2, 1987

Liver-specific F antigen in the serum of patients with liver diseases and the detection of early stage hepatocellular carcinoma
Mori Y., Iwama S., Mori T. et al. 127
Relationships between the lipid content and the rheological properties of airway secretions in cystic fibrosis
Galabert C., Jacquot J., Zahm J.M., Puchelle E. 139
Relationship between albumin and fructosamine concentration in diabetic and non-diabetic sera
Johnson R.N., Metcalf P.A., Baker J.R. 151
Plasma amino-acids analysis: Effects of delayed samples preparation and of storage
Schaefer A., Piquard F., Haberey P. 163
A simple chromogenic assay for arylsulfatase A
Lee-Vaupel M., Conzelmann E. 171
Determination of erythrocyte glucose 1,6-bisphosphate — A comparison of two methods using a centrifugal analyzer
Thorburn D.R., Kuchel P.W. 181
Improved between-laboratory agreement for specific protein assays in serum following introduction of a common reference preparation (SPS-01) demonstrated in an external quality assessment scheme
Chambers R.E., Bullock D.G., Whicher J.T. 189

Clinical application of enzyme immunoassay in the analysis of citrullinemia
Imamura Y., Kobayashi K., Yamashita T. et al. 201
An alternative way of presenting quantitative amino acid results
Briddon A., Oberholzer V.G. 209
Removal of ascorbate from urine prior to assaying with a commercial oxalate kit
Kasidas G.P., Rose G.A. 215
Serum angiotensin converting enzyme activity in ex-smokers
Ninomiya Y., Kioi S., Arakawa M. 223
An evaluation of four methods for the detection of heterozygous cystinuria
Giugliani R., Ferrari I., Greene L.J. 227
A comparison of the utility of Dowex resin and polybenzimidazole Aurorez resin in the determination of urinary porphobilinogen concentrations
Galbraith R.A., Sassa S., Kappas A. 235

Vol. 164/3, 1987

Lipid peroxidation and malondialdehyde in the synovial fluid and plasma of patients with rheumatoid arthritis
Wade C.R., Jackson P.G., Highton J., Van Rij A.M. 245
The molecular abnormality of albumin Niigata: 269 Asp → Gly
Sugita O., Endo N., Yamada T. et al. 251
Identification of glutarylcarnitine in glutaric aciduria type 1 by carboxylic acid analyzer with an ODS reverse-phase column
Kidouchi K., Sugiyama N., Morishita H. et al. 261
Modified phosphoribosylpyrophosphate (PRPP) radioenzymatic assay: Increased sensitivity, technical simplification and new applications
Ghitis J., Waxman S. 267
5'-Nucleotide phosphodiesterase isoenzymes in human serum: Quantitative measurement and some biochemical properties
Luthje J., Ogilvie A. 275
Rapid suppression of plasma alkaline phosphatase activity after renal transplantation in patients with osteodystrophy
Cundy T., Kanis J.A. 285
Purification of placental protein PP12 from human amniotic fluid and its comparison with PP12 from placenta by immunological, physicochemical and somatomedin-binding properties
Koistinen R., Huhtala M.-L., Stenman U.-H., Seppala M. 293
A rapid and precise assay for calmodulin and cyclic AMP phosphodiesterase on a centrifugal analyser
Walker S.W., Milner S.W., Howie A.F. 305
Effects of purified fish oil on platelet lipids and function in diabetic women
Tilvis R.S., Rasi V., Viinikka L. et al. 315
Reduced plasma and white blood cell selenium levels in haemodialysis patients
Foote J.W., Hinks L.J., Lloyd B. 323
The effect of fatty acid chain length on the rate of arylester hydrolysis by various albumins
Wolfbeis O.S., Gurakar A. 329
Isoenzymes of urinary N-acetyl-β-D-glucosaminidase (NAG) in patients with renal transplants
Yuen C.-T., Corbett C.R.R., Kind P.R.N. et al. 339

Vol. 165/1, 1987

Animal models of inherited hematologic disease
Kaneko J.J. 1
Apoenzyme content of serum aminotransferases in patients with a renal allograft treated with

cyclosporine A and azathioprine
Hafkenscheid J.C.M., Rosier J.G.M.C., Van Dijk C.M.C.E. 21
Fatty acid composition of phospholipids in bile in man: promoting effect of deoxycholate on arachidonate
Van Berge Henegouwen G.P., Van der Werf S.D.J., Ruben A.T. 27
A new simple screening method for the diagnosis of medium chain acyl-CoA dehydrogenase deficiency
Bhuiyan A.K.M.J., Watmough N.J., Turnbull D.M. et al. 39
The direct barbituric acid assay for nicotine metabolites in urine: A simple colorimetric test for the routine assessment of smoking status and cigarette smoke intake
Barlow R.D., Stone R.B., Wald N.J., Puhakainen E.V.J. 45
Correlation between the cystic fibrosis protein detected by isoelectric focusing and a protein of 12 kDa detected by SDS-polyacrylamide gel electrophoresis
Barthe C., Figarella C., Galabert C., Guy-Crotte O. 53
Thermal desorption — gas chromatography of plasma isovaleraldehyde in hepatic encephalopathy in man
Al Mardini H., O'Brien C.J., Bartlett K. et al. 61
Conditions for a reliable application of the lysoplate method in the determination of lysozyme
Gupta D.K., Von Figura K., Hasilik A. 73
Value of glial fibrillary acidic protein determination in amniotic fluid for prenatal diagnosis of neural tube defects
Van Regemorter N., Gheuens J., Noppe M. et al. 83
Comparison of fat agglutination slide test and latex test for C-reactive protein
Hulman G., Fuller M. 89
Approved IFCC methods. Reference method (1986) for pH measurement in blood
Maas A.H.J., Weisberg H.F., Burnett R.W. et al. 97
Approved recommendation (1986) on the theory of reference values. Part 1. The concept of reference values
Solberg H.E. 111

Vol. 165/2–3, 1987

Thyroid function in leukaemia patients after allogeneic bone marrow transplantation
Frolich M., De Planque M.M., Goslings B.M., Meinders A.E. 127
Interference of blood clotting factors in the determination of HDL_2- and HDL_3-cholesterol by the dual precipitation method
Demacker P.N.M., Hak-Lemmers H.L.M., Van Heijst P.J. 133
Human serum sialo transferrins in diseases
Van Eijk H.G., Van Noort W.L., De Jong G., Koster J.F. 141
Determination of polyamines and metabolites in cerebrospinal fluid by isotope dilution mass fragmentography, and a clinical application
Van Den Berg G.A., Schaaf J.M., Nagel G.T. et al. 147
Does the degree of polymerization of haptoglobin influence the results of the radial immunodiffusion assay?
Van Rijn H.J.M., Wisman J.P.W., Gmelig Meyling F. 155
Catalytic and immunologic activities of placental-like alkaline phosphatase in clinical studies. The value of PLAP in follow-up of ovarian cancer
Haije W.G., Van Driel J., Van Der Burg M.E.L. 165
Therapeutic drug monitoring of amitriptyline: impact of age, smoking and contraceptives on drug and metabolite levels in bulimic women
Edelbroek P.M., Zitman F.G., Knoppert-Van Der Klein E.A.M., et al. 177
Assay of oestrogen and progestin receptors in human meningioma cytosols using immunological methods
Blankenstein M.A., Van Der Meulen-Dijk C., Thijssen J.H.H. 189

Urinary excretion of 2-methyl-2,3-butanediol and 2,3-pentanediol in patients with disorders of propionate and methylmalonate metabolism
Duran M., Ketting D., Bruinvis L. et al. 197
Pregnancy and guinea-pig isotransferrins — isolation and characterization of both isotransferrins
Bierings M.B., Heeren J.W.A., Van Noort W.L. et al. 205
Urinary polyamines and their metabolites during new combination chemotherapeutic treatments of high grade non-Hodgkin lymphoma
Muskiet F.A.J., Van Imhoff G.W., Van Den Berg G.A. et al. 213
Laser diffraction ellipsometry of erythrocytes under controlled shear stress using a rotational viscosimeter
Hardeman M.R., Goedhart P., Breederveld D. 227
Influence of gastric emptying on the PABA test
Hoek F.J., Sanders G.T.B., Tytgat G.N.J. 235
Reinvestigations into the formation and assay of C3bBbP complexes
Asghar S.S., Barendsen C.Y., Van Der Helm H.J. 243
Glucagon and insulin response to meals in non-obese and obese Dutch women
Hill P., Garbaczewski L., Koppeschaar H. et al. 253
Analytical and clinical evaluation of commercial protein C assays
Sturk A., Morrien-Salomons W.M., Huisman M.V. et al. 263
Differential diagnosis of genetic disease by DNA restriction fragment length polymorphisms
Bolhuis P.A., Defesche J.C., Van Der Helm H.J. 271
ELISA of ceruloplasmin and alpha-2-macroglobulin in paired bronchoalveolar lavage fluid and serum samples
Out T.A., Jansen H.M., Van Steenwijk R.P. et al. 277
Immunoglobulin D in cerebrospinal fluid
Out T.A., Hische E.A., Van Walbeek H.K., Van Der Helm H.J. 289
Quantification of cholesterol nucleation promoting activity in human gallbladder bile
Drapers J.A.G., Groen A.K., Stout J.P.J. et al. 295
Prenatal diagnosis of Zellweger syndrome by measurement of very long chain fatty acid (C26:0) β-oxidation in cultured chorionic villous fibroblasts: implications for early diagnosis of other peroxisomal disorders
Wanders R.J.A., Van Wijland M.J.A., Van Roermund C.W.T. et al. 303
Alanine glyoxylate aminotransferase and the urinary excretion of oxalate and glycollate in hyperoxaluria type I and the Zellweger syndrome
Wanders R.J.A., Van Roermund C.W.T., Westra R. et al. 311
X-linked adrenoleukodystrophy: defective peroxisomal oxidation of very long chain fatty acids but not of very long chain fatty acyl-CoA esters
Wanders R.J.A., Van Roermund C.W.T., Van Wijland M.J.A. et al. 321
A sensitive radioimmunoassay of atrial natriuretic peptide in human plasma, using a tracer with an immobilized glycouril agent
Rosmalen F.M.A., Tan A.C.I.T.L., Tan H.S., Benraad T.J. 331
A symmetric dialysis method for the determination of free testosterone in human plasma
Swinkels L.M.J.W., Ross H.A., Benraad T.J. 341
Determination of urinary cystine for monitoring anticystinuric therapy
Backer E.T., Van Den Ende A., Geus W.P. 351
Effect of ritodrine on thyroid hormone concentrations
Menheere P.P.C.A., Essed G.G.M., Sels J.P.J.E. 359
β-Aminoisobutyric acid as a marker of thymine catabolism in malignancy
Van Gennip A.H., Van Bree-Blom E.J., Abeling N.G.G.M. et al. 365
The effect of oral contraceptives on plasma-free and salivary cortisol and cortisone
Meulenberg P.M.M., Ross H.A., Swinkels L.M.J.W., Benraad T.J. 379

Vol. 166/1, 1987

Automated enzymatic standardized lipid analyses for plasma and lipoprotein fractions
McNamara J.R., Schaefer E.J. 1

Application of transformation algorithm and nonparametric calculation in determining the reference intervals of some urine constituents and characteristics
Soliman S.A., Abdel-Hay M.H., Tayeb O.S., Sulaiman M.I. 9
Immunochemical determination of acetylcholinesterase in amniotic fluid — An evaluation of eleven monoclonal antibodies
Rasmussen A.G., Sorensen K., Selmer J. et al. 17
Agarose gel isoelectrofocusing of UDP-galactose pyrophosphorylase and galactose-1-phosphate uridyltransferase. Developmental aspect of UDP-galactose pyrophosphorylase
Shin Y.S., Niedermeier H.-P., Endres W. et al. 27
Atrial natriuretic peptide in human plasma — Comparison of radioreceptor versus radioimmunoassay
Sagnella G.A., Buckley M.G., Markandu N.D., MacGregor G.A. 37
Age-related accumulation of phytanic acid in plasma from patients with the cerebro-hepato-renal (Zellweger) syndrome
Wanders R.J.A., Smit W., Heymans H.S.A. et al. 45
Serum human pancreatic elastase I levels in pancreatic disease
Buamah P.K., Cornell C., Cassells-Smith A.J., Skillen A.W. 57
Detection of photobilirubin in urine of jaundiced infants supporting the diagnosis of 'Bronze Baby Syndrome'
Meisel P., Jahrig D., Meisel M. 61
A simple latex agglutination test for urinary albumin screening
Paoli C., Bardelli F., Tarli P. et al. 67
Isoelectric focusing of the cytosolic human creatine kinase isoenzymes
Nealon D.A. 73
Serum aromatic L-amino acid decarboxylase activity in patients with acute hepatitis
Nishigaki I., Nagatsu T. 79
Turbidimetric measurement of microalbuminuria
Harmoinen A., Vuorinen P., Jokela H. 85
Determination of aromatic and neutral aminoacids by HPLC in blood specimens collected on filter paper
Schrynemackers-Pitance P., Schoos-Barbette S. 91

Vol. 166/2–3, 1987

A comparison of dopamine and homovanillic acid excretion, as prognostic indicators in malignant phaeochromocytoma
Tippett P.A., West R.S., McEwan A.J. et al. 123
Comparison of N-acetyl-β-glucosaminidase and albuminuria with clinical finding of microangiopathy in type I diabetes mellitus
Skrha J., Perusicova J., Stolba P. et al. 135
Evidence for an increased glycation of IgG in diabetic patients
Danze P.-M., Tarjoman A., Rousseaux J. et al. 143
Tubular maximum for calcium reabsorption: Lack of diagnostic usefulness in primary hyperparathyroidism and familial hypocalciuric hypercalcaemia
Kent G.N., Bhagat C.I., Garcia-Webb P., Gutteridge D.H. 155
A simple enzymatic fluorimetric method for the determination of branched-chain L-amino acids in microlitre volumes of plasma
Gleeson M., Maughan R.J. 163
Blood and liver selenium concentrations in patients with liver diseases
Valimaki M., Alfthan G., Pikkarainen J. et al. 171
Studies of the sialylation and microheterogeneity of human serum α_1-acid glycoprotein in health and disease
Moule S.K., Peak M., Thompson S., Turner G.A. 177
Comparison of the precision of seven analytical methods for the H_2O concentration in human serum and urine
De Jong Th. G.M., Huizenga J.R., Wolthers B.G. et al. 187

Separation of plasma carotenoids and quantitation of β-carotene using HPLC
Gatautis V.J., Pearson K.H. 195
Purification and characterization of glutathione S-transferases in human kidney
Tateoka N., Tsuchida S., Soma Y., Sato K. 207
A fluorometric micromethod for estimation of carnosinase in dried blood samples
Orfanos A.P., Guthrie R., Jinks D.C. 219
Estimation and characterization of glycosylated carbonic anhydrase I in erythrocytes from patients with diabetes mellitus
Kondo T., Murakami K., Ohtsuka Y. et al. 227
Sandwich enzyme immunoassay for human interleukin-1β (hIL-1β) in urine
Tanaka K., Ishikawa E., Ohmoto Y., Hirai Y. 237
Detection of microalbuminuria in diabetic patients using a simple latex agglutination test
Spooner R.J., Weir R.J., Frier B.M. 247
Peroxisomal very long-chain fatty acid β-oxidation in human skin fibroblasts: Activity in Zellweger syndrome and other peroxisomal disorders
Wanders R.J.A., Van Roermund C.W.T., Van Wijland M.J.A. et al. 255
Specific assays for human alkaline phosphatase isozymes
Hirano K., Matsumoto H., Tanaka T. et al. 265
Improved method for determination of high density lipoprotein cholesterol using a sensitive reagent and a centrifugal analyzer
Moshides J.S. 275
Diode-array spectrophotometry for simultaneous measurement of hemoglobin pigments
Fogh-Andersen N., Siggaard-Andersen O., Lundsgaard F.C., Wimberley P.D. 283
Spectrophotometric determination of hemoglobin pigments in neonatal blood
Fogh-Andersen N., Siggaard-Andersen O., Lundsgaard F.C., Wimberley P.D. 291
Estimation of whole body bone resorption rate: A comparison of urinary total hydroxyproline excretion with two radioisotopic tracer methods in osteoporosis
Deacon A.C., Hulme P., Hesp R. et al. 297
Immunoglobulin light-chain immunoblots or urine proteins from patients with tubular and Bence-Jones proteinuria
Norden A.G.W., Fulcher L.M., Flynn F.V. 307
Hemoglobin A_1 and serum fructosamine levels in hyperthyrodism
Ford H.C., Lim W.C., Crooke M.J. 317
Detection of wheat gliadin contamination of gluten-free foods by a monoclonal antibody dot immunobinding assay
Freedman A.R., Galfre G., Gal E. et al. 323
Estimation of low levels of phosphatidylglycerol in aspirates of newborns using the AmnioStat FLM agglutination test
Egberts J., Contant J., Wijnands J.B.G. 329

Vol. 167/1, 1987

Rheumatoid factors
Dorner R.W., Alexander Jr. R.L., Moore T.L. 1
Direct analysis of tetrahydrobiopterin in cerebrospinal fluid by high-performance liquid chromatography with redox electrochemistry: Prevention of autoxidation during storage and analysis
Howells D.W., Hyland K. 23
The crypt-villus distribution of the 10000 M(r) intestinal calcium binding protein in human jejunum
Staun M. 31
Lack of relationship between beta-hexosaminidase activity and retinopathy in insulin dependent diabetics
Agardh C.-D., Agardh E., Hultberg B. 37
Immunochemical characterization and quantification of the sex steroid-binding protein (SBP) in human amniotic fluid
Auzelle M.-P., Tardivel-Lacombe J., Domingo M., Egloff M., Degrelle H. 43

Cholinesterase activities in resected bowel specimens from children with Hirschsprung's disease
Causse E., Vaysse P., Fabre J., Valdiguie P., Thouvenot J.P. 51
Relationship between effects of added albumin, initial free thyroxine value and endogenous serum-binding protein concentrations on Amerlex® free thyroxine estimations
Midgley J.E.M., Winton M.R.J., Wilkins T.A. 67
Guanidino compounds in serum and cerebrospinal fluid or non-dialyzed patients with renal insufficiency
De Deyn P.P., Marescau B., Cuykens J.J., Van Gorp L., Lowenthal A., De Potter W.P. 81
Underestimation of serum albumin by the bromcresol purple method and a major endogenous ligand in uremia
Mabuchi H., Nakahashi H. 89
Determination of total cholesterol in serum by flow injection analysis with immobilized enzymes
Fernandez-Romero J.M., Luque De Castro M.D., Valcarcel M. 97
A rapid method for determining catalase in human blood
Lamoureux G., Bourdeau S., Dubois G., Charbonneau R., Gagnon M., Grad B.R. 105

Vol. 167/2, 1987

Faecal lysozyme: Determination, reference intervals and some data in gastro-intestinal disease
Costongs G.M.P.J., Hemrika M.H., Engels L.G.J.B. et al. 125
CSF concentration and CSF/blood ratio of fuel related components in children after prolonged fasting
Lamers K.J.B., Doesburg W.H., Gabreels F.J.M. et al. 135
Isoenzyme specificities of immunoglobulins isolated from lactate dehydrogenase-immunoglobulin complexes
Kanemitsu F. 147
Development and application of an enzyme-linked immunosorbent assay for acid stable trypsin inhibitor (ASTI) in human plasma
Yoshida E., Sumi H., Maruyama M. et al. 155
Quantitative detection of a variant prealbumin associated with type 1 familial amyloidotic polyneuropathy (Japanese type) by high performance liquid chromatography
Ikegawa S., Tanase S., Araki S., Morino Y. 165
Evaluation of the average of patients: Application to endocrine assays
Douville P., Cembrowski G.S., Strauss J.F. 173
Intracellular magnesium of mononuclear cells from venous blood of clinically healthy subjects
Reinhart R.A., Marx Jr. J.J., Haas R.G., Desbiens N.A. 187
HDL cholesterol and the acute phase reaction following myocardial infarction and acute pancreatitis
Fahie-Wilson M., Mills R., Wilson K. 197
4-Nitrophenylalkaline phosphatase inhibition in muscular dystrophy
Goldemberg A.L., Sanchez J.J., Garcia A.M. et al. 211
A simple, reproducible and cheap batch method for the analysis of serum glycoproteins using a Sepharose-coupled lectins and silver staining
Thompson S., Latham J.A.E., Turner G.A. 217
An improved spot-test for the detection of paraquat and diquat in biological samples
Woollen B.H., Mahler J.D. 225
Analysis of 25-hydroxyvitamin D in plasma using solid phase extraction
Shephard G.S., Carlini S.M., Hanekom C., Labadarios D. 231

Vol. 167/3, 1987

Serum aldolase isoenzymes in benign and malignant liver disease
Royds J.A., Kennedy H.J., Little P.V., Taylor C.B., Triger D.R. 237

Effects of severe burns on glycan microheterogeneity of four acute phase proteins
Mallet B., Franc J.L., Miquel M., Arnaud C. 247
Inorganic phosphate transport across the red blood cell membrane: The effect of exposure to hyperoxia
Webster N.R., Toothill C. 259
Clinical usefulness of cimetidine for the treatment of acute intermittent porphyria — a preliminary report
Horie Y., Udagawa M., Hirayama C. 267
Vitamin D-binding protein synthesized by a human hepatocellular carcinoma cell line
Egawa T., Yamamoto H., Kishimoto S. 273
Glycylproline dipeptidyl aminopeptidase and γ-glutamyl transpeptidase in human hepatic cancer and embryonal tissues
Kojima J., Ueno Y., Kasugai H., Okuda S., Akedo H. 285
A large molecular form of glucagon-like peptide-1 (GLP-1) immunoreactivity is co-released with glucagon from pancreas by arginine in normal subjects
Hirota M., Shimizu I., Ohboshi C., Nishino T., Shima K. 293
An increased apo A-IV serum concentration of patients with chronic renal failure on hemodialysis
Seishima M., Muto Y. 303
A simple differential immunoprecipitation assay of urinary acid and neutral α-glucosidases for glycogenosis II
Tsuji A., Yang R.-C., Omura K., Imabayashi T., Suzuki Y. 313
Hydrolysis of phosphatidate by human placental alkaline phosphatase
Sumikawa K., Saeki K., Okochi T., Adachi K., Nishimura H. 321
Macromolecules inhibit calcium oxalate crystal growth and aggregation in whole human urine
Edyvane K.A., Hibberd C.M., Harnett R.M., Marshall V.R., Ryall R.L. 329
Competitive enzyme immunoassay for human serum apolipoproteins C1, C11 and C111
Fu M., Liu B., Liu W., Fang D. 339
Structurally abnormal transthyretin causing familial amyloidotic polyneuropathy in Sweden
Nakazato M., Steen L., Holmgren G., Kurihara T., Matsukura S., Kangawa K., Matsuo H. 341

Vol. 168/1, 1987

Detergent induced lysis of erythrocytes in kwashiorkor
Rao A., Onuora C.U., Cherian A. 1
Abnormal degradation of erythrocyte membrane proteins in hereditary spherocytosis
Gaczynska M., Bartosz G., Judkiewicz L., Rosin J. 7
Immunoradiometric assay of serum thyroid stimulating hormone (TSH) as a first-line test in a mixed hospital population
Ronnov-Jessen D., Skov L., Faber J. 13
Changes in magnesium, zinc and calcium in men and women after an acute myocardial infarction
Speich M., Gelot S., Robinet N., Arnaud P., Nicolas G. 19
Black-white differences in serum Lp(a) lipoprotein levels
Parra H.-J., Luyeye I., Bouramoue C., Demarquilly C., Fruchart J.-C. 27
Interference of an α_2 component in immunological determinations of pregnancy-specific β_1 glycoprotein in serum
Pala A., Di Ruzza A., Curtilli L., Strom R., Moro M., Reverberi L., Carenza L. 33
Evaluation of RU-27987 as a ligand to determine the progesterone receptor
Abendstein B., Marth C., Daxenbichler G. 47
Differential reactivity of α-fetoprotein with lectins and its usefulness in the diagnosis of various liver diseases
Buamah P.K., Cornell C., Cassells-Smith A.J., Skillen A.W. 69
Increased serum lactate dehydrogenase isoenzyme 1 and macro creatine kinase type 2 in a patient with lung cancer
Lentjes E.G.W.M., Backer E.T. 75

Enzyme-linked immunosorbent assay (ELISA) for human epidermal growth factor (hEGF)
Abe Y., Sagawa T., Sakai K., Kimura S. 87
A highly sensitive enzyme immunoassay of anti-insulin antibodies in human serum
Kohno T., Ishikawa E., Sugiyama S., Kamamo M., Kuzuya H., Imura H. 97
Effects of surfactants on human serum biotinidase
Hayakawa K., Oizumi J. 109

Vol. 168/2, 1987

Hemoglobin J-Baltimore (β16(A13)Gly → Asp): Interference with the assay of HbA(1c)
Vandenesch F., Baklouti F., Francina A., Vianey-Liaud C., Bertrand A., Le Devehat C., Delaunay J. 121
Detection of glucose-6-phosphate dehydrogenase deficiency in erythrocytes: A spectrophotometric assay and a fluorescent spot test compared with a cytochemical method
Wolf B.H.M., Weening R.S., Schutgens R.B.H., Van Noorden C.J.F., Vogels I.M.C., Nagelkerke N.J.D. 129
Measurement of fetal surfactant production by fluorescence polarization of amniotic fluid in complicated pregnancies
Masturzo P., Salmona M., Rossi G., Luerti M. 137
Stable isotope dilution analysis of pipecolic acid in cerebrospinal fluid, plasma, urine and amniotic fluid using electron capture negative ion mass fragmentography
Kok R.M., Kaster L., De Jong A.P.J.M., Poll-The B., Saudubray J.-M., Jakobs C. 143
The molecular nature of cholecystokinin in human plasma
Cantor P., Rehfeld J.F. 153
Direct determination of lead in blood by differential pulse stripping voltammetry
Nedeljkovic M., Stojadinovic L., Matovic V., Koricanac Z. 169
Histamine uptake by human platelets
Gill D.S., Barradas M.A., Mikhailidis D.P., Dandona P. 177
Studies with an automated kinetic assay for plasma angiotensin-converting enzyme activity and its potentiation by zinc ion
Roulston J.E., Allan D. 187
Purification and characterization of a pancreas cancer-related antigen
Furukawa J., Shimano T., Inaji H., Saito M., Matsuura N., Monden T., Mori T. 219
Determination of acid phosphatase in biological fluids using a new substrate, 2,6-dichloro-4-nitrophenyl phosphate
Teshima S., Hayashi Y., Ando M. 231
Development of an automated kinetic Jaffe method designed to minimise bilirubin interference in plasma creatinine assays
Knapp M.L., Mayne P.D. 239
Fructosamine measurements in serum and plasma
MacDonald D., Pang C.-P., Cockram C.S., Swaminathan R. 247

Vol. 168/3, 1987

Stability of radioiodinated ferritin
Covell A.M., Cook J.D. 261
A second albumin variant in an Irish population
Leahy D.T., McLaughlin H. 273
Plasma fibronectin in normolipidaemic and hyperlipidaemic uraemic patients treated with haemodialysis
De Giorgio L.A., Maffucci G., Toscano G., Gironi A., Vignoli S., Seghieri G. 281
Diagnostic significance of different urinary enzymes in patients suffering from chronic renal disease
Jung K., Schulze B.-D., Sydow K. 287

Microheterogeneity of serum alkaline phosphatase isoenzymes as revealed by isoelectric focusing
Rosendahl K., Waldenlind L., Onica D. 297
Determination of erythrocyte protoporphyrin by magnetic circular dichroism
Ivanetich K.M., Jeans D.R. 307
Laboratory and clinical features of self-poisoning with salbutamol and terbutaline
Jarvie D.R., Thompson A.M., Dyson E.H. 313
Determination of selenium in human spermatozoa and prostasomes using base digestion and electrothermal atomic absorption spectrophotometry
Suistomaa U., Saaranen M., Vanha-Perttula T. 323
Assay of sodium-lithium countertransport rate by direct measurement of erythrocyte lithium concentrations
Siebers R.W.L., Maling T.J.B. 329
A modified statistical approach for the detection of outlying values in external quality control: Comparison with other techniques
Thienpont L.M.R., Steyaert H.L.C., De Leenheer A.P. 337

Vol. 169/1, 1987

Biochemical tests in the diagnosis, classification, and management of patients with malignant lymphoma and leukemia
Goldberg D.M., Brown D. 1
New conjugated urinary metabolites in intermediate type maple syrup urine disease
Hagenfeldt L., Naglo A.S. 77
Plasma glutathione S-transferase measurements in patients with alcoholic cirrhosis
Beckett G.J., Hayes P.C., Hussey A.J., Bouchier I.A.D., Hayes J.D. 85
Lactate dehydrogenase, isoenzyme patterns and cation levels in human breast gross cyst fluid
Biagioni S., Mannello F., Stella F., Battistelli S., Marcheggiani F., Cerroni L., Artico M., Terzano C., Troccoli R. 91
Improved method for the quantification of aminoglutethimide in urine samples
Van Tellingen O., Bonfrer J.M.G., Nooyen W.J., Bruning P.F. 99
Detection of 3β-hydroxysteroid dehydrogenase deficiency in a newborn by means of urinary steroid analysis
Wolthers B.G., De Vries I.J., Volmer M., Nagel G.T. 109
Removal of interference by antihypertensive drugs in the spectrophotometric assay of metanephrines
Crawford G.A., Gallery E.D.M., Gyory A.Z. 117
A simple diagnostic method of adrenoleukodystrophy: Total fatty acid analysis of erythrocyte membranes
Antoku Y., Sakai T., Tsukamoto K., Imanishi K., Ohtsuka Y., Iwashita H., Goto I. 121
Urinary bile alcohols in liver dysfunction
Hiraoka T., Kihira K., Kohda T., Kosaka D., Kajiyama G., Hoshita T. 127
Effect of insulin, S-adenosylhomocysteine, phospholipase C, n-butanol and Triton X-114 on alkaline phosphatase from isolated rat adipocyte plasma membranes
Sykes E., Ghag S., Epstein E., Kiechle F.L. 133
Sandwich enzyme immunoassay of total insulin in the serum containing insulin antibodies
Tominaga M., Honda M., Itoh Y., Mokuda O., Ikeda T., Mashiba H. 141
Problems and promises arising from measuring salivary testosterone
Navarro M.A., Villabona C.M., Bonnin M.R. 143

Vol. 169/2–3, 1987

Increased excretion of propan-1,3-diol and 3-hydroxypropionic acid apparently caused by abnormal bacterial metabolism in the gut
Pollitt R.J., Fowler B., Sardharwalla I.B., Edwards M.A., Gray R.G.F. 151

Source of elevated serum mitochondrial creatine kinase activity in patients with malignancy
Okano K., Yamamoto K., Ohba Y., Matsumura K., Miyaji T. 159
Alkaline phosphatase in human milk: A new heat-stable enzyme
Chuang N.N. 165
A new assay for choline-containing phospholipids in amniotic fluid by an enzyme sensor
Campanella L., Tomassetti M., De Angelis G., Sammartino M.P., Cordatore M. 175
Separation of 5'-nucleotide phosphodiesterase isoenzymes of human serum by anion exchange high pressure liquid chromatography
Luthje J., Ogilvie A. 183
Comparison of plasminogen activator inhibitor activity and antigen in plasma samples
Urden G., Hamsten A., Wiman B. 189
Quantitation of serum carotenoid concentrations in healthy inhabitants by high-performance liquid chromatography
Ito Y., Sasaki R., Minohara M., Otani M., Aoki K. 197
Chromatographic profiles of urinary isoenzymes in healthy children
Matsukura H., Suzuki Y., Takai R., Okada T., Naiki S., Sakuragawa N. 209
Sulfation of minoxidil by human platelet sulfotransferase
Johnson G.A., Baker C.A. 217
Radioimmunoassay of glycosylated albumin with monoclonal antibody to glucitol-lysine
Ohe Y., Matsuura M., Nakajima Y., Shin S., Hashimoto F., Hirota Shima M.K. 229
An enzyme immunoassay for pepsinogen II (PGII): chronological changes in serum PGII concentrations
Matsusako K., Itoh M., Yokochi K., Miyamoto T., Joh T., Takeuchi T., Moriyama A., Kato T., Tanaka R. 239
Isolation and quantitation of apolipoproteins A-I and A-II from human high-density lipoproteins by fast-protein liquid chromatography
Weisweiler P., Friedl C., Ungar M. 249
Evaluation by immunoblotting of the criteria for tubular lesions diagnostic with urinary SDS-PAGE
Perini J.-M., Dehon B., Marianne Th., Foulard M., Roussel Ph. 255
Improved methods for the detection of unique sequences in Southern blots of mammalian DNA by non-radioactive biotinylated DNA hybridization probes
Gregersen N., Koch J., Kolvraa S., Bruun Petersen K., Bolund L. 267
Characterization of human middle ear mucus glycoprotein in chronic secretory otitis media (CSOM)
FitzGerald J.E., Green G.G.R., Stafford F.W., Birchall J.P., Pearson J.P. 281
Interactions of foetal steroid binding protein with other binding proteins in human serum
Forbes A., Wilkinson M.L., Jawed Iqbal M., Williams R. 299
Faecal haem assay — some practical modifications of the haemoquant assay for haemoglobin in faeces
Van den Berg J.W.O., Edixhoven-Bosdijk A., Koole-Lesuis R., Wilson J.H.P. 319
Decrease of serum malondialdehyde in patients treated with chlorpromazine
Bindoli A., Rigobello M.P., Cavallini L., Dalla Libera A., Galzigna L. 329

Vol. 170/1, 1987

Biochemical and functional properties of protein C and protein S
Strickland D.K., Kessler C.M. 1
Protein C and protein S clinical perspectives
Kessler C.M., Strickland D.K. 25
Production, purification and characterization of antibodies to 1,25-dihydroxyvitamin D raised in chicken egg yolk
Bauwens R.M., Kint J.A., Devos M.P., Van Brussel K.A., De Leenheer A.P. 37
Serum concentrations of a granulocyte-derived calcium-binding protein in cystic fibrosis patients and heterozygotes
Hayward C., Glass S., Van Heyningen V., Brock D.J.H. 45

Low molecular mass trypsin inhibitors in normal and osteoarthritic human articular cartilage
Lindenhayn K., Haupt R., Heilmann H.-H. 57
Neural tube defect-specific acetylcholinesterase: Its properties and quantitation in the detection of anencephaly and spina bifida
Bonham J.R., Dale G., Atack J.R. 69
High performance liquid chromatography of plasma aminoacids using orthophthalaldehyde derivatisation
Blundell G., Brydon W.G. 79
Modification of transferrin by polysaccharides in cancer patients
Narita K., Kanamaru I., Makino Y., Konno K. 85
Effect of short-term digoxin treatment on ouabain-sensitive sodium transport of erythrocytes
Smith J.B., Luft F.C., Wade M.B., Fineberg N.S., Weinberger M.H. 89
Sandwich enzyme immunoassay for human interleukin-1α produced in vitro by peripheral blood mononuclear cells
Tanaka K., Ishikawa E., Ohmoto Y., Hirai Y. 97
A simple assay for serum arylesterase using a specific substrate
Tanaka H., Kitamura M., Inoue Y. 105
Zinc contamination due to the collection tube
De Haan K.E.C., De Groot C.J., Boxma H., Van Den Hamer C.J.A. 111

Vol. 170/2–3, 1987

Identification and partial purification of a (Na-K)ATPase stimulating serine protease from plasma of insulin-dependent diabetics
Finotti P., Verbaro R. 121
Altered calmodulin activity in buccal epithelial cells from cystic fibrosis patients
McPherson M.A., Tiligada E., Bradbury N.A., Goodchild M.C. 135
Relationship of branched-chain α-keto-acids and lipids in sera of hypertriglyceridemic subjects
Penttila A.M., Herranen J., Lampainen E., Voutilainen E. 143
The quantification of gamma-aminobutyric acid in the cerebrospinal fluid by a radioreceptorassay
Mousah H., Jacqmin P., Lesne M. 151
Low molecular weight thiol content in glutathione synthetase-deficient human fibroblasts
Debieu D., Deschavanne P.J., Malaise E.P. 161
In vivo metabolism of apolipoprotein S in humans. Comparison with apolipoprotein A-I metabolism
Malmendier C.L., Lontie J.-F., Delcroix C. 169
Characterization of non-TCA-precipitable 'Lowry protein' in plasma of patients with acute renal failure
Haag M., Meyer H.E., Schollmeyer P., Horl W.H. 181
A new micro-assay for human liver alanine: Glyoxylate aminotransferase
Allsop J., Jennings P.R., Danpure C.J. 187
Primary hyperparathyroidism and hyperuricaemia are associated but not correlated with indicators of bone turnover
Broulik P.D., Stepan J.J., Pacovsky V. 195
Detection of alpha-1-antitrypsin deficiency variants by synthetic oligonucleotide hybridization
Klasen E.C., Hofker M.H., Van Paassen H.M.B., Verlaan-De Vries M., Bos J.L., Frants R.R. 201
Stability of protein and mRNA in human postmortem liver-analysis by two-dimensional gel electrophoresis
Finger J.M., Mercer J.F.B., Cotton R.G.H., Danks D.M. 209
Effect of bromelain on alkaline phosphatases of intestinal and non-intestinal tissues and serum
Kalimanovska V., Whitaker K.B., Moss D.W. 219
Development of a reference method for determining urinary oxalate by means of isotope dilution-mass spectrometry (ID-MS) and its usefulness in testing existing assays for urinary oxalate
Koolstra W., Wolthers B.G., Hayer M., Elzinga H. 227

An improved high performance liquid chromatographic method for determining urinary oxalate, making use of an ID-MS reference method
Koolstra W., Wolthers B.G., Hayer M., Rutgers H.M. 237
Determination of transcortin in serum by polyethylenglycol enhanced immunonephelometry. A comparison with equilibrium dialysis and radioimmunoassay
Eriksson C.-G., Eneroth P. 245
An automated colorimetric assay for urine nicotine metabolites: a suitable alternative to cotinine assays for the assessment of smoking status
Puhaikanen E.V.J., Barlow R.D., Salonen J.T. 255
Effect of long preincubation on the two forms of human erythrocyte prolidase
Myara I. 263
Semi-automated enzyme-linked immunosorbent assay (ELISA) for the quantification of apolipoprotein B using monoclonal antibodies
Bojanovski M., Gregg R.E., Wilson D.M., Brewer Jr. H.B. 271
Screening for adenine and hypoxanthine phosphoribosyltransferase deficiencies in human erythrocytes by high-performance liquid chromatography
Sakuma R., Nishina T., Kitamura M., Yamanaka H., Kamatani N., Nishioka K. 281
Collagenase-like (CL) peptidase activity in synovial fluid from patients with rheumatoid arthritis
Ito A., Hagihara M., Nagatsu T., Iwata H., Miura T. 291
Increased urinary excretion of argininosuccinate in type II citrullinemia
Saheki T., Kobayashi K., Inoue I., Matuo S., Hagihara S., Noda T. 297
Use of GC/MS/SIM for rapid determination of plasma levels of o,p′-DDD, o,p′-DDE and o,p′-DDA
Inouye M., Mio T., Sumino K. 305
A new radioimmunoassay of free thyroxine using ^{12}A5I-labelled thyroxine-protein complex uninfluenced by albumin and thyroxine-binding globulin
Tsutsumi s., Ishibashi K., Miyai K., Nagase S., Ito M., Amino N., Endo Y. 315
Accumulation of phosphorylated dolichol in several tissues in ceroid-lipofuscinosis (Batten disease)
Hall N.A., Patrick A.D. 323
Pharmacokinetics of desferrioxamine and of its iron and aluminium chelates in patients on haemodialysis
Allain P., Chaleil D., Mauras Y., Beaudeau G., Varin M.C., Poignet J.L., Ciancioni C., Ang K.S., Cam G., Simon P. 331
Correlation between qualitative and quantitative faecal occult blood tests
Robertson J.D., Maughan R.J., Davidson R.J.L. 339
Increased urinary excretion of the retinol-binding protein in insulin-dependent diabetes mellitus in the absence of microalbuminuria
Holm J., Hemmingsen L., Nielsen N.V., Thomsen M. 345
Hormone-induced redistribution of protein kinase C in human pituitary adenomas
Ishizuka T., Ito Y., Murayama M., Miura K., Nagao S., Nozawa Y. 351

Vol. 171/1, 1988

A rapid and sensitive enzyme immunoassay for Cu/Zn superoxide dismutase with polyclonal and monoclonal antibodies
Porstmann T., Wietschke R., Schmechta H., Grunow R., Porstmann B., Bleiber R., Pergande M., Stachat S., Von Baehr R. 1
Analysis of the fatty acid composition of erythrocyte phospholipids by a base catalysed transesterification method — Prevention of formation of dimethylacetals
Laryea M.D., Cieslicki P., Diekmann E., Wendel U. 11
A sensitive, direct colorimetric assay of serum iron using the chromogen, nitro-PAPS
Makino T., Kiyonaga M., Kina K. 19
Quantitation and characterization of urinary glycosaminoglycans in healthy subjects and patients with mastocytosis
Kettelhut B.V., Metcalfe D.D. 29

Development of fluoroimmunoassays for the specific detection of morphine in urine
Colbert D.L., Gallacher G., Ayling P., Turner G.J. 37

A new method for the determination of L-DOPA and 3-O-methyldopa in plasma and cerebrospinal fluid using gas chromatography and electron capture negative ion mass spectrometry
De Jong A.P.J.M., Kok R.M., Cramers C.A., Wadman S.K., Haan E. 49

Organ specific properties for human urinary alkaline phosphatases
Miura M., Koyama I., Matsuzaki H., Sakagishi Y., Ikezawa H., Komoda T. 63

Red cell pyrimidine 5′ nucleotidase deficiency: Determination of nucleotidase activity and nucleotide content using HPLC
Adair C.G., Elder G.E., Lappin T.R.J., Bridges J.M., Nelson M.G. 75

Effect of renal insufficiency on the concentration of free retinol-binding protein in urine and serum
Bernard A., Vyskocyl A., Mahieu P., Lauwerys R. 85

3-Hydroxy-3-methylglutaryl-CoA lyase in human skin fibroblast: Study of its properties and deficient activity in 3-hydroxy-3-methylglutaric aciduria patients using a simple spectrophotometric method
Wanders R.J.A., Schutgens R.B.H., Zoeters P.H.M. 95

Changes in the bone and liver isoenzymes of alkaline phosphatase in postmenopausal women being treated with norethisterone
Deeny M., Farish E., Tillman J., Dagen M., Hart D.M., Fletcher C.D. 103

Estimation of pyruvate dehydrogenase (E_1) activity in human skeletal muscle; three cases with E_1 deficiency
Van Laack H.L.J.M., Ruitenbeek W., Trijbels J.M.F., Sengers R.C.A., Gabreels F.J.M., Janssen A.J.M., Kerkhof C.M.C. 109

Vol. 171/2–3, 1988

Flow cytometry in the clinical laboratory
Ryan D.H., Fallon M.A., Horan P.K. 125

Combined enzymatic assay of phosphatidylglycerol and phosphatidylcholine in amniotic fluid
Farquharson J., Jamieson E.C., Berry E., Sutton A., Logan R.W. 175

Separation and partial characterization of two galactosyltransferase isoforms from malignant ascitic fluid
Boyle F.A., Cook N.D., Peters T.J. 187

Altered erythrocyte nucleotide patterns are characteristic of inherited disorders of purine or pyrimidine metabolism
Simmonds H.A., Fairbanks L.D., Morris G.S., Webster D.R., Harley E.H. 197

GC and GC-MS procedures for simultaneous phenotyping with dextromethorphan and mephenytoin
Baumann P., Jonzier-Perey M. 211

Stable isotope dilution analysis of succinylacetone using electron capture negative ion mass fragmentography: An accurate approach to the pre- and neonatal diagnosis of hereditary tyrosinemia type I
Jakobs C., Dorland L., Wikkerink B., Kok R.M., De Jong A.P.J.M., Wadman S.K. 223

Assessment of taurine deficiency in cystic fibrosis
Thompson G.N. 233

The effect of renal failure and haemodialysis on the concentration of free apolipoprotein A-1 in serum and the implications for the catabolism of high-density lipoproteins
Neary R.H., Gowland E. 239

Optimization of histamine radio enzyme assay with purified histamine-N-methyltransferase
Harvima R.J., Harvima I.T., Fraki J.E. 247

Newborn screening for cystic fibrosis using an enzyme linked immunosorbent assay (ELISA) technique
Bowling F.G., Brown A.R.D. 257

The characterisation of molecular forms of acetylcholinesterase in Hirschsprung's disease
Bonham J.R., Dale G., Scott D.J., Wagget J., Atack J.R. 263
Microheterogeneity of paraproteins. I. Diagnostic value of isoelectric focusing followed by immunoblotting
Schipper H.I., Bertram G., Kaboth U. 271
Purine metabolism in normal and ITP-pyrophosphohydrolase-deficient human erythrocytes
Van Waeg G., Niklasson F., Ericson A., De Verdier C.-H. 279
Effect of insulin treatment on fatty acids of plasma and erythrocyte membrane lipids in type 2 diabetes
Tilvis R.S., Taskinen M.-R., Miettinen T.A. 293
Apoprotein AII in acute pancreatitis: Intriguing improvement in the prediction of fatal outcome
Schneider J., Meier M., Maroske D., Kaffarnik H. 305
A simple fluorometric method for the determination of serum free carnitine
Maehara M., Kinoshita S., Watanabe K. 311
Improved purification of human liver alkaline phosphatase by phenyl-Sepharose column chromatography
Takata K.-I., Sumikawa K., Saeki K., Okochi T., Adachi K. 317
An alpha-fetoprotein radioimmunoassay for use in maternal serum screening for Down's syndrome: Limitations and improvements
Mooney R.A., McGorty B.L., French C.A., Arvan D.A. 325

Vol. 172/1, 1988

Laboratory analyses for steroid hormone receptors, and their applications to clinical medicine
Mayer T.K., Mooney R.A. 1
A sensitive kinetic assay of serum albumin based on its enzyme-like hydrolytic activity, using a new chromogenic and fluorogenic substrate
Gurakar A.R., Wolfbeis O.S. 35
Fetal lung maturity tests on the 10000 x g pellet
McCulloch J.C., Mendelsohn D. 47
Sialic acid concentration in erythrocyte membrane subfractions in patients with myotonic dystrophy and healthy controls
Sydow O. 57
Molecular analysis of peroxisomal β-oxidation enzymes in infants with Zellweger syndrome and Zellweger-like syndrome: Further heterogeneity of the peroxisomal disorder
Suzuki Y., Shimozawa N., Orii T., Igarashi N., Kono N., Hashimoto T. 65
Re-evaluation of normal human urinary proteins fractionated by one-dimensional sodium dodecyl sulphate-polyacrylamide gel electrophoresis
Shiba K.S., Kanamori K., Cho H., Furuhata N., Harada T., Shiba A., Nakao M. 77
Neutrophil chemiluminescence induced by platelet activating factor and suppressed by C-reactive protein
Tatsumi N., Hashimoto K., Okuda K., Kyougoku T. 85
Determination of cytochromes in human liver contaminated with hemoglobin
Tanaka A., Maki A., Kamiyama Y., Yamaoka Y., Ozawa K., Orii Y. 93
A new methjod of inhibiting glycolysis in blood samples
Uchida K., Matuse R., Toyoda E., Okuda S., Tomita S. 101
Concanavalin A- or phorbol ester-induced translocation of protein kinase C in thymoma cells from a patient with myasthenia gravis
Ishizuka T., Ito Y., Mori H., Miura K., Nagao S., Nagata K.-I., Nozawa Y. 109
Rapid high-performance liquid chromatography for pseudouridine assay in serum and urine
Amuro Y., Nakaoka H., Shimomura S., Tamura S., Hada T., Higashino K. 117
Elevated excretion of N-acetylated branched-chain amino acids in maple syrup urine disease
Lehnert W., Werle E. 123

Vol. 172/2–3, 1988

An enzymatic method for the α-amylase assay which comprises a new procedure for eliminating glucose and maltose in biological fluids
Kondo H., Shiraishi T., Nagata K., Tomita K. 131
Characterization of an adenine phosphoribosyltransferase deficiency
Chiba P., Zwiauer K., Muller M.M. 141
Electroimmunoassay of human serum lipoproteins containing Apo A-I by monoclonal antibodies
Ayrault-Jarrier M., Petit E., Bekaert E.D., Robin H., Pastier D., Beucler I., Polonovski J., Hervaud E., Pau B. 149
A rapid electrochemical assay of lecithin in amniotic fluid using a fluoride ion-sensitive electrode
Diomede L., Masturzo P., Agosti S., Ornaghi F., Salmona M. 161
Enzyme-linked immunosorbent assay for ileal human goblet cell mucin
Roberton A.M., Dalzell H., Harding C.A., Pearce S., Ram S.P., McKay E.J., Wesley A.W. 171
Direct quantitation of serum high density lipoprotein subfractions separated by gradient gel electrophoresis
Gambert P., Farnier M., Bouzerand C., Athias A., Lallemant C. 183
Urinary albumin excretion in healthy adult subjects: Reference values and some factors affecting their interpretation
Watts G.F., Morris R.W., Khan K., Polak A. 191
The presence and molecular forms of cardiodilatin immunoreactivity in the human and rat right atrium
Meleagros L., Ghatei M.A., Anderson J.V., Wharton J., Taylor K.M., Krikler D.M., Meijler F.L., Polak J.M., Bloom S.R. 199
Glutathione S-transferase levels in autoimmune chronic active hepatitis: A more sensitive index of hepatocellular damage than aspartate transaminase
Hayes P.C., Hussey A.J., Keating J., Bouchier I.A.D., Williams R., Beckett G.J., Hayes J.D. 211
Thyroid dysfunction and hypercholesterolaemia in the general population of Glasgow, Scotland
Series J.J., Biggart E.M., O'Reilly St. D.J., Packard C.J., Shepherd J. 217
Anomeric specificity of D-glucose metabolism in human erythrocytes
Malaisse-Lagae F., Sterling I., Sener A., Malaisse W.J. 223
Determination of medium chain fatty acids in serum
Vici C.D., Bachmann C., Gradwoht M., Colombo J.P. 233
Whole blood storage solution for erythrocyte sodium and sodium-lithium countertransport rate determination
Siebers R.W.L., Maling T.J.B. 239
The isolation and identification of indigoid pigments from urine
Jackson A.H., Jenkins R.T., Grinstein M., Ferramola de Sancovich A.-M., Sancovich H.A. 245
Identification of the carboxypeptidase responsible for the post-synthetic modification of creatine kinase in human serum
Hendriks D., Soons J., Scharpe S., Wevers R., Van Sande M., Holmquist B. 253
The distinction of bone and liver isoenzymes of alkaline phosphatase in serum using a monoclonal antibody
Seabrook R.N., Bailyes E.M., Price C.P., Siddle K., Luzio J.P. 261
A monoclonal antibody capture assay for intestinal alkaline phosphatase and the measurement of this isoenzyme in pregnancy
Bailyes E.M., Seymour P.M., Fulton I., Price C.P., Luzio J.P. 267
Fatty acid composition and kinetic behaviour of liver retinyl esters in vitamin A sufficient and deficient rats
Periquet B., Lambert W., Bailly A., Tomatis I., Ghisolfi J., De Leenheer A.P., Thouvenot J.P. 275
Inhibition of galactosyltransferase activity by cacodylate buffer
Boyle F.A., Cook N.D., Peters T.J. 291
Application of heparin-conjugated Sepharose for the measurement of cobalamin-saturated and unsaturated transcobalamin II
Van Kapel J., Wouters N.M.H., Lindemans J. 297

Analysis of cell surface glycoconjugates in fibroblasts from patients with cystic fibrosis
Sarras Jr. M.P., Huff J.K. 311
Presence of activator proteins for the enzymatic degradation of glucosylceramide in several human tissues
Vaccaro A.M., Ciaffoni F., Mandara I., Suzuki K. 323
Role of serum pepsinogen in detecting ulcer disease
Vianello F., Plebani M., Piccoli A., Tessaro P., Di Mario F., Naccarato R., Burlina A. 335

Vol. 173/1, 1988

The place of reference materials in clinical enzymology
Moss D.W. 1
Standardization by reference methods: A German viewpoint
Schmidt E.S., Schmidt F.W. 9
γ-Glutamyltransferase reference material: An example of international cooperation
Siest G., Schiele F., Muller J., Wellman M. 19
Molecular titration as a means of calibrating enzyme reference materials
Roth M., Selz L. 27
Influence of isoenzyme patterns on commutability in enzyme determinations
Burlina A., Bonvicini P., Plebani M., Zaninotto M. 35
Glutamate dehydrogenase: Biochemical and clinical aspects of an interesting enzyme
Schmidt E.S., Schmidt F.W. 43
Clinical biochemistry of peroxisomal disorders
Kaiser E., Kramar R. 57
Esterase D: Evaluation of a potential derived gene marker for hereditary retinoblastoma
Van Der Heiden C., Geurtzen A.F.M., Brink W., De Klerk J.B.C., Beemer F.A. 81
Lactate dehydrogenase M-subunit deficiency: A new type of hereditary exertional myopathy
Kanno T., Sudo K., Maekawa M., Nishimura Y., Ukita M., Fukutake K. 89

Vol. 173/2, 1988

Determination of zinc in blood serum by electrothermal atomic absorption spectrometry with matrix modification
Accominotti M., Pegon Y., Vallon J.-J. 99
Identification of renal calculi by computerized infrared spectroscopy
Lehmann C.A., McClure G.L., Smolens I. 107
The estimation of dihydropteridine reductase in human blood cells
Webber S., Hural J.A., Whitely J.M. 117
Accumulation of furancarboxylic acids in uremic serum as inhibitors of drug binding
Niwa T., Takeda N., Maeda K., Shibata M., Tatematsu A. 127
Similarity of the sugar moiety of human alkaline phoshphatases between the kidney cortex and duodenum, or medulla and ileum
Koyama I., Arai Y., Miura M., Matsuzaki H., Sakagishi Y., Komoda T. 139
Interleukin-2-dependent T lymphocytes for the diagnosis and investigation of inherited metabolic disorders
Adolph K., Kucey M.T., Hodges S.D., Carter R.J., Snyder F.F. 147
Diagnostic significance of estimation of serum apolipoprotein a along with α-fetoprotein in alcoholic cirrhosis and hepatocellular carcinoma patients
Nayak S.S., Kamath S.S., Kundaje G.N., Aroor A.R. 157
In vitro glycation of dried serum
Ch'ng S.L., Marinah T.A. 165
Alkaline phosphatase isoenzyme pattern in human amniotic fluid is dependent on the level of total activity. Implications in cystic fibrosis diagnosis
Dictus-Vermeulen C., Ameryckx J., Gueuning C., Van Bogaert E., Graff G.L.A. 173

A new microtechnique for the analysis of the human hepatic microsomal glucose-6-phosphatase system
Burchell A., Hume R., Burchell B. 183
Enzyme pattern and lipid peroxides in endomyocardial biopsies from patients with cardiomyopathy and myocarditis
Lehmann I., Papies B., Parsi R.A., Romaniuk P., Schimke I., Parsi E., Konig M.-L. 193
Comparison of separation techniques in radioimmunoassays for 17-hydroxyprogesterone
El-Gamal B.A., Landon J. 201
Monoclonal IgMκ with rheumatoid factor activity and cryoprecipitability identified only by immunofixation electrophoresis after 2-mercaptoethanol treatment
Sorensen S. 217
Agarose isoelectric focusing of arylsulfatase A producing a permanent record on slab gels.
Menna-Perper M., Nora R., Goldstein L., Manowitz P. 225
Pterins as biochemical tests for diagnosis and management of patients with malignant lymphoma and leukemia
Reibnegger G., Fuchs D., Hausen A., Werner E.R., Wachter H. 233
Impaired production of leukotrienes by peripheral blood polymorphonuclear leukocytes from children with chronic renal failure
Hamasaki T., Sakano T., Kobayashi M., Sakura N., Usui T., Ueda K. 235
The relationship between serum fructosamine and albumin
Senecal P.-E., Douville P., Simard S., Coulombe R. 239

Vol. 173/3, 1988

Minimal myocardial injury after uncomplicated coronary bypass surgery. Various sources of overestimation
Hermens W.T., Wilems G.M., Van Der Vusse G.J. 243
Early diagnosis of clostridial gas gangrene using sialidase antibodies
Roggentin P., Gutschker-Gdaniec G.H.M., Hobrecht R., Schauer R. 251
Urinary acylcarnitines in a patient with neonatal multiple acyl-CoA dehydrogenation deficiency, quantified by a carboxylic acid analyzer with a reversed-phase column
Kidouchi K., Niwa Nohara T.D., Asai K., Sugiyama N., Morishita H., Kobayashi M., Wada Y. 263
Reference limits of bone and liver alkaline phosphatase isoenzymes in the serum of healthy subjects according to age and sex as determined by wheat germ lectin affinity electrophoresis
Kuwana T., Sugita O., Yakata M. 273
A high-performance liquid chromatographic assay for acid ceramidase activity in cultured fibroblasts from patients with Farber's disease and from controls
Mitsuo K., Kobayashi T., Shinnoh N., Goto I. 281
Reactivities of human urinary prokallikrein and kallikrein toward rabbit antibodies with special reference to enzyme immunoassay
Irie A., Takahashi S., Katayama Y., Shibata Y., Miyake Y. 289
Random distribution of iron among the two binding sites of transferrin in patients with various hematologic disorders
Beguin Y., Huebers H., Finch C.A. 299
Succinic acidemia: a new syndrome of organic acidemia associated with congenital lactic acidosis and decreased NADH-cytochrome c reductase activity
Asano K., Miyamoto I., Matsushita T., Murakami Y., Minouri S., Wagatsuma T., Oshima M. 305
Pharmacokinetics of desferrioxamine and of its iron and aluminum chelates in patients on peritoneal dialysis
Allain P., Mauras Y., Varin M.C., Ang K.S., Cam G., Simon P. 313
Biochemical investigations on prolidase and prolinase in erythrocytes from patients with prolidase deficiency
Kodama H., Mikasa H., Ohhashi T., Ohno T., Arata J. 317

Correlation between serum lipid peroxides and the lesion size in cerebrovascular disease
Huang Z.-S., Lu F.-J., Lee T.-K. 325
The carbohydrate composition of human serum amyloid P component
Hess D., Ohishi H., Skinner M., Cohen A.S., Schmid K. 331
The chemiluminescence response of polymorphonuclear leukocytes from febrile patients
Miyata H., Moriguchi N., Kinoshita T. 337
Lack of association between plasma lecithin: cholesterol acyltransferase concentration and plasma sex hormone concentrations in men
Nanjee M., Albers J.J., Miller N.E. 343

Vol. 174/1, 1988

Methyltransferase and phopholipase A_2 activity in the cell membrane of neutrophils and lymphocytes from patients with Behcet's disease, systemic lupus erythematosus, and rheumatoid arthritis
Niwa Y., Miyachi Y., Sakane T., Kanoh T., Taniguchi S. 1
Elimination of undesirable immunoglobulin contaminants including aggregated IgG from γ-globulin preparations by jackfruit lectin affinity chromatography
Kondoh H., Kobayashi K. 15
Characterization of amylase linked immunoglobulin G to distinguish human salivary and pancreatic isoamylases
Moriyama T., Takebe T., Nobuoka M., Makino M. 25
Construction and clinical validation of a sensitive and specific assay for serum mature calcitonin using monoclonal anti-peptide antibodies
Motte P., Vauzelle P., Gardet P., Ghillani P., Caillou B., Parmentier C., Bohuon C., Bellet D. 35
DSC, X-ray and NMR properties of cholesterol-cholestanol-dihydrate crystals
Hirsch D., Azoury R., Sarig S. 55
Applications of immunoperoxidase techniques in specificity testing of monoclonal antibodies (Mabs) against von Willebrand factor (vWf)
Ingerslev J., Bukh A., Moller N.P.H., Stenbjerg S., Zeuthen J. 65
Enzyme immunoassay of cortisol in human plasma using penicillinase as label
Shrivastav G.S., Kumari G.L., Rao P.N. 83
Modified dual label Schilling test for pancreatic exocrine function
Leung J.W.C., Frost R.A., Burgess R., Braganza J.M., Slater D.M., Cotton P.B. 93
The Philadelphia variant of galactokinase: Impaired [1-^{14}C]galactose oxidation by intact erythrocytes
Soni T., Brivet M., Moatti N., Dahan N., Lemonnier A. 101
Effect of insulin on globin synthesis in human reticulocytes
Ballas S.K., Tran T.H. 111
Magnesium, zinc, calcium, potassium, lipoproteins and creatine kinases on the first day after acute myocardial infarction: Statistical analysis
Auget J.-L., Speich M., Arnaud P. 115

Vol. 174/2, 1988

A sensitive method to assay blood complement $C\bar{1}$; Inhibitor activity
Drouet C., Alibeu C., Ponard D., Arlaud G.J., Colomb M.G. 121
A comparative study on the diagnostic value of prostatic acid phosphatase (PAP) and prostatic specific antigen (PSA) in patients with carcinoma of the prostate gland
Van Dieijen-Visser M.P., Delaere K.P.J., Gijzen A.H.J., Brombacher P.J. 131
Ca^{2+} influx in normal and spherocytic red cells
Johnsson R., Saris N.-E. 141
Monoclonal antibodies to human kidney gamma-glutamyltransferase
Donati R., Sabolovic N., Wellman M., Artur Y., Siest G. 149

Evaluation of serum aminoterminal procollagen type III propeptide as an index of portal hypertension and esophageal varices in chronic liver diseases
Gressner A.M., Tittor W., Kropf J. 163
Quality of performance of assays for serum growth hormone in the United Kingdom (UK): Evidence from the UK external quality assessment scheme, 1980–1987
Seth J., Hanning I., Bacon R.R.A., Hunter W.M. 171
Factors associated with the quality of laboratory performance in the United Kingdom external quality assessment scheme for serum growth hormone
Seth J., Hanning I. 185
Plasma and biliary cholestanol related to steroid metabolism in familial hypercholesterolemia patients with and without ileal exclusion
Koivisto P.V.I., Miettinen T.A. 197
Quantification of soluble serum HLA Class I antigens in healthy volunteers and AIDS patients
Ferreira A., Villar M.L., Alvarez Cermeno J.C., Revilla Y., Garcia Rodriguez M.C., Fontan G., Gonzalez-Porque P. 207
Serum plant sterols and lathosterol related to cholesterol absorption in coeliac disease
Vuoristo M., Tilvis R., Miettinen T.A. 213
A new and rapid immunochemiluminometric assay for the measurement of Tamm-Horsfall glycoprotein
Horton J.K., Davies M., Woodhead J.S., Weeks I. 225

Vol. 174/3, 1988

Renal handling of cathodic trypsin-like immunoreactivity in man
Moller-Petersen J., Ladefoged J. 241
Albumin determination in frozen urines — Underestimated results
Erman A., Rabinov M., Rosenfeld J. 255
On the fluorometric assay of circulating lipoperoxides
Hendriks T., Assman R.F.T.A. 263
Sepiapterin reductase in human amniotic and skin fibroblasts, chorionic villi, and various blood fractions
Ferre J., Naylor E.W. 271
An altered calmodulin binding protein in cystic fibrosis — A clue to the biochemical defect
Shori D.K., Bradbury N.A., Goodchild M.C., Dormer R.L., McPherson M.A. 283
Effect of chronic ethanol ingestion on alpha-mannosidase isoenzymes in rat liver
Lauro P., Lechner P.S., Okolo A., Eagon P.K., Glew R.H. 291
Distribution of tocopherol among human plasma lipoproteins
Ogihara T., Miki M., Kitagawa M., Mino M. 299
Enzymatic analysis of total- and HDL-cholesterol: Comparison with the standardized Liebermann-Burchard method used by the lipid research clinics program
Bachorik P.S., Virgil D.G., Derby C., Widman D., McMahon R., Fulwood R.P., Ezzati T. 307
α-Amylase assay with use of a benzyl derivative of p-nitrophenyl α-maltopentaoside, BG5P
Satomura S., Sakata Y., Omichi K., Ikenaka T. 315
A comparative study of the accumulated sialic acid-containing oligosaccharides from cultured human galactosialidosis and sialidosis fibroblasts
Van Pelt J., Kamerling J.P., Vliegenthart J.F.G., Hoogeveen A.T., Galjaard H. 325
Urinary purines, pyrimidines and nucleosides in uridine-treated orotic aciduria
Mills G.C. 337
A pepstatin-immunometric assay for active renin in human plasma
Higaki J., Ogihara T., Nishiura M., Murakami K., Kumahara Y. 345
Stabilization of ascorbic acid in plasma
Khanduja K.L., Koul A. 351

Vol. 175/1, 1988

Serum lysosomal hydrolases in cystic fibrosis
Krall E.A., Basu A., Gloninger M.F., Glew R.H., Humphries L. 1

Enzyme immunoassay for α-human atrial natriuretic polypeptide — Direct measurement of plasma level
Hashida S., Ishikawa E., Nakao K., Mukoyama M., Imura H. 11

3-Hydroxyoctanoic aciduria: Identification of a new organic acid in the urine of a patients with non-ketotic hypoglycemia
Kelley R.I., Morton D.H. 19

Studies on serum immunoreactive prolyl 4-hydroxylase in liver disease — Its elevation both in hepatocellular damage and cholestatic disease
Nagai Y., Yoshiba M. 27

Enzyme-linked immunosorbent assays for serum pepsinogens I and II using monoclonal antibodies — With data on peptic ulcer and gastric cancer
Huang S.C., Miki K., Furihata C., Ichinose M., Shimizu A., Oka H. 37

Homogeneous immunoassay based on dihexadecyl-dimethylammonium bilayer membranes
Ihara H., Imamura T., Kunitake T., Hirayama C. 51

Assay of stable isotope labelled urea in biological fluids by selected ion monitoring
Nelson J.E., Ruo T.I. 59

Criteria for identyfying endogenous compounds as digoxin-like immunoreactive factors in humans
Lau B.W.C., Valdes Jr. R. 67

Urinary C_6-C_{12} dicarboxylic acylcarnitines in Reye's syndrome
Tracey B.M., Cheng K.N., Rosankiewicz J., Stacey T.E., Chalmers R.A. 79

Oestrogen receptors, lactate dehydrogenase and cellularity in human breast cancer
Hawkins R.A., Killen E., Tesdale A.L., Sangster K., Thomson M., Steele R.J.C., Blackie R.A.S. 89

The Philadelphia variant of galactokinase in human erythrocytes: Physicochemical and catalytic properties
Soni T., Brivet M., Moatti N., Lemonnier A. 97

Total and free quinidine by fluorescence polarization immunoassay and comparison with high performance liquid chromatography
Chen B.H., Taylor E.H., Kennedy E., Ackerman B., Olsen K., Pappas A.A. 107

Creatine kinase BB and neuron-specific γ-enolase are enhanced in leiomyosarcoma
Usui A., Kato K., Abe T., Hachisuka K., Yamaguchi A. 109

Measurement of platelet factor 4 and β-thromglobulin by an enzyme-linked immunosorbent assay
Takahashi H., Yoshino N., Shibata A. 113

Vol. 175/2, 1988

Isoelectric focusing of human neutrophil alkaline phosphatase isoenzyme in agarose gel
A. Brisson-Lougarre, H. Vergnes, J. Grozdea, R. Biorme 121

Behaviour of L-γ-glutamyl-4-nitroanilide and L-γ-glutamyl-3-carboxy-4-nitroanilide with respect to γ-glutamyltransferases of different origin
P.M.M. Verhoeff, F.P.A.M.N. Peters, H. Steigstra, J.C.M. Hafkenscheid 129

Clinical utility of serum fructosamine in diabetes mellitus compared with hemoglobin A_{1c}
P. Jerntrop, G. Sundkvist, G. Fex, J.-O. Jeppsson 135

Determination of cholic acid and chenodeoxycholic acid pool sizes and fractional turnover rates by means of stable isotope dilution technique, making use of deuterated cholic acid and chenodeoxycholic acid
B.J. Koopman, F. Kuipers, C.M.A. Bijleveld, J.C. Van der Molen, G.T. Nagel, R.J. Vonk, B.G. Wolthers 143

Tumour markers in the cerebrospinal fluid of patients with central nervous system metastases from extracranial malignancies
A.P. van Zanten, A. Twijnstra, B.W. Ongerboer de Visser, A.A.M. Hart, W.J. Nooyen 157

Kinetic studies on nitrite and nitrate in rats by ion-pair chromatography
C. Stein, H.-G. Classen, G. Schwedt 167
A more simple, rapid and sensitive fluorimetric method for the determination of isoniazid and acetylisoniazid in serum. Application for acetylator phenotyping
P.C. Ioannou 175
Concentration differences in isoforms of transferrin in blood from alcoholics during abuse and abstinence
S. Petrén, O. Vesterbery 183
gamma-Glutamyl-transpeptidase in diabetics: a case control study
G. Colleredo-Mels, G. Bettale, G. Bellati, M. Guanziroli, V. Bertone, G. Angeli, G. Idéo 189
Letter to the Editor
Colorimetric measurement of plasma 3-hydroxybutyrate
J.F. Buttery, B.R. Chamberlain 197

Vol. 175/3, 1988

Factors influencing the production of creatinine: implications for the determination and interpretation of urinary creatinine and creatine in man
N.J. Fuller, M. Elia 199
Plasma and serum zinc concentrations: effect of time between collection and separation
J.L. English, K.M. Hambidge 211
Fast functional assay of protein C in whole plasma using a snake venom activator: evaluation in patients with congenital and acquired protein C deficiencies
H. Takahashi, M. Hanano, W. Tatewaki, A. Shibata 217
The regional variability of enzymes in the brain: relevance to CSF enzyme determinations
D.A. Armbruster, H.-D. Gruemer 227
Simultaneous ion-pair and partition liquid chromatography of acetaminophen, theophylline and salicylate with application to 500 toxicologic specimens
J. Osterloh, S. Yu 239
Effects of acetaldehyde on human red cell metabolism: evidence for the formation of enzyme inhibitors
L.R. Solomon 249
A sensitive time-resolved immunofluorometric assay of ferritin in serum with monoclonal antibodies
M.J. Khosravi, M.A. Chan, A.C. Bellem, E.P. Diamandis 267
Lipoprotein cholesterol concentrations in patients with Behçet's disease
T. Mitamura, S. Ohno, H. Ariga, T. Ohsaka, N. Iwasaki, H. Matsuda, H. Matsumiya 277
Doxorubicin-induced altered glycolytic patterns in the leukemic cell studied by proton spin echo NMR
J. Reglinski, W.E. Smith, C.J. Suckling, M. Al-Kabban, M.J. Stewart, I.D. Watson 285
Plasma prolidase and prolinase activity in alcoholic liver disease
B. Brosset, I. Myara, M. Fabre, A. Lemonnier 291
Development of a simple, fluoroimmunoassay for paraquat
R.C. Coxon, C. Rae, G. Gallacher, J. Landon 297
In vivo and in vitro ionized calcium variations induced by acute respiratory acid base disturbances
C.B. Kancir, P.H. Petersen, T. Madsen, A.S. Olesen 307
Measurement of endotoxin in cerebrospinal fluid from neonates and infants using a new endotoxin-specific chromogenic test
M. Inagaki, M. Nakajima, Y. Ando, S. Takashima, K. Takeshita, S. Tanaka, H. Tamura, T. Obayashi 315
Detection and purification of an abnormal protein from plasma of patients with rheumatoid arthritis
O. Oda, K. Maeda, H. Yamaga, Y. Kanamono, H. Iwata, T. Miura, M. Yoshida, S. Miyata, K. Ohta, N. Yamanaka 321

Stability of vitamin E in long-term stored serum
E.W. Gunter, W.J. Driskell, P.R. Yeager 329
An enzyme antigen immunoassay for the determination of neuron-specific enolase in serum samples
K. Sorensen, U. Brodbeck, E. Paus, B. Norgaard-Pedersen 337
Letters to the Editor
Diagnosis of Zellweger syndrome by rectal biopsy: immunoblot of peroxisomal β-oxidation enzyme and activity of dihydroxyacetone phosphate acyltransferase in rectal mucosa
N. Shimozawa, Y. Suzuki, T. Orii, J. Hashimoto 345
The frequency of C_5 + cholinesterase in the normal Japanese population
M. Oimomi, J. Ohkawa, S. Saeki, S. Baba 349

Vol. 176/1, 1988

Circadian and circannual rhythms of several enzymes of lysosomal origin in human plasma
G. Goi, A. Fabi, A. Lombardo, A.B. Burlina, G. Tettamanti, N. Montalbetti, M. Cavalleri, F. Halberg 1
Modification in Ramsay's method for correct measurement of total iron-binding capacity
H. Baadenhuijsen, L.G.J. Deimann, A.P. Jansen 9
An enzymatic method for the determination of butyrobetaine via conversion to carnitine after isolation by high performance liquid chromatography
A. Sandor, P.E. Minkler, S.T. Ingalls, C.S. Hoppel 17
Composition and surface properties of the bronchial liquid in adult patients with cystic fibrosis
H. Gilljam, O. Anderson, Å. Ellin, B. Robertson, B. Strandvik 29
Urinary excretion of a glucose-containing tetrasaccharide. A parameter for increased degradation of glycogen
J. Kumlien, M.A. Chester, B.S. Lindberg, P. Pizzo, D. Zopf, A. Lundblad 39
Modification of Concanavalin A patterns of α1-acid glycoprotein and α2-HS glycoprotein in alcoholic liver disease
M. Jezequel, N.S. Seta, M.M. Corbic, J.M. Feger, G.M. Durand 49
Acute phase reactants in the elderly
N. Milman, N. Graudal, H.C. Anderson 59
Determination of phospholipids in biological samples by an improved densitometric method on thin-layer chromatograms
R. Masella, A. Cantafora 63
Evaluation of urinary oestrogen assays after the menopause and their potential for screening
M.H. Rae, P.A. Mole, C.R. Paterson 71
Effect of methionine loading and endogenous hypermethioninaemia on blood mercaptans in man
H. Al Mardini, J. Leonard, K. Bartlett, S. Lloyd, C.O. Record 83
Antibody immobilization on swollen polystyrene tubes for the radioimmunological determination of total human serum IgE
J.E. Causse, I. Ricordel, P. Taulelle, M.T. Chateau, F. Moussel, A.M. Dupuy, B. Descomps, J.P. Yver 91
Plasma fructosamine assay in children with insulin-dependent diabetes mellitus
N. Shimojo, M. Hirai, K.i. Naka, C. Yoshikawa, K. Okuda, S. Aono, G. Isshiki, S. Fuji 101
Diamine and polyamine levels in the urine of healthy adults
Y. Yodiat, M. Weiser, M. Kreisel, U. Bachrach 107
I-cell disease: evidence for a mannose 6-phosphate independent pathway for translocatoin of lysosomal enzymes in lymphoblastoid cells
A. Tsuji, K. Omura, Y. Suzuki 115

Vol. 176/2, 1988

Selection and performance of monoclonal anti-C-reactive protein in ELISA quantitative assay
J.A. Martinez, J.M. Coll 123

Use of ^{86}Rb and ^{22}Na in assaying active and cotransport activities in human erythrocytes in a biracial population
C. Sharma, E.R. Dalferes Jr., D.S. Freedman, A. Asamoah, G.S. Berenson 133
Immunoaffinity purification of human erythrocyte prolidase
F. Endo, A. Tanoue, T. Ogata, K. Motohara, I. Matsuda 143
IgE pyroglobulinemia in multiple myeloma
M.L. Gallango, R. Suinaga, M. Ramirez 151
Validation of a radioreceptor assay for 1,25-dihydroxyvitamin D using selected ion monitoring GC-MS
H. Oftebro, J.A. Falch, I. Holmberg, E. Haug 157
Chromatography of serum on Sep-pak C_{18} corrects falsely elevated vitamin D metabolic levels measured by protein binding assay
B. Dean, M.S. Kolavcic, J.D. Wark, L.C. Harrison 169
Characterisation of a slow component of normal human serum albumin
S.O. Brennan, P.M. George, R.J. Peach 179
A kinetic procedure for the estimation of arginine in serum using arginine kinase
C.H. Konings 185
Changes of serum apo A-IV levels in a newborn with chylothorax treated with medium-chain triglyceride milk
M. Scishima, A. Noma, S. Yamaguchi, T. Orii, Y. Muto 195
Serum indole and skatole in patients with various liver diseases
Y. Suyama, C. Hirayama 203
Measurement of myo-inositol in diabetic sera by GC/MS/SIM using *n*-butylboronate derivatives
M. Hasegawa, K. Dol, S. Baba 207
Enzyme-linked immunosorbent assay (ELISA) for human tumor necrosis factor (hTNF)
Y. Abe, M. Miyake, T. Sagawa, S. Kimura 213
Elastase-like protease and elastolytic activities expressed in cultured dermal fibroblasts derived from lesional skin of patients with pseudoxanthoma elasticum, actinic elastosis, and cutis laxa
E. Schwartz, F.A. Cruickshank, M.G. Lebwohl 219
Letters to the Editor
Effect of serum albumin concentration on free thyroxin (Amerlex FT4) values in healthy non-pregnant and pregnant women
S. Sakata, T. Komaki, S.-i. Shiraki, N. Kumasaki, H. Iwata 225
A pitfall in the analysis of the fatty acid composition of plasma and erythrocyte membrane lipids
C. Popp-Snijders 227
Evaluation of sex hormone binding globulin among patients with thyroxine binding globulin excess
J. Wortsman, B.N. Premachandra, I.K. Williams, W. Rosner 229
Separation and purification of creatine kinase BB isoenzyme and neuron-specific enolase on DEAE-Sephadex A-50 chromatography
J.-G. Geng, H.-Z. Chen, Y.-F. Yang, C.Y. Jian, Z.-H. Qian, Q.-Q. Han, Y.-M. Yao 233
Choline supplementation increases serum alkaline phosphatase activity in rats
J. Wuarin-Bierman, L. Wecker 237

Vol. 176/3, 1988

The prolamin antibody reactivity against hordein polypeptides in sera from patients with coeliac disease
S.U. Friis, H. Sjöström, O. Norén, N. Rüdiger, D. Anthonsen 241
Sialic acid as a marker for differentiation between benign and malignant intracranial tumors
E. Marth, G. Flaschka, S. Stiegler, J.R. Möse 251
Synthesis of pyrenesulfonylamido-sphingomyelin and its use as substrate for determining sphingomyelinase activity and diagnosing Niemann-Pick disease
R. Klar, T. Levade, S. Gatt 259
Ion-exchange chromatography and clinical criteria in the screening of the aminoacidopathies
P.R. Parvy, J.I. Bardet, D.M. Rabier, J.-M. Saudubray, P.P. Kamoun 269

Amino acid composition of peptides remaining after in vitro digestion of a gliadin sub-fraction with duodenal mucosa from patients with coeliac disease
H.J. Cornell 279
Solid phase immunoassay for high molecular weight alkaline phosphatase in human sera using a specific monoclonal antibody
J.T. Deng, P.G. Parsons 291
Determination of human erythropoietin by radioimmunoassay. Method and clinical data
O.J. Nielsen 303
Field measurement of plasma and erythrocyte cholinesterases
R.A. Magnotti, Jr., K. Dowling, J.P. Eberly, R.S. McConnell 315
Enhanced urinary trypsin inhibitory activity in gentamicin-induced nephrotoxicity in rats
S. Smetana, S. Khalef, Z. Nitsan, N. Hurwitz, A. Miskin, Y. Bar-Khayim, Y. Birk 333
Measurement of cytochrome *c* oxidase activity in human liver specimens obtained by needle biopsy
Y. Sakai, A. Tanaka, I. Ikai, Y. Yamaoka, K. Ozawa, Y. Orii 343

Vol. 177/1,2, 1988

Assessment of the role of pancreatic proteases in human abdominal aortic aneurysms and occlusive disease
M.A. Dubick, G.C. Hunter, E. Perez-Lizano, G. Mar, M.C. Geokas 1
Effect of colchicine on collagen synthesis by liver fibroblasts in murine schistosomiasis
M.M. Mansour, M.A. Dunn, L.A. Salah 11
Strong activity of acid-stable trypsin inhibitor in bovine thrombin for clinical use
H. Sumi, H. Hamada, S. Maehara, H. Mihara 21
Enzyme-linked immunosorbent assay of lipoprotein(a) in serum and cord blood
A. Abe, S. Maeda, K. Makino, M. Seishima, K. Shimokawa, A. Noma, M. Kawade 31
Elevated nucleotide pyrophosphatase activity in cultured skin fibroblasts from patients with Lowe's syndrome
K. Horie, T. Yano, I. Funakoshi, I. Yamashima 41
Clinical relevance of increase neuron-specific enolase concentration in cerebrospinal fluid
C. Jacobi, H. Reiber 49
Stabilization and quantitative determination of disulfiram in human plasma samples
B. Johansson 55
Purification of acidic glutathione S-transferase from human lung, placenta and erythrocyte and the development of a specific radioimmunoassay for their measurement
A.F. Howie, J.D. Hayes, G.J. Beckett 65
The diagnostic value of urinary transferrin compared to serum prostatic specific antigen (PSA) and prostatic acid phosphatase (PAP) in patients with prostatic cancer
M.P. van Dieijen-Visser, M.W.G. Hendriks, K.P.J. Dekaere, A.H.J. Gijzen, P.J. Brombacher 77
Iron, copper and zinc concentrations in human sweat and plasma: the effect of exercise
O.J. Aruoma, T. Reilly, D. MacLaren, B. Halliwell 81
A method to detect the in vitro modification of serum beta-2-microglobulin in health and disease
M.A. Forbes, A.M. Cox, E.H. Cooper 89
Relationship between the urinary excretion of albumin and retinol-binding protein in insulin-dependent diabetics
J. Holm, L. Hemmungsen, N.V. Nielsen 101
Micromethod for determination of ATPase activities using bioluminescence technique
M. Steiner, H. Bauer, F. Krassnigg, W.-B. Schill, H. Adam 107
Letters to the Editors
Second antibodies covalently linked to Sepharose with 1,1′-carbonyldimidazole — effect of storage on radioimmunoassay separation efficiency
M.G. McConway, R.S. Chapman 115
Platelet intracellular calcium in patients with Duchenne muscular dystrophy
C. Lechi, D. Sinigaglia, P. Guzzo, G. Bonadonna, M. Zatti 119

Vol. 177/3, 1988

A simplified method for determination of uroporphyrinogen decarboxylase activity in human blood
Adjarov D.G., Elder G.H. 123
Levels of 5′-nucleotide phosphodiesterase isoenzymes in normal and pathological sera
Luthje J., Pickert S., Ogilvie A., Hornemann E., Siegfried W., Waldherr A., Domschke W. 131
Serum procollagen III N-terminal peptide and laminin P1 fragment concentrations in alcoholic liver disease and primary biliary cirrhosis
Van Zanten R.A.A., Van Leeuwen R.E., Wilson J.H.P. 141
Plasma alkaline phosphatase isoenzymes in diabetes mellitus
Tibi L., Collier A., Patrick A.W., Clarke B.F., Smith A.F. 147
Correlation between concentrations of magnesium, zinc, and potassium in plasma, erythrocytes and muscles
Ladefoged K., Hagen K. 157
Diagnostic efficiency in discriminating liver malignancies from cirrhosis by serum gamma-glutamyltransferase isoforms
Sacchetti L., Castaldo G., Cimino L., Budillon G., Salvatore F. 167
Fat macroglobule formation from chylomicrons and non-traumatic fat embolism
Hulman G. 173
Dipeptidyl peptidase III and alanyl aminopeptidase in the human seminal plasma: origin and biochemical properties
Vanha-Perttula T. 179
A specific method for the direct determination of lipoprotein cholesterol in electrophoretic patterns
Aufenanger J., Haux P., Weber U., Kattermann R. 197

Vol. 177/3, 1988

Urinary neopterin excretion in pulmonary sarcoidosis: Correlation to clinical course of the disease
Prior C., Frank A., Fuchs D., Hausen A., Judmaier G., Reibnegger G., Werner E.R., Wachter H. 211
Precipitation of LDL with sulfated polyanions: Three methods compared
Siekmeier R., Marz W., Gross W. 221
Hb G-Philadelphia, or [α68(E17)Asn → Lys], in North Sardinia: Detection by isoelectric focusing and identification by HPLC of tryptic peptides
Manca L., Demuro P., Masala B. 231
Discovery of methoxyacetylcarbamide in the urine of normal adults and phenylketonuric children
Antoshechkin A.G., Tatur V.Y., Maximova L.A. 239
Accurate and specific HPLC assay of coproporphyrinogen III oxidase activity in human peripheral leucocytes
Guo R., Lim C.K., Peters T.J. 245
Quantitative determination of biocytin in urine of patients with biotinidase deficiency using high-performance liquid chromatography (HPLC)
Suormala T.M., Baumgartner E.R., Bausch J., Holick W., Wick H. 253
The effect of age and sex on β-hexosaminidase in urine
Hultberg B., Isaksson A., Berg B., Tryding N., Ekman S., Nilsson J.-E. 271
Approved recommendation (1988) on the theory of reference values. Part 3. Preparation of individuals and collection of specimens for the production of reference values
Solberg H.E. S3

Vol. 178/1, 1988

Apolipoprotein measurements in clinical biochemistry and their utility vis-a-vis conventional assays
Bachorik P.S., Kwiterovich Jr. P.O. 1

Colorimetric method for assay of serum γ-glutamyltransferase activity with some L-γ-glutamyl-carboxyanilides
Szewczuk A., Kuropatwa M., Lang D. 35
Serum noncholesterol sterols related to cholesterol metabolism in familial hypercholesterolemia
Gylling H., Miettinen T.A. 41
Sodium-lithium countertransport activity and its sensitivity to inhibitors with erythrocyte ageing in man
Carr S., Thomas T.H., Wilkinson R. 51
Determination of 3,4-dihydroxyphenylalanine (DOPA) in plasma and cerebrospinal fluid by high performance liquid chromatography with electrochemical detection
Boomsma F., Van der Hoorn F.A.J., Man in 't Veld A.J., Schalekamp M.A.D.H. 59
A sensitive non-radioactive assay for pyrimidine nucleoside phosphorylase using reversed-phase high performance liquid chromatography
Laurensse E.J., Pinedo H.M., Peters G.J. 71
Interaction of horse-spleen ferritin with aluminium citrate
Cochran M., Chawtur V. 79
Plasma acetate levels in response to intravenous fat or glucose/insulin infusions in diabetic and non-diabetic subjects
Akanji A.O., Ng L., Humphreys S. 85
Thermostability of human serum biotinidase activity
Hayakawa K., Oizumi J. 95
Allelic polymorphism of human placental alkaline phosphatase in three races: sugar chain heterogeneities
Sato M., Furiya K., Tanaka S., Sakagishi Y., Komoda T. 101

Vol. 178/2, 1988

Correlation of two different assays for urinary kallikrein in normotensive and hypertensive subjects
Maier M., Jerabek I., Reissert G., Holtzl E., Binder B.R. 127
An immunobioluminescence assay for gamma-gamma enolase activity in human serum and cerebrospinal fluid
Wevers R.A., Theunisse A.W.G., Rijksen G. 141
Serum pseudouridine as a biochemical marker in patients with hepatocellular carcinoma
Amuro Y., Nakaoka H., Shimomura S., Fujikura M., Yamamoto T., Tamura S., Hada T., Higashino K. 151
Competitive enzyme-linked immunosorbent assay (ELISA) for the quantitation of apolipoprotein A-I using a monoclonal antibody
Bojanovski M., Gregg R.E., Wilson D.M., Brewer Jr. H.B. 159
A 3,3',5-triiodothyronine autoantibody (IgG, lambda) in a case of Hashimoto's thyroiditis
Muratsugu M., Tsuchiya Y., Makino M. 171
Cholesterol monomer activity correlates with nucleation time in model bile
Chijiiwa K., Kiyosawa R., Nakayama F. 181
Enzyme immunoassay of medullasin in peripheral blood
Aoki Y., Yoshida M., Kominami E. 193
A sensitive two-site immunoassay for human pancreatic secretory trypsin inhibitor (PSTI) using monoclonal antibodies
Kurobe M., Kono M., Yoshida N., Hayashi K. 205
Comparison of the performance and clinical utility of a carboxy-terminal assay and an intact assay for parathyroid hormone
St. John A., Davies C., Riley W.J., Kent G.N., Brown R.C., Aston J.P., Weeks I., Woodhead J.S. 215
Serum osteocalcin and other indices of bone formation: An 8-decade population study in healthy men and women
Worsfold M., Sharp C.A., Davie M.W.J. 225

Vol. 178/3, 1988

Betaine metabolism in human neonates and developing rats
Davies S.E.C., Chalmers R.A., Randall E.W., Iles R.A. 241
Triglyceride lowering effect of MaxEPA fish lipid concentrate: a multicentre placebo controlled double blind study
Miller J.P., Heath I.D., Choraria S.K., Shephard N.W., Gajendragadkar R.V., Harcus A.W., Batson G.A., Smith D.W., Saynor R. 251
Enzyme-linked immunosorbent assay for quantitation of cereal proteins toxic in coeliac disease
Friis S.U. 261
Increase of erythrocyte resistance to hemolysis and modification of membrane lipids induced by hemodialysis
Peuchant E., Salles C., Vallot C., Wone C., Jensen R. 271
Level of plasma fibronectin in patients with breast cancer
Ruelland A., Kerbrat P., Clerc C., Legras B., Cloarec L. 283
Characterisation of galactosyltransferase isoforms by ion-exchange and lectin affinity chromatography
Boyle F.A., Peters T.J. 289
Carbamylated haemoglobin in chronic renal failure
Smith W.G.J., Holden M., Benton M., Brown C.B. 297
A second antibody solid-phase enzyme immunoassay of prostaglandin E_2 in human gastric juice
Orsini B., Calabro A., Renzi D., Pradelles P., Fedi P., Surrenti C. 305
Is the ex vivo rat gastric chamber model suitable for studying the gastrotoxicity of refluxed duodenal contents? Initial results using deoxycholic acid
Armstrong D., Farrell M., Hanby A., Murphy G.M., Dowling R.H. 313
Detection of lysozyme and alpha$_2$-macroglobulin-lysozyme complexes by immunoblotting
Morsky P. 327
Alterations in specific activity of lysosomal alpha-glucosidase in cystic fibrosis
Vladutiu G.D., Kewin C.C. 337
Enzymatic antioxidative defence of erythrocytes in an Italian family with Hb Volga or $\alpha_2\beta_2$ 27 (B9) Ala → Asp
Falcioni G., Grelloni F., De Sanctis G., Pierani P., Felici L., Coppa G.V. 345
Purine metabolism in a bone-marrow transplanted adenosine deaminase deficient patient
Boulieu R., Bory C., Souillet G. 349

Vol. 179/1, 1989

Evaluation of a direct assay for atrial natriuretic peptide
Tan A.C.I.T.L., Rosmalen F.M.A., Hofman J.A., Kloppenborg P.W.C., Benraad Th. J. 1
Specific determination of plasma dexamethasone by HPLC and RIA - application to standard dexamethasone suppression test in psychiatric patients
Lejeune-Lenain C., Bosson D., Linkowski P. 13
Clinical significance of a new isoform of serum alanine aminopeptidase; relationship with liver disease and alcohol consumption
Sanderink G.-J., Artur Y., Paille F., Siest G. 23
Association between increased concentrations of free thyroxine and unsaturated free fatty acids in non-thyroidal illnesses: role of albumin
Tikanoja S.H., Joutti A., Liewendahl B.K. 33
Complement activation after prolonged exercise
Dufaux B., Order U. 45
Urinary excretion of 2,3-butanediol and acetoin by babies on a special care unit
Mills G.A., Walker V. 51
Isolation and characterization of dipeptidyl aminopeptidase IV from human kidney cortex
Wolf G.B., Scherberich J.E., Fischer P., Schoeppe W. 61

Stability of alpha$_1$-microglobulin, beta$_2$-microglobulin and retinol binding protein in urine
Donaldson M.D.C., Chambers R.E., Woolridge M.W., Whicher J.T. 73
Validation of the Friedewald formula for the measurement of low density lipoprotein cholesterol in insulin-dependent diabetes mellitus
Winocour P.H., Ishola M., Durrington P.N. 79
Characterization of two monoclonal antibodies to human apolipoprotein A-I that also bind to rabbit apolipoprotein A-I. Application to ELISA evaluation of rabbit serum apo A-I
Leroy A., Olivier P., Vu-Dac N., Fruchart J.C. 85
The determination of plasma oxalate concentrations using an enzyme/bioluminescent assay. 2. Co-immobilisation of bioluminescent enzymes and studies of in vitro oxalogenesis
Parkinson I.S., Channon S.M., Tomson C.R.V., Adonai L.R., Ward M.K., Laker M.F. 97
Variation in alpha-1-antitrypsin phenotypes associated with penicillamine therapy
Whitehouse D.B., Lovegrove J.U., Hopkinson D.A. 109

Vol. 179/2, 1989

Effect of fish oil supplementation on erythrocyte lipid pattern, malondialdehyde production and glutathione-peroxidase activity in psoriasis
Corrocher R., Ferrari S., De Gironcoli M., Bassi A., Olivieri O., Guarini P., Stanzial A., Barba A.L., Gregolini L. 121
Human plasma contains low molecular weight factors which stimulate active sodium transport in erythrocytes
Tepper T., Huisman R.M., Jilderda J.F., Van der Hem G.K., De Zeeuw D. 133
α-Fetoprotein antibody-lectin enzyme immunoassay to characterize sugar chains for the study of liver diseases
Kinoshita N., Suzuki S., Matsuda Y., Taniguchi N. 143
Severity of GM1 gangliosidosis and urinary oligosaccharide excretion
Takahashi Y., Orii T. 153
Racial effect on serum creatine-kinase: Implications for estimation of heterozygosity risks for females at-risk for Duchenne dystrophy
Passos-Bueno M.R., Rabbi-Bortolini E., Azevedo E., Zatz M. 163
Sandwich enzyme immunoassay for serum amyloid A protein (SAA)
Yamada T., Uchiyama K., Yakata M., Gejyo F. 169
A sensitive and specific chemiluminescence method for estimating the ability of human granulocytes and monocytes to generate O_2^-
Nishida A., Kimura H., Nakano M., Goto T. 177
Isolation of apolipoprotein A-I-containing lipoproteins by immunoaffinity chromatography using a formylated derivative of cellulose gel
Ohta T., Hattori S., Murakami M., Horiuchi S., Nishiyama S., Matsuda I. 183
Determination of tellurium in biological fluids by means of electrothermal vaporization-inductively coupled plasma-mass spectrometry (ETV-ICP-MS)
Newman R.A., Osborn S., Siddik Z.H. 191
Insulin binding to erythrocytes and fatty acid composition of erythrocyte membrane phospholipids in healthy men
Pelikanova T., Kohout M., Hilgertova J., Rase J., Farska I., Valek J. 197
Carboxyl terminal glycine extended progastrin (gastrin-G) in human gastric mucosa and gastrinomas
Azuma T., Inokuchi H., Kawai K., Fujimoto S., Nakajima M., Taggart R.T., Walsh J.H. 201
Role of variant prealbumin in the pathogenesis of familial amyloidotic polyneuropathy: Fate of normal and variant prealbumin in the circulation of rats
Ando Y., Ikegawa S., Miyazaki A., Inoue M., Morino Y., Araki S. 205
Serum and ascitic fluid concentrations of apolipoproteins A and B in alcoholic cirrhosis patients
Nayak S.S., Bhat K.R., Nair A., Rao S., Aroor A.R. 207

Vol. 179/3, 1989

Diagnosis of subtypes of GM1 gangliosidosis in vitro and in vivo — Using urinary oligosaccharides as substrates
Takahashi Y., Orii T. 219
Improved competitive enzyme-linked immunoassay (ELISA) for albuminuria
Neumann R.G., Cohen M.P. 229
Fibrinogen sevilla, a congenital dysfibrinogenemia characterized by an abnormal monomer aggregation and a defective plasmin lysis
Fernandez F.J., Noguerol P., Sosa R., Cuesta B., Paramo J.A., Rocha E. 239
Assessment of antithyroglobulin and antimicrosomal autoantibodies in patients with autoimmune thyroid disease: Comparison of haemagglutination assay, enzyme-linked immunoassay and radioligand assay
Engler H., Staub J.-J., Althaus B., Ryff-deLeche A., Gerber H. 251
Application of solid-phase antibodies to radioimmunoassay. Improved performance of Dynospheres® following affinity purification of antisera
McConway M.G., Chapman R.S. 265
Relationship between hair, serum and bone aluminium in hemodialyzed patients
Chappuis P., De Vernejoul M.-C., Paolaggi F., Rousselet F. 271
The subunit composition of factor XIII proteins in normal and factor XIII deficient plasma and serum analysed by line immunoelectrophoresis
Kroll J. 279
Changes in the immunoreactivity of Apo A-I during storage
Wang X.L., Dudman N.P.B., Blades B.L., Wilcken D.E.L. 285
Erythrocyte and plasma cholesterol exchange in sickel cell anemia
Ngogang J., Mouray H., Lebreton de Vonne T., Raisonnier A. 295
Microbial influences on urinary polyamine excretion
Satink H.P.W.M., Hessels J., Kingma A.W., Van Den Berg G.A., Muskiet F.A.J., Halie M.R. 305
Procollagen III peptides and fibronectin in alcohol-related chronic liver disease: Correlations with morphological features and biochemical tests
Gabrielli G.B., Faccioli G., Casaril M., Capra F., Bonazzi L., Falezza G., Tomba A., Barcchino F., Corrocher R. 315
Kinetics of serum glutamate dehydrogenase during short-term alcohol withdrawal
Weill J., Schellenberg F., Le Goff A.-M. 323
The lysosomal β-D-N-acetylglucosaminidase isozymes in human plasma during pregnancy: Separation and quantification by a simple automated procedure
Goi G., Fabi A., Lombardo A., Bairati C., Bovati L., Burlina A.B., Agosti S., Serio C., Tettamanti G. 327
Retention of antimony in hair during leishmaniasis treatment
Dorea J.G., Merchan-Hamann E., Ryan D.E., Holzbecher J. 341

Vol. 180/1, 1989

Monoclonal gammopathies — Their identification and biological significance
Osterland C.K. 1
Inherited occurrence of a heat stable alkaline phosphatase in the absence of malignant disease
Onica D., Rosendahl K., Waldenlind L. 23
Lipid lowering actions of 7-(2-methylene butyryl)-(2H)-1,4-benzoxazin-3-(4H)-one derivatives in mice, rats and Syrian hamsters
Moussavi Z., Plancke M.O., Olivier Ph., Lesieur D., Fruchart J.C., Sauzieres J. 35
Clinical significance of mucin-like high molecular weight glycoprotein originated from lung cancer as tumor marker
Imokawa M., Okabe H., Ochi Y., Kajita Y. 45
Identification of an extra lactate dehydrogenase band in human erytrocytes
Hirano T., Matsuzaki H., Sekine T. 59

Characteristics and partial purification of prolidase and prolinase from leukocytes of a normal human and a patient with prolidase deficiency
Kodama H., Ohhashi T., Ohba Ch., Ohno T., Arata J., Kubonishi I., Miyoshi I. 65
Development of an RIA-based primary screen for IL-1 antagonists
Maloff B.L., Shaw J.E., Di Meo T.M., Fox D., Bruin E.M. 73
Determination of the biological activity of antithrombin III related antigen in urine
Ueshima S., Yamamoto T., Okada K., Tanaka H., Matsuo O. 79
Comparison of unitages for total free erythrocyte protoporphyrin (total EP) in anemia screening
Harada K., Omori S., Ueno T., Misumi J., Miura H. 87
Direct measurement of whole blood glucose by a needle-type sensor
Yamasaki Y., Ueda N., Nao K., Sekiya M., Kawamori R., Shichiri M., Kamada T. 93
Aspartylglucosamine excretion in heterozygous carriers of aspartylglycosaminuria
Mononen T.K. 99
Comparative study: The antioxidant enzymes and lipid peroxidation in cord and maternal red blood cell haemolysates
Novak Z., Kovacs L., Pal A., Pataki L., Varga Sz. I., Matkovics B. 103

Vol. 180/2, 1989

Plasma lipoproteins and adrenocortical hormones in men — Positive association of low density lipoprotein cholesterol with plasma cortisol concentration
Nanjee M.N., Miller N.E. 113
Glycated hemoglobin in hypoglycemia
Niketic V., Djurdjic V., Stojkovic V., Maric S., Djordjevic P. 121
Single step affinity chromatographic purification of human α-amylase from aspirated duodenal juice and its application in the measurement of pancreatic α-amylase synthesis rates in man
Ogden J.M., O'Keefe S.J.D., Ehlers M.R.W., Kirsch R.E., Marks I.N. 129
Amniotic fluid phospholipids. Study of the cold acetone precipitation effect
Budria R.J., Gonzalez F.E., Sanz H.M.T., Ostariz S.J.L. 141
Variability in lipoprotein concentrations in serum after prolonged storage at -20°C
Tiedink H.G.M., Katan M.B. 147
Measurement of 25-hydroxyvitamin D_2, 25-hydroxyvitamin D_3, 24,25-dihydroxyvitamin D_2 and 25,26-dihydroxyvitamin D_2 in a single plasma sample by mass fragmentography
Coldwell R.D., Trafford D.J.H., Varley M.J., Kirk D.N., Makin H.L.J. 157
Metabolism of pyrazinamide and allopurinol in hereditary xanthine oxidase deficiency
Yamamoto T., Higashino K., Kono N., Kawachi M., Nanahoshi M., Takahashi S., Suda M., Hada T. 169
Differences between human liver- and bone-type alkaline phosphatases
Miura M., Matsuzaki H., Bailyes E.M., Koyama I., Sakagishi Y., Sekine T., Komoda T. 177

Vol. 180/3, 1989

The simultaneous determination of oxidized and reduced glutathiones in liver tissue by ion pairing reverse phase high performance liquid chromatography with a coulometric electrochemical detector
Harvey P.R.C., Ilson R.G., Strasberg S.M. 203
Binding of glycosaminoglycan inhibitors to calcium oxalate crystals in relation to ionic strength
Fellstrom B., Lindsjo M., Danielson B.G., Karlsson F.A., Ljunghall S. 213
Renin substrate in human amniotic fluid
Metsarinne K., Fyhrquist F., Rosenlof K., Autero M., Pekonen F., Rutanen E.-M. 221
Effect of drugs on histamine radio-enzyme assay
Harvima R.J., Harvima I.T., Kajander E.O., Penttila I.M., Horsmanheimo M., Fraki J.E. 231
The effect of folic acid and 5-formyltetrahydrofolic acid on folate-deficient HEp 2 cells in culture
Leeming R.J., Hamon C.G.B., Tomkyns J.G. 241

Circadian and seasonal variations of electrolytes in aging humans
Touitou Y., Touitou C., Bogdan A., Reinberg A., Motohashi Y., Auzeby A., Beck H. 245
Correction of subclinical ascorbate deficiency in patients receiving dialysis: Effects on plasma oxalate, serum cholesterol, and capillary fragility
Tomson C.R.V., Channon S.M., Parkinson I.S., McArdle P., Qureshi M., Ward M.K., Laker M.F. 255
Interactions with hemoglobin: A source of error in measurements of transketolase activity in hemolysates
Grudzinski A., Waltham M., Ioannoni B., Price J., Nixon P.F. 265
A multiwell lectin-binding assay using Lotus tetragonolobus for measuring different glycosylated forms of haptoglobin
Thompson S., Stappenbeck R., Turner G.A. 277
A two-site immunoradiometric assay for C-reactive protein in serum
Shapiro D., Shenkin A. 285
Detection and identification of the plant lignans lariciresinol, isolariciresinol and seco-isolariciresinol in human urine
Bannwart C., Adlercreutz H., Wahala K., Brunow G., Hase T. 293
α-L-Fucosidase activity in normal human lymphocytes
Roger L., Bernard M.-A., Percheron F., Foglietti M.-J. 303

Vol. 181/1, 1989

Isolation of creatine kinase BB isoenzyme with high specific activity and adequate purity for radioimmunoassay from human placenta on preparative polyacrylamide gel electrophoresis
Geng J.-G., Cheng H.-Z., Yang Y.-F., Qian Z.-H., Jiang C.-Y. 1
Alterations of acidic phospholipids in bronchoalveolar lavage fluids of patients with pulmonary alveolar proteinosis
Honda Y., Kataoka K., Hayashi H., Takahashi H., Suzuki A., Akino T. 11
An enzyme immunoassay of slow moving protease (SMP) in human gastric mucosa
Matsusako K., Itoh M., Joh T., Miyamoto T., Yokochi K., Takeuchi T., Moriyama A., Kato T., Tanaka R. 19
Sensitive enzyme immunoassay for human aldolase B
Haimoto H., Kurobe N., Hosoda S., Kato K. 27
Mechanism of inhibition of erythrocyte glutathione reductase by Mitomycin-C
Okuda J., Hirai Y., Hayazaki T. 37
Glycan microheterogeneity of α_1-acid glycoprotein in sera and dialysates of patients on continuous ambulatory peritoneal dialysis
Fraeyman N.H., Van de Velde E.J., Belpaire F.M., Lameire N.H. 47
Increased phosphoinositide breakdown by phospholipase C in erythrocyte membranes from patients with cystic fibrosis
Doughney C., McPherson M.A., Goodchild M.C., Dormer R.L. 55
Aberrant free radical activity in cystic fibrosis
Salh B., Webb K., Guyan P.M., Day J.P., Wickens D., Griffin J., Braganza J.M., Dormandy T.L. 65
Uptake of trihalomethanes by patients during hemodialysis
Cailleux A., Subra J.F., Riberi P., Allain P. 75
The serum activities of AP, gamma-GT, GLDH, GPT and CHE after complete biliary obstruction and choledochocaval fistula in the rat
Putzki H., Reichert B., Heymann H. 81
Serum unconjugated bile acids in patients with small bowel bacterial overgrowth
Bolt M.J.G., Stellaard F., Sitrin M.D., Paumgartner G. 87
Metabolism of androstenedione by placental microsomes in pregnancy hypertension
Hahnel M.E., Martin J.D., Michael C.A., Hahnel R. 103
Reversed-phase high-performance liquid-chromatography of 2,8-dihydroxyadenine in serum and urine with electrochemical detection
Kojima T., Nishina T., Kitamura M., Kamatani N., Nishioka K. 109

Vol. 181/2, 1989

A new case with hereditary xanthinuria: Response to exercise
Landaas S., Borch K., Aagaard E. 119
Biochemical diagnosis of Hunter syndrome on Epstein-Barr virus-transformed lymphoblastoid cell lines
Morabito E., Giambarrasi I., Rocchi M., Di Natale P. 125
Comparison between the chromate inhibition test and a cytochemical method for the determination of glucose-6-phosphate dehydrogenase deficiency in erythrocytes
Jonges G.N., Hagen H., Van Noorden C.J.F., Weening R.S., Roos D. 135
Measurement of plasma uroporphyrin during treatment and long-term follow-up of patients with Porphyria cutanea tarda
Adjarov D.G., Kerimova M.A. 143
Concentration and physicochemical characterisation of unsaturated cobalamin binding proteins in amniotic fluid
Gueant J.L., Hambaba L., Vidailhet M., Schaefer C., Wahlstedt V., Nicolas J.P. 151
Patients with hereditary fructose intolerance have normal erythrocyte aldolase activity
Gitzelmann R., Steinmann B., Tuchschmid 163
Phosphatidylcholine storage in mucolipidosis IV
Bargal R., Bach G. 167
Carnitine deficiency associated with anticonvulsant therapy
Rodriguez-Segade S., Alonso de la Pena C., Tutor J.C., Paz J.M., Fernandez M.P., Rozas I., Del Rio R. 175
Plasma and serum lactoferrin levels in cystic fibrosis. Relationship with the presence of cystic fibrosis protein
Barthe C., Galabert C., Guy-Crotte O., Figarella C. 183
Determination of $NADH_2$-ferricyanide oxidoreductase (cytochrome b_5 reductase, diaphorase) activity of human erythrocytes by an analysis of the time dependence of $NADH_2$ oxidation
Papandreou P., Rakitzis E.T. 189
Correlations among blood levels of renin-angiotensin-aldosterone, bradykinin, and catecholamines in healthy subjects
Yamamoto T., Hiromoto J., Kikuchi Y., Oka M., Takatori O. 197
Two new serum immunoassay (RIA and EIA) for the determination of M_2 antimitochondrial antibodies in primary biliary cirrhosis
Plebani M., Chiaramonte M., Floreani A., Bertorelle R., Naccarato R., Burlina A. 201
An improved spectrophotometric bromide assay for the estimation of extracellular water volume
Trapp S.A., Bell E.F. 207
Simultaneous assay of zinc protoporphyrin and protoporphyrin IX in dried blood
Orfanos A.P., Guthrie R., Jinks D.C., Vollmer D.W. 213
A micromethod for the determination of fructose in human erythrocytes and plasma
Liu S.-Q., Das B., Srivastava S.K. 219
Nonspecific reaction in the sandwich immunoassay for human tumor necrosis factor-α (hTNF-α)
Abe Y., Miyake M., Miyazaki T., Horiuchi A., Sagawa T., Ono H., Kimura S. 223
Analysis of acylcarnitines in normal human urine with the radioisotope exchange-high performance liquid chromatography (HPLC) method
Schmidt-Sommerfeld E., Penn D., Kerner J., Bieber L.L. 231

Vol. 181/3, 1989

Proteinuria in sickle cell trait and disease: An electrophoretic analysis
Lonsdorfer A., Comoe L., Yapo A.E., Lonsdorfer J. 239
Tetrahydrobiopterin metabolism in the streptozotocin induced diabetic state in rats
Hamon C.G., Cutler P., Blair J.A. 249
Pharmacokinetic study of butyric acid administered in vivo as sodium and arginine butyrate salts
Daniel P., Brazier M., Cerutti I., Pieri F., Tardivel I., Desmet G., Baillet J., Chany C. 255

An improved formula for the judgement of intrathecally produced IgG in the presence of blood brain barrier damage
Ohman S., Forsberg P., Nelson N., Vrethem M. 265
Hydroxycholesterols in serum from hypercholesterolaemic patients with an without bile acid sequestrant therapy
Van Doormaal J.J., Smit N., Koopman B.J., Van der Molen J.C., Wolthers B.G., Doorenbos H. 273
Serum human chorionic gonadotrophin levels in early pregnancy
Ooi D.S., Perkins S.L., Claman P., Muggah H.F. 281
One step sandwich enzyme immunoassay for human type IV collagen using monoclonal antibodies
Obata -i. K., Iwata K., Ichida T., Inoue K., Matsumoto E., Muragaki Y., Ooshima A. 293
Detection of elevated amounts of urinary pseudouridine in cancer patients by use of a monoclonal antibody
Itoh K., Mizugaki M., Ishida N. 305
Assay of serum hyaluronic acid in clinical application
Chichibu K., Matsuura T., Schichijo S., Yokoyama M.M. 317
The quantitation of human ventricular myosin light chain 1 in serum after myocardial necrosis and infarction
Wang J., Shi Q., Wu T.-W., Jackowski G., Mickle D.A.G. 325
A new approach for measuring the erythrocyte life span with a nonradioisotope
Dever M., Smith J.E., Hausler D.W. 337

Vol. 182/1, 1989

Radioisotopic assay for erythrocyte adenosine 5′-monophosphate deaminase
Dale G.L. 1
Carbohydrate removal fails to eliminate the heterogeneity of human prostatic acid phosphatase
Morris M.F., Waheed A., Risley J.M., Van Etten R.L. 9
Immunoglobulins from the sera of immunologically activated persons with pairs of electrophoretic restricted bands show a greater tendency to aggregate than normal
Levinson S.S., Keren D.F. 21
Inhibition of neutrophil and natural killer cell function by human seminal fluid acid phosphatase
Mukhopadhyay N.K., Saha A.K., Smith W., Dowling J.N., Hiserodt J., Glew R.H. 31
Effects of N-hydroxyethyl-1-deoxynojirimycin (BAY m 1099) on the activity of neutral- and acid α-glucosidases in human fibroblasts and HepG2 cells
Wisselaar H.A., Van Dongen J.M., Reuser A.J.J. 41
Simultaneous determination of hemoglobin and coproporphyrin by second derivative differential spectrophotometry: Application to the diagnosis of meconium aspiration
Ficheux H., Levillain P., Francoual J., Leluc R., Lemonnier A. 53
Adenylate degradation products release from the human myocardium during open heart surgery
Smolenski R.T., Skladanowski A.C., Perko M., Zydowo M.M. 63
Binding of fibronectin to human lipoproteins
Cseh K., Karadi I., Rischak K., Szollar L., Janoki G., Jakab L., Romics L. 75
A spectrophotometric method for the determination of free and esterified carnitine
Schafer J., Reichmann H. 87
Simple enzymatic screening assay for ethylene glycol (ethane-1,2-diol) in serum
Hansson P., Masson P. 95
Quantitative estimation of an α_2 pregnancy-associated protein in normal pregnancy by electro-immunoassay
Nakai M., Atsumi M., Sone Y., Arai M. 103
Elimination of free glycerol interference in a colorimetric enzymic triglyceride assay
Artiss J.D., Strandbergh D.R., Zak B. 109
New enzymatic endpoint assay of serum sialic acid
Horiuchi T., Kurokawa T. 117

Vol. 182/2, 1989

The effect of hypoalbuminaemia, hyperbilirubinaemia and renal failure on serum fructosamine concentration in non-diabetic individuals
Dominiczak M.H., Orrell J.M., Finlay W.E.I. 123

Measurement of glomerular filtration rate in children with insulin-dependent diabetes mellitus
Gibb D.M., Dalton N.R., Barratt M.T. 131

Measurement of propionate turnover in vivo using sodium [$^{2}H_{5}$]propionate and sodium [^{13}C]propionate
Walter J.H., Thompson G.N., Leonard J.V., Heatherington C.S., Bartlett K. 141

Detection of point mutations in amplified single copy genes by biotin-labelled oligonucleotides: Diagnosis of variants of alpha-1-antitrypsin
Gregersen N., Winter V., Petersen K.B., Koch J., Kolvraa S., Rudiger N., Heinsvig E.-M., Bolund L. 151

Red cell membrane phosphatidylinositol kinase activity in hemolytic anemias and myeloproliferative diseases
Boivin P., Galand C. 165

Evidence that free gamma carboxyglutamic acid circulates in serum
Fournier B., Gineyts E., Delmas P.D. 173

Serum CK-BB as a tumor marker in patients with carcinoma confirmed histologically
Arenas J., Diaz A.E., Alcaide M.J., Santos I., Martinez A., Culebras J.M. 183

Plasma protein S in disseminated intravascular coagulation, liver disease, collagen disease, diabetes mellitus, and under oral anticoagulant therapy
Takahashi H., Takewaki W., Wada K., Shibata A. 195

Human serum immuno-reactive copper, zinc-superoxide dismutase assayed with an enzyme monoclonal immunosorbent in patients with digestive cancer
Oka S., Ogino K., Matsuura S., Yoshimura S., Yamamoto K., Okazaki Y., Takemoto T., Kato N., Uda T. 209

Serum selenium in liver cirrhosis: Correlation with markers of fibrosis
Casaril M., Stanzial A.M., Gabrielli G.B., Capra F., Zenari L., Galassini S., Moschini G., Liu N.Q., Corrocher R. 221

Measurement of free kappa immunoglobulin light chains in the cerebrospinal fluid by a competitive avidin-biotin ELISA
Lolli F., Amaducci L. 229

Vol. 182/3, 1989

The effects of Oradexone and Ambroxol pretreatment on the oxidative sensitivity of the red blood cells in preterm infants
Novak Z., Varga S.I., Kovacs L., Pal A., Pataki L., Matkovics B., 241

A simple, rapid assay for plasma oxalate in uraemic patients using oxalate oxidase, which is free from vitamin C interference
Rolton H.A., McConnell K.N., Modi K.S., Macdougall A.I. 247

Binding of calcium by organic anions, determined by perturbation of the equilibrium solubility of [^{14}C]calcium oxalate
Geller D.A., Ostrow J.D., Moore E.W., Celic L., Nancollas G.H. 255

Reference values for amino acids in cerebrospinal fluid of children determined using ion-exchange chromatography with fluorimetric detection
Gerrits G.P., Trijbels F.J., Monnens L.A., Gabreels F.J., De Abreu R.A., Theeuwes A.G., Raay-Selten B. 271

N-acetyl-β-glucosaminidase and albuminuria in normal and diabetic pregnancies
Skrha J., Perusicova J., Sperl M., Bendl J., Stolba P. 281

Identification of octadeca-9,11-dienoic acid and its significance in the diagnosis of cervical intraepithelial neoplasia
Parham D.M., Ross P.E., Gunstone F.D., Robertson A.J., Duncan I.D. 289

Application of two monoclonal antibodies to either an immunosorbent enzyme assay or a competitive binding enzyme immunoassay for human serum pancreatic lipase
Ohkaru Y., Kurooka S., Kitamura T. 295
Sickle cell screening using a rapid automated HPLC system
Shapira E., Miller V.L., Miller J.B., Qu Y. 301

Vol. 183/1, 1989

Carnitine: Metabolism and clinical chemistry
Siliprandi N., Sartorelli L., Ciman M., Di Lisa F. 3
Clinical biochemistry of neuron specific enolase
Kaiser E., Kuzmits R., Pregant P., Burghuber O., Worofka W. 13
The importance of objective data in the diagnosis of pancreatitis
Lott J.A., Ellison E.C., Applegate D. 33
Pancreatic α-amylase assay employing the synergism of two monoclonal antibodies
Rauscher E., Gerber M. 41
Monoclonal antibodies to enzymes
Rosalki S.B. 45
Serum phospholipase A in hospitalized patients
Hoffmann G.E., Hiefinger R., Steinbrueckner B. 59
Increase of creatine kinase and lactate dehydrogenase in the serum of rats submitted to experimental intestinal infarction
Roth M., Jaquet P.-Y., Rohner A. 65
Serum arylesterase (paraoxonase) activity following myocardial infarction
Secchiero S., Mussap M., Zaninotto M., Bertorelle R., Burlina A. 71
Assay of HMG-CoA reductase activity in the evaluation of cholesterol synthesis in man
Carulli N., Tripodi A., Carubbi F. 77
Glutathione-related enzyme activities in testis of patients with malignant diseases
Aceto A., Di Ilio C., Angelucci S., Tenaglia R., Zezza A., Caccuri A.M., Federici G. 83
Methods for serum cholinesterase assay and classification of genetic variants
Panteghini M. 87
Caffeine as indicator of metabolic functions of microsomal liver enzymes
Varagnolo M., Plebani M., Mussap M., Nemetz L., Paleari C.D., Burlina A. 91
Evaluation of pancreatic amylase immunoassay in acute pancreatitis
Sacchetti L., Cavalcanti E., Cerasuolo D., Oriani G., Uomo G., Rabitti P.G., Laccetti M., Visconti M., Salvatore F. 95
Assays and structure of reverse transcriptase of the AIDS virus
Bertazzoni U., Lori F., Achilli G., Cattaneo E. 101

Vol. 183/2, 1989

Calcium absorption in children estimated from single and double stable calcium isotope techniques
Miller J.Z., Smith D.L., Flora L., Peacock M., Johnston Jr. C.C. 107
Time resolved immunofluorometric assay for placental protein 14
Riitinen L., Stenman U.-H., Alfthan H., Suikkari A.-M., Bohn H., Seppala M. 115
Detection of kinetically abnormal argininosuccinate synthase in neonatal citrullinemia by conversion of citrulline to arginine in intact fibroblasts
Ben-Yoseph Y., Mitchell D.A. 125
Human cerebrospinal fluid acetylcholinesterase and butyrylcholinesterase. Evidence for identity between the serum and cerebrospinal fluid butyrylcholinesterase
Rao R.V., Gnanamuthu C., Balasubramanian A.S. 135
Fluorometric assay of protein in native human bile
Harvey P.R.C., Upadhya G.A., Toth J.L., Strasberg S.M. 147

Immunoassay of β-hexosaminidase isoenzymes in serum in patients with raised total activities
Isaksson A., Hultberg B. 155
Automated enzymatic assays for the determination of intestinal permeability probes in urine. 2. Mannitol
Lunn P.G., Northrop C.A., Northrop A.J. 163
A convenient enzymatic method for the determination of 4-methyl-2-oxopentanoate in plasma: comparison with high performance liquid chromatographic analysis
Schadewaldt P., Hummel W., Trautvetter U., Wendel U. 171
A rapid, qualitative ELISA test for the specific detection of morphine in serum or urine
Laurie D., Manson A.J., Rowell F.J., Seviour J. 183
A competitive microtitre plate enzyme immunoassay for plasma testosterone using polyclonal anti-testosterone immunoglobulins
Rajkowski K.M., Hanquez Ch., Bouzoumou A., Cittanova N. 197
An improved rapid thin-layer chromatographic-gas-liquid chromatographic procedure for the determination of free fatty acids in plasma
Tichelaar H.Y., Spinnler Benade A.J., Daubitzer A.K., Kotze T.J.V.W. 207
Salivary corticosteroids in the study of adrenal function
Bolufer P., Gandia A., Rodriguez A., Antonio P. 217
ELISA for measuring porphobilinogen deaminase in human erythrocytes
Lannfelt L., Wetterberg L., Lilius L., Thunell S., Gellerfors P. 227

Vol. 183/3, 1989

Markers of collagen metabolism in sera of patients with various rheumatic diseases
Myllyla R., Becvar R., Adam M., Kivirikko K.I. 243
Serum lipids and cholesterol metabolism during guar gum, plantago ovata and high fibre treatments
Miettinen T.A., Tarpila S. 253
Increased furosemide-sensitive sodium influx into erythrocytes in non-insulin-dependent diabetes mellitus associated with hypertension
Chimori K., Miyazaki S., Goi R., Dodo S., Hayashi H., Izai M., Goto S., Yasuda K., Miura K. 263
Partial purification and characterization of human hepatoma-derived growth factor
Nakamura H., Kambe H., Egawa T., Kimura Y., ito H., Hayashi E., Yamamoto H., Sato J., Kishimoto S. 273
IgA isotyping of antigliadin antibodies. A possible clue for a less invasive diagnosis of coeliac disease
Elewaut A., Dacremont G., Robberecht E., Leroy J., De Baets M.H. 285
CSF-IgM measurement in neurovenereological disease
Van Kamp G.J., Wolters Ch. E. 295
Glycosyltransferase activities in normal and leukaemic monocytic cells
Francina A., Gateau-Roesch O., Couprie N., Leculier C., Col J.-F., Archimbaud E., Campos L., Louisot P., Richard M. 301
Selective determination of hydroxyproline in urine by high-performance liquid chromatography using precolumn derivatization
Teerlink T., Tavenier P., Netelenbos J.C. 309
Small bowel biopsy for disaccharidase levels: Evidence that endoscopic forceps biopsy can replace the Crosby capsule
Smith J.A., Mayberry J.F., Ansell I.D., Long R.G. 317
Plasminogen activator activity and plasminogen independent amidolytic activity in tear fluid from healthy persons and patients with anterior segment inflammation
Tozser J., Berta A., Punyiczki M. 323
Alterations of purine metabolism in mononuclear cells of individuals at risk of developing type I (insulin-dependent) diabetes mellitus
Schafer J.H., Jira T.B., Michaelis D., Rjasanowski I.M. 333

Insulin and glucagon degradation in liver are not affected by hepatic cirrhosis
Antoniello S., La Rocca S., Calvacanti E., Auletta M., Salvatore F., Cacciatore L. 343
Increased acylphosphatase levels in erythrocytes from hyperthyroid patients
Nassi P., Liguri G., Nediani C., Taddei N., Piccinni P., Degl'Innocenti D., Gheri R.G., Ramponi G. 351
Immunodepletion of albumin from human serum samples
Burgess-Cassler A., Johansen J.J., Kendrick N. 359

Vol. 184/1, 1989

A high performance liquid chromatography method for the analysis of ^{35}S-cystine: Application to the diagnosis of cystinosis
Hall N.A., Young E.P. 1
Human fibrinogen heterogeneity. A study of limited fibrinogen degradation
Reganon E., Vila V., Aznar J., Laiz B. 7
Purification and characterization of endometrial protein PP14 from mid-trimester amniotic fluid
Riittinen L., Julkunen M., Seppala M., Koistinen R., Huhtala M.-L. 19
Immunoreactive anionic and cationic trypsin in human serum
Kimland M., Russick C., Marks W.H., Borgstrom A. 31
Analysis of maple syrup urine disease in cell culture: Use of substrates
Schadewaldt P., Beck K., Wendel U. 47
3-Methylglutaconyl-CoA hydratase, 3-methylcrotonyl-CoA carboxylase and 3-hydroxy-3-methylglutaryl-CoA lyase deficiencies: A coupled enzyme assay useful for their detection
Narisawa K., Gibson K.M., Sweetman L., Nyhan W.L. 57
Radioimmunoassay of human platelet-derived growth factor using monoclonal antibody toward a synthetic 73-97 fragments of its B-chain
Shiraishi T., Morimoto S., Itoh K., Sato H., Sugihara K., Onishi T., Ogihara T. 65
The sugar-chain heterogeneity of human γ-glutamyl transferases from the reproductive system and kidney
Arai K., Yoshida K.-I., Komoda T., Kobayashi N., Saitoh H., Sakagishi Y. 75
Serum F protein: A new sensitive and specific test of hepatocellular damage
Foster G.R., Goldin R.D., Oliveira D.B.G. 85
3-Methylisoxazol-5-one, an artefact from acetoacetic acid formed during urinary organic acid analysis
Mills G.A., Walker V., Mellor J.M. 93
Removal of bilirubin from jaundiced serum by immunoaffinity chromatography
Izumi Y., Nakajima H. 99
Serum lipolytic activity in Reye's syndrome
Kang E.S., Matsuo N., Nagai T., Greenhaw J., Williams P.L. 107
Serum deoxyribonuclease I determined by a radioimmunoassay and an enzymatic assay in malignant diseases
Miyauchi K., Ogawa M., Murata A., Nakano I., Funakoshi A., Mori T. 115

Vol. 184/2, 1989

Con A affinity of rat α1-acid glycoprotein (rAGP): Changes during inflammation, dexamethasone or phenobarbital treatment as detected by crossed affino immunoelectrophoresis (CAIE) are not only a reflection of biantennary glycan content
Pos O., Drechou A., Durand G., Bierhuizen M.F.A., Van der Stelt M.E., Van Dijk W. 121
Fragmented fibronectin and other synovial fluid proteins in chronic arthritis: Their relation to immune complexes
Griffiths A.M., Herbert K.E., Perrett D., Scott D.L. 133
The effect of clomipramine treatment on plasma lipoproteins and high density lipoprotein subfractions in healthy subjects
Skinner E.R., Watt C., Reid I.C., Besson J.A.O., Ashcroft G.W. 147

Oxalate, phosphate and sulphate determination in serum and urine by ion chromatography
Politi L., Chiaraluce R., Consalvi V., Cerulli N., Scandurra R. 155
Glucocorticoid receptors in monocytes in Type 1 diabetes mellitus
Damm P., Binder C. 167
Pyrimidine 5'-nucleotidase deficiency: Improved detection of carriers
De Korte D., Sijstermans J.M., Seip M., Van Doorn C.C.H., Van Gennip A.H., Roos D. 175
Plasma lipoproteins in infantile visceral Leishmaniasis: Deficiency of apolipoproteins A-I and A-II
Bekaert E.D., Kallel R., Bouma M.-E., Lontie J.-F., Mebazaa A., Malmendier C.L., Ayrault-Jarrier M. 181
Effect of renal failure on plasma dihydroxyphenylglycol, 3-methoxy-4-hydroxyphenylglycol, and vanillylmandelic acid
Hoeldtke R.D., Israel B.C., Cavanaugh S.T., Krishna G.G. 195

Vol. 184/3, 1989

Development of an immunoassay for all human isoferritins, and its application to serum ferritin evaluation
Cozzi A., Levi S., Bazzigaluppi E., Ruggeri G., Arosio P. 197
Amplification of the proto-oncogenes int-2, c-erb B-2 and c-myc in human breast cancer
Machotka S.V., Garrett C.T., Schwartz A.M., Callahan R. 207
The estimation of total serum lipids by a completely enzymatic 'summation' method
Akins J.R., Waldrep K., Bernert Jr. J.T. 219
Rapid homogeneous enzyme immunoassay of plasma protein C
Suzuki K., Hayashi T., Koyama M., Yoshimura T., Shimamoto M., Kimura N., Kita M. 227
Use of 4-heptylumbelliferyl-β-D-glucoside to identify Gaucher's disease heterozygotes
Butcher B.A., Gopalan V., Lee R.E., Richards T.C., Waggoner A.S., Glew R.H. 235
Determination of succinylacetone and succinylacetoacetate in physiological samples as the common product 5(3)-methyl-3(5)-isoxazole propionic acid using an isotope dilution method and mass spectrometry
Schierbeek H., Berger R. 243
Abnormally-fucosylated serum haptoglobins in patients with inflammatory joint disease
Thompson S., Kelly C.A., Griffiths I.D., Turner G.A. 251
Effect of ischaemia on the activities of human hepatic acyl-CoA: cholesterol acyltransferase and other microsomal enzymes
Smith J.L., Pillay S.P., De Jersey J., Hardie I.R. 259
Glutathione S-transferase in human bile
Howie A.F., Hayes P.C., Bouchier I.A.D., Hayes J.D., Beckett G.J. 269
Determination of bone Gla protein (osteocalcin) by enzyme-linked immunosorbent assay
Egsmose C., Daugaard H., Lund B. 279
Enhanced net mass transfer of HDL cholesteryl esters to Apo B-containing lipoproteins in patients with peripheral vascular disease
Ruhling K., Zabel-Langhennig R., Till U., Thielmann K. 289
Time course of serum amyloid A response in myocardial infarction
Bausserman L.L., Sadaniantz A., Saritelli A.L., Martin V.L., Nugent A.M., Sady S.P., Herbert P.N. 297
Assay of methylmalonyl CoA mutase with high-performance liquid chromatography
Kikuchi M., Hanamizu H., Narisawa K., Tada K. 307
Enzyme-labelled immunoassay for urinary gonadotropins using penicillinase
Desai M., Khatkhatay I., Sankolli G., Meherji P., Joshi U. 315
Effect of low dose acetylsalicylic acid upon plasma thromboxane B_2 levels and platelet aggregation in ischemic stroke patients
Lee T.-K., Chen Y.-C., Kuo T.-L., Lien I.-N., Liu M.-C., Huang Z.-S. 323

Vol. 185/1, 1989

Carbon dioxide tensions in physiological salt solutions: direct measurements
Abrahamsen J., Norrie B., Andersen P.K., Stokke D.B., Nedergaard O.A. 1
A sensitive, direct colorimetric assay of serum copper using 5-Br-PSAA
Makino T. 7
Bioluminescent assay for serum adenosine deaminase with immobilized bacterial luciferase
Oda K., Yoshida S., Hirose S., Takeda T. 17
The relationship of plasma acetate with glucose and other blood intermediary metabolites in non-diabetic and diabetic subjects
Akanji A.O., Humphreys S., Thursfield V., Hockaday T.D.R. 25
Serum alkaline DNase activity in normal or nonhospitalised individuals
Avall-Lundqvist E., Economidou-Karaoglou A., Sjovall K., Lans M., Taper H.S., Roberfroid M., Eneroth P. 35
A preliminary study of serum class II levels in healthy individuals and bone-marrow transplant patients
Thompson S., Wareham M., Pearson A.D.J., Sviland L., Turner G.A. 45
Intraerythrocytic water relaxation rates in amyotrophic lateral sclerosis. An NMR investigation
Monti J.P., Braguer D., Guelton C., Crevat A., Serratrice G. 53
Radioimmunoassay of neuropeptide Y (NPY) in biological fluids. Circulating NRY levels depend on renal function
Stridsberg M., Lundqvist G. 61
Direct extraction and assay of collagenase from human osteoarthritic articular cartilage
Blanckaert A., Mazieres B., Eeckhout Y., Vaes G. 73
Elastase binding capacity of α_2-macroglobulin in plasma of patients with asthma or chronic obstructive pulmonary disease, without α_1-protease inhibitor deficiency
Kilroe-Smith T.A., Dowdeswell R.J., Gaillard M.C. 81
Biochemical and neurophysiological parameters in hemodialyzed patients with chronic renal failure
Schoots A.C., De Vries P.M.J.M., Thiemann R., Hazejager W.A., Visser S.L., Oe P.L. 91
The effect of vitamin E therapy on plasma and erythrocyte lipid peroxidation in chronic hemodialysis patients
Yalcin A.S., Yurtkuran M., Dilek K., Kilinc A., Taga Y., Emerk K. 109
Reactive oxygen species production in whole blood from AIDS patients
Guillard O., Sonnet J., Lauwerys R. 113
Oxidative hemolysis in erythrocytes with unstable hemoglobins
Falcioni G., Grelloni F., De Sanctis G., Bonfigli A.R., Pierani P., Felici L., Coppa G.V. 115

Vol. 185/2, 1989

Bound amino acids in uremic sera: Search for peptidic middle molecules by preparative polyacrylamide gel electrophoresis and high performance liquid chromatography
Kaplan B., Gotfried M., Cojocaru M., Pras M., Ravid M. 119
The quantitation of oxalate in amniotic fluid by ion-chromatography
Wandzilak T.R., Hanson F.W., Williams H.E. 131
Effect of 8-anilino-1-naphthalene sulfonic acid (ANS) on the interaction between thyroid hormone and anti-thyroid hormone antibodies
Matsuda M., Sakata S., Komaki T., Nakamura S., Kojima N., Takuno H., Miura K. 139
A rapid electrophoretic method for the detection of serum Lp(a) lipoprotein
Kawakami K., Tsukada A., Okubo M., Tsukada T., Kobayashi T., Yamada N., Murase T. 147
Evaluation and performance characteristics of a novel ELISA using monoclonal antibody to glycated albumin
Hud E., Cohen M.P. 157

Gas-liquid chromatographic method for routine detection of plasma sulfo-, gluco-, and free 3-methoxy-4-hydroxyphenylglycol
Piletz J.E., Halaris A. 165
Radioimmunoassay for erythropoietin using anti-recombinant erythropoietin antibody with high affinity
Matsubara K., Yoshimura T., Kamachi S., Fukushima M., Hino M., Morii H. 177
Lectin binding characteristics of a cholesterol nucleation promoting protein
Harvey P.R.C., Upadhya A., Toth J.L., Strasberg S.M. 185
Automated analysis of urinary hydroxyproline
Ho K.-C., Pang C.-P. 191
Microassay for amino acids in human liquid blood and dried blood
Ho J.W. 197
The effect of non-enzymatically glycosylated collagen on normal human platelets
Yarat A., Emerkli N. 203
A simplified procedure for the study of LDL kinetics
Samman S., Khosla P., Carroll K.K. 207
A combined lectin precipitation method for differentiating modified bone alkaline phosphatase in serum
Kuwana T., Sugita O., Okada M., Yakata M. 211
Binding of nitroblue tetrazolium to human serum albumin in a fructosamine test
Sakurai T., Tsuchiya S. 213
Unusual 1β-hydroxylated bile acids in children with a paucity in interlobular bile ducts
Kimura A., Mahara R., Tohma M., Ushijima K., Yuge K., Ono E., Yamashita F. 215
German Society for clinical chemistry: Workshop on the standardization of enzyme determinations in medicine
Schmidt E. S1
1989 Alkaline phosphatase symposium
Moss D. S2
11th National Congress of the Italian Society of clinical enzymology
Moss D. S3

Vol. 185/3, 1989

Preface
Okuda K. 221
Development and applications of ultrasensitive enzyme immunoassays for antigens and antibodies
Ishikawa E., Hashida S., Tanaka K., Kohno T. 223
A new homogenous enzyme immunoassay using recombinant enzyme fragments
Khanna P.L., Dworschack R.T., Manning W.B., Harris J.D. 231
Microbial enzymes for creatinine assay: A review
Shimizu S., Kim J.-M., Yamada H. 241
Diagnostic application of mitochondrial enzymes and isoenzymes
Schmidt F.W., Schmidt E. 253
Different patterns of leakage of cytosolic and mitochondrial enzymes
Kamiike W., Fujikawa M., Koseki M., Sumimura J., Miyata M., Kawashima Y., Wada H., Tagawa K. 265
Immunological and ultrastructural analysis of loss of tubular membrane-bound enzymes in patients with renal damage
Scherberich J.E. 271
Structure of the ornithine transcarbamylase (OTC) gene and DNA diagnosis of OTC deficiency
Matsuda I., Hata A., Matsuura T., Tsuzuki T., Shimada K. 283
Genetic analysis of familial amyloidotic polyneuropathy, an autosomal dominant disease
Sakaki Y., Sasaki H., Yoshioka K., Furuya H. 291
Laboratory and clinical features of lactate dehydrogenase subunit deficiencies
Maekawa M., Kanno T. 299

Phosphoinositide metabolism and oncogenes
Takenawa T., Fukami K. 309
Separation of hepatoma-associated γ-glutamyltransferase isoenzyme on cellulose acetate media with Triton X-100 and concanavalin A
Yoshikawa C., Shimojo N., Naka K., Okuda I., Ohkawa J. 317
N-Acetylglucosaminyltransferase III in human serum, and liver and hepatoma tissues: Increased activity in liver cirrhosis and hepatoma patients
Ishibashi K., Nishikawa A., Hayashi N., Kasahara A., Sato N., Fujii S., Kamada T., Taniguchi N. 325
Regulation of protein kinase C by sphingosine and lysosphingolipids
Hannun Y.A., Bell R.M. 333
Small molecular weight GTP-binding proteins and signal transduction
Yamamoto K., Tanimoto T., Kim S., Kikuchi A., Takai Y. 347
Pathogenic potentials of bacterial proteases
Maeda H., Molla A. 357

Vol. 186/1, 1989

A lack of coordination in the release of urinary lysosomal and brush border enzymes following renovascular surgery
Holmes R.P., Craddock G., Espeland M.A., Assimos D.G., Dean R.H. 1
Hyperhistaminemia in patients with histidinemia due to increased decarboxylation of histidine
Tanabe M., Sakura N. 11
Immunochemical characterization of corticosteroid-binding globulin in human bronchoalveolar fluid
Loric S., Egloff M., Domingo M., Lacronique J., Degrelle H. 19
Ornithine carbamoyltransferase deficiency. A new variant with subnormal enzyme activity
Rabier D., Benoit A., Petit F., Chekoury A., Bonnefont J.P., Saudubray J.M., Kamoun P. 25
Measurement of the unbound fraction of long chain nonesterified fatty acids (NEFA) in serum: Methodological considerations
Lim C.-F., Wynne K.N., Stockigt J.R. 31
Origin of serum catalase activity in acute pancreatitis
Goth L. 39
A rapid and sensitive method for estimating low concentrations of albumin in human urine
Horton J.K., Davies M., Woodhead J.S., Weeks I. 45
Octadeca-9,11-dienoic acid in cervical intraepithelial neoplasia: A colposcopic study
Fairbank J., Hollingworth A., Griffin J., Ridgway E., Wickens D., Singer A., Dormandy T. 53
Microheterogeneity of the carbohydrate moity of human alpha 1-acid glycoprotein in two benign liver diseases: Alcoholic cirrhosis and acute hepatitis
Biou D., Chanton P., Konan D., Seta N., N'Guyen H., Feger J., Durand G. 59
Progress and problems in immunoassays for serum pituitary gonadotrophins: Evidence from the UK external quality assessment schemes (EQAS) 1980–1988
Seth J., Hanning I., Bacon R.R.A., Hunter W.M. 67
The evaluation of a chemiluminescent assay for free thyroxine by comparison with equilibrium dialysis in clinical samples
Beaman J., Woodhead J.S., Liewendahl K., Mahonen H. 83
On optimising the diagnostic yield of secretin pancreozymin tests
Hunt L.P., Braganza J.M. 91
A simple method for the measurement of total and glycated apolipoprotein B and its relevance to apolipoprotein-B metabolism in diabetes mellitus
Kortlandt W., Benschop C., Erkelens D.W., Thijssen J.H.H. 109
Characterization of trypsin-activatable latent collagenases and their inhibitors in rheumatoid arthritis synovial fluid
Tagawa M., Kobayashi J., Akama T. 119

Vol. 186/2, 1990

On the importance of being (stereo) specific
Fishman W.H. 129
The human alkaline phosphatases: What we know and what we don't know
Harris H. 133
Purification and some properties of the fast migrating alkaline phosphatase in FL-amnion cells (the Kasahara isoenzyme) and its cDNA cloning
Higashino K., Muratani K., Hada T., Imanishi H., Amuro Y., Yamamoto Y., Furuyama J., Hirano K., Hong Y.-M., Shimokura M., Hirano T., Kishimoto T. 151
Comparison of the three PLAP-related genes on human chromosome 2
Martin D., Tucker D., Campbell I., Trowsdale J. 165
Cloning and characterization of a cDNA encoding alkaline phosphatase in mouse embryonal carcinoma
Hahnel A.C., Schultz G.A. 171
Structure of alkaline phosphatases
Kim E.E., Wyckoff H.W. 175
An active site mutant of human placental alkaline phosphatase with deficient enzymatic activity and preserved immunoreactivity
Narisawa S., Sowadski J.M., Millan J.L. 189
Purified tetrameric alkaline phosphatase: The effect of treatments with phosphatidylinositol phospholipase C and sodium dodecyl sulfate
Hawrylak K., Kihn L., Rutkowski D., Stinson R.A. 197
Allelic and ectopic polymorphisms in human placental alkaline phosphatases: Sugar chain heterogeneities
Komoda T., Sato M., Furiya K., Sakagishi Y., Sekine T. 203
Secretion of hepatic and intestinal alkaline phosphatases: Similarities and differences
Alpers D.H., Eliakim R., DeSchryver-Kecskemeti K. 211
Human fetal intestinal alkaline phosphatase: Molecular heterogeneity and immunological detection in amniotic fluids
Verpooten G.F., Hoylaerts M.F., Nouwen E.J., De Broe M.E. 225
Alkaline phosphatase of rat intestinal lymph: Its characterization and the effect of fat administration
Miura S., Yoshioka M., Hamada Y., Morita A., Asakura H., Tsuchiya M. 239
Alkaline phosphatase releasing activity in human tissues
Hamilton B.A., McPhee J.L., Hawrylak K., Stinson R.A. 249
Enzymatic heterogeneity of seminomas
Koshida K., Wahren B.E. 255
Purification and partial characterization of the placental-like alkaline phosphatase in human lung tissue
Hirano K., Koyama I., Stigbrand T. 265
Serum placental alkaline phosphatase (PLAP) in gynecologic malignancies — With special reference to the combination of PLAP and CA54/61 assay
Nozawa S., Udagawa Y., Ohkura H., Negishi Y., Akiya K., Inaba N., Takamizawa H., Kimura E., Terashima Y. 275
Placental-type alkaline phosphatase in peritoneal fluid of women with endometriosis
Kang J.O., Hudak W.A., Crowley W.J., Criswell B.S. 285
Placental-like alkaline phosphatase from ovary and ascites of a patient with a malignant ovarian tumour
Koyama I., Stendahl U., Stigbrand T. 295
Histochemical, ultracytochemical and biochemical study of alkaline phosphatase activity during gastric carcinogenesis
Hirota N., Sakai T., Komoda T. 301
An alkaline phosphatase reacting with both monoclonal antibodies to intestinal and placental isoenzymes
Imanishi H., Hada T., Muratani K., Hirano K., Higashino K. 309

The preparation of monoclonal antibodies which react preferentially with human bone alkaline phosphatase and not liver alkaline phosphatase
Hill C.S., Wolfert R.L. 315

Vol. 186/3, 1990

Two ELISA's for measurement of protein S, and their use in the laboratory diagnosis of protein S deficiency
Deutz-Terlouw P.P., Ballering L., Van Wijngaarden A., Bertina R.M. 321
Transmembrane cation fluxes and fatty acid composition of erythrocytes in psoriatic patients
Corrocher R., Bassi A., Gandini A., Guarini P., Trevisan M.T., Schena D., Olivieri O., Ferrari S. 335
Experimental studies on unconjugated bilirubin binding by human erythrocytes
Hayer M., Piva M.-T., Sieso V., Magnan de Bornier B. 345
Serial measurements of Ca antigen in sera from patients with advanced malignant breast disease
Searle F., Chantler S.M., Goodall A.B., Adam T., Bagshawe K.D., Mitchell H.D.C., Boxer G.M., Southall P.M. 351
Human liver nicotinamide N-methyltransferase: Ion-pairing radiochemical assay, biochemical properties and individual variation
Rini J., Szumlanski C., Guerciolini R., Weinshilboum R.M. 359
Quantitation of hydrogen peroxide formed during UV-visible irradiation of protoporphyrin, coproporphyrin and uroporphyrin
Menon I.A., Becker M.A.C., Persad S.D., Haberman H.F. 375
Quantitative assay of free and total carnitine using tandem mass spectrometry
Kodo N., Millington D.S., Norwood D.L., Roe C.R. 383
Enzyme immunoassay for urinary albumin at low concentration in diabetes mellitus
Niwa T., Katsuzaki T., Tatemichi N., Kato K., Yazawa T., Maeda K. 391
Low biotinidase activities in the sera of patients with impaired liver function: evidence that the liver is the source of serum biotinidase
Grier R.E., Heard G.S., Watkins P., Wolf B. 397

Vol. 187/1, 1990

Characterization of prolidase I and II from erythrocytes of a control, a patient with prolidase deficiency and her mother
Ohhashi T., Ohno T., Arata J., Sugahara K., Kodama H. 1
Sensitive enzyme immunoassay for human Cu/Zn superoxide dismutase
Kurobe N., Suzuki F., Okajima K., Kato K. 11
Immunochemical assays of serum proteins: A European external quality assessment survey and the effects of calibration procedures on interlaboratory agreement
Bullock D.G., Dumont G., Vassault A., Aguzzi F., Chambers R.E., Milford Ward A., Whicher J.T., Bienvenu J. 21
Validation of a method to measure the uptake of [^{3}H]dopamine by human platelets
Dean B., Kulkarni J., Copolov D.L. 37
Influence of angiotensin converting enzyme inhibitor (captopril) on kidney epithelial cells in vitro: Studies on potassium (^{86}Rb) influx and cellular proliferation
Bauer H., Lametschwandtner A., Steiner M., Bauer H.C. 47
Accumulation of mannosyl-β(1→4)-N-acetylglucosamine in fibroblasts and leukocytes of patients with a deficiency of β-mannosidase
Van Pelt J., Hokke C.H., Dorland L., Duran M., Kamerling J.P., Vliegenthart J.F.G. 55

Vol. 187/2, 1990

Detection of neuron specific enolase concentrations in cerebrospinal fluid from patients with neurological disorders by means of a sensitive enzyme immunoassay
Vermuyten K., Lowenthal A., Karcher D. 69

Automated enzymatic assays for the determination of intestinal permeability probes in urine. 1. Lactulose and lactose
Northrop C.A., Lunn P.G., Behrens R.H. 79

A new sensitivity microassay for the measurement of erythrocyte glycogen
Farquharson J., Jamieson E.C., MacPhee G.B., Logan R.W. 89

Development and validation of an automated enzyme assay for paracetamol (acetaminophen)
Morris H.C., Overton P.D., Ramsay J.R., Campbell R.S., Hammond P.M., Atkinson T., Price C.P. 95

Characterization of aminopeptidases in human kidney soluble fraction
Mantle D., Lauffat B., McDermott J., Gibson A. 105

Post-transcriptional modification of serum creatine kinase in infected intensive care patients
Delanghe J.R., De Buyzere M.L., De Scheerder I.K., Van Rostenberghe H.L., Faust U., Rodenbach J.M., Kruse-Jarres J.D., Wieme R.J. 115

Comparison of serum and urine neopterin concentrations in patients with HIV-1 infection
Fuchs D., Jaeger H., Popescu M., Reibnegger G., Werner E.R., Kaboth W., Wachter H. 125

A fluorimetric enzyme assay for the diagnosis of Morquio disease type A (MPS IV A)
Van Diggelen O.P., Zhao H., Kleijer W.J., Janse H.C., Poorthuis B.J.H.M., Van Pelt J., Kamerling J.P., Galjaard H. 131

Elevation of hormone-binding globulins in acute intermittent porphyria
Herrick A.L., McColl K.E.L., Wallace A.M., Moore M.R., Goldberg A. 141

Dye-stained gelatin microcarriers as insoluble chromolytic substrates for the determination of proteolytic activity
Safarik I. 149

Enzymatic erythrocyte creatine determinations as an index for cell age
Buysse A.M., Delanghe J.R., De Buyzere M.L., De Scheerder I.K., De Mol A.M., Noens L. 155

A microalbuminuria assay using bromphenol blue
Jung K., Nickel E., Pergande M. 163

A sensitive and specific assay for granulocyte elastase in inflammatory tissue fluid using L-pyroglutamyl-L-prolyl-L-valine-p-nitroanilide
Tanaka H., Shimazu T., Sugimoto H., Yoshioka T., Sugimoto T. 173

Alterations in manganese and iodide contents in human thyroid tumors; a correlation between the contents of essential trace elements and the states of malignancy
Nishida M., Sakurai H., Tezuka U., Kawada J., Koyama M., Takada J. 181

Spectrophotometric evaluation of results of the ascorbate and the ascorbate-cyanide screening tests for glucose-6-phosphate dehydrogenase deficiency
Rakitzis E.T., Manolakou K., Papandreou P. 189

Vol. 187/3, 1990

Kawasaki disease and oxidative metabolism
Lebranchu Y., Malvy D., Richard M.J., Arnaud J., Favier A., Bardos P. 193

Sialyllacto-N-fucopentaose II (sialylated Le^a) coupled to human serum albumin used as standard in immunoassays of tumour associated antigens CA 19-9 and CA-50 in serum
Masson P., Lundblad A., Wieslander J. 199

Characterization of a monoclonal antibody specific for prostatic secretory protein of 94 amino acids (PSP94) and development of a two-site binding enzyme immunoassay for PSP94
Von Der Kammer H., Krauhs E., Aumuller G., Scheit K.H. 207

Demonstration of a combined deficiency of xanthine oxidase and aldehyde oxidase in xanthinuric patients not forming oxipurinol
Reiter S., Simmonds H.A., Zollner N., Braun S.L., Knedel M. 221

Microheterogeneity of α_1-acid glycoprotein: Variation during the menstrual cycle in healthy women, and profile in women receiving estrogen-progestogen treatment
Succari M., Foglietti M.-J., Percheron F. 235
Comparative evaluation of the diagnosis of acute pancreatitis based on serum and urine enzyme assays
Gwozdz G.P., Steinberg W.M., Werner M., Henry J.P., Pauley C. 243
Anti-skeletal muscle antibodies in the sera from myasthenic patients with thymoma: Identification of anti-myosin, actomysin, actin, and α-actinin antibodies by a solid-phase radioimmunoassay and a Western blotting analysis
Ohta M., Ohta K., Itoh N., Kurobe M., Hayashi K., Nishitani H. 255
Sensitive enzyme immunoassay for human aldolase A
Okajima K., Kurobe N., Shimizu K., Kato K. 265
Variations in serum alpha-L-fucosidase activity during childhood and pregnancy
Vaysse J., Pilardeau P., Gattegno L. 273
A routine method for the measurement of the sodium, potassium, magnesium and calcium content of human lymphocytes
Weissberg N., Brooks B.A., Schwartz G., Eylath U., Abraham A.S. 281
How should urinary cotinine concentrations be adjusted for urinary creatinine concentration?
Thompson S.G., Barlow R.D., Wald N.J., Van Vunakis H. 289
Isoelectric focusing versus quantitative measurements in the detection of intrathecal local synthesis of IgG
Luxton R.W., McLean B.N., Thompson E.J. 297
Non-M CK — A practical measure of creatine kinase isoenzymes in cancer patients
McGing P.G., Teeling M., McCann A., Kyne F., Carney D.N. 309
Measurement of L-carnitine and acylcarnitines in body fluids and tissues in children and in adults
De Sousa C., English N.R., Stacey T.E., Chalmbers R.A. 317
Early amniocentesis and amniotic fluid organic acid levels in the prenatal diagnosis of organic acidemias
Coude M., Chadefaux B., Rabier D., Kamoun P. 329
A carry-over study in three different instruments for the determination of serum glucose: The importance of student's t test
Yucel D., Kantaroglu N., Saydam G.S., Balk M., Top S. 333

Vol. 188/1, 1990

Metal inhibition of ferrochelatase activity in human lymphocytes
Rossi E., Attwood P.V., Garcia-Webb P., Costin K.A. 1
Isolation, identification, and analysis of 4-acetylaminophenol-glucuronide in body fluids of dialyzed renal patients; a molecular mass marker for peritoneal diffusive transport
Schoots A.C., Koomen G.C.M., Struijk D.G., Krediet R.T., Arisz L. 15
The effect of synthetic very low calorie diets on the GH-IGF-1 axis in obese subjects
Gama R., Teale J.D., Marks V. 31
A new hemoglobin variant found during investigations of diabetes mellitus: Hb Pavie (α 135 (H18) Val→Glu)
Wajcman H., Blouquit Y., Riou J., Kister J., Poyart C., Soria J., Galacteros F. 39
A simple serum galactosyltransferase assay method suitable for routine use
Sichel F., Malas J.-P., Gauduchon P., Aubert M., Bar E., Le Talaer J.-Y. 49
A comparison of serum mucoprotein with serum α_1 acid glycoprotein, haptoglobin, and α_1 antitrypsin assays in monitoring inflammatory bowel disease
Lubega J., Davies T.J. 59
Does electrophoresis reliably screen for high serum lipoprotein Lp(a)?
Bruckert E., Truffert J., De Gennes J.L., Lagarde J.P., Federspiel C.B.M.C., Giral P., Dairou F., Bernard C. 71
Urinary paraquat in acute intoxication
Kuo T.-L. 79

Urinary organic acids in elderly Japanese patients with ketosis and encephalopathy who have taken panto-yl-gamma-aminobutyrate, calcium salt (calcium hopantenate, HOPA)
Nakanishi T., Funahashi S., Shimizu A., Hayashi A. 85
Newsletter of the Societa Italiana di Biochimica Clinica
Deledda R. 92

Vol. 188/2, 1990

Urinary excretion of brush-border antigen and plasma proteins in early stages of diabetic nephropathy
Mutti A., Alinovi R., Ghiggeri G.M., Bergamaschi E., Candiano G., Rasi A., Gusmano R., Franchini I., Borghetti A. 93
Early alterations of plasma free amino acids in chronic renal failure
Ceballos I., Chauveau P., Guerin V., Bardet J., Parvy P., Kamoun P., Jungers P. 101
Identification of an intermediate form of β-N-acetylhexosaminidase in chorionic villi
Pieretti M., Beccari T., Cosmi E.V., Stirling J.L., Orlacchio A. 109
The increased plasma Lp(a): B lipoprotein particle concentration in angina pectoris is not associated with hypofibrinolysis
Alessi M.C., Parra H.J., Joly P., Vu-Dac N., Bard J.M., Fruchart J.C., Juhan-Vague I. 119
Usefulness of a prognostic inflammatory and nutritional index in pediatric clinical practice
Pressac M., Vignoli L., Aymard P., Ingenbleek Y. 129
Decreased xanthine oxidase activities and increased urinary oxypurines in heterozygotes for hereditary xanthinuria
Kawachi M., Kono N., Mineo I., Yamada Y., Tarui S. 137
Quantitative determination of IgM-rheumatoid factor by enzyme immunoassay — Standardization using a serum from a rheumatoid arthritis patient
Oka H., Hirohata S., Inoue T., Iwamoto S., Miyamoto T. 147
Increased urinary excretion of β-hydroxyisovaleric acid in ketotic and nonketotic type II diabetes mellitus
Yu W., Kuhara T., Inoue Y., Matsumoto I., Iwasaki R., Morimoto S. 161
Nuclear magnetic resonance studies of lithium transport in erythrocyte suspensions of hypertensives
Ramasamy R., Mota de Freitas D., Bansal V.K., Dorus E., Labotka R.J. 169
Improved specific assay for serum arylesterase using a water-soluble substrate
Kawai H., Sakamoto F., Inoue Y. 177
Elevated spermidine level in pleural fluid of patients with lung cancer
Abe K., Saito T., Kitamura S., Konaka R. 183
Stability of gonadotropins in urine
Khatkhatay I., Desai M., Sankolli G., Joshi U. 185

Vol. 188/3, 1990

Immunochemical determination of lipoprotein Lp(a): Comparison of Laurell electrophoresis and ELISA
Kostner G.M., Gries A., Pometta M., Molinari E., Pichler P., Aicher H. 187
Additional fucosyl residues on membrane glycoproteins but not a secreted glycoprotein from cystic fibrosis fibroblasts
Wang Y.-M., Hare T.R., Won B., Stowell C.P., Scanlin T.F., Glick M.C., Hard K., Van Kuik J.A., Vliegenthart J.F.G. 193
The phospholipid composition of erythrocyte ghosts and plasma lipoproteins in diabetes type 1 in children
Watala C., Jozwiak Z. 211
O-Phosphohydroxylysinuria: A new inborn error of metabolism?
Dorland L., Duran M., De Bree P.K., Smith G.R., Horvath A., Tibosch A.S., Wadman S.K. 221

Diurnal variation of erythrocyte sodium-lithium countertransport rate and intracellular cation concentrations
Siebers R.W.L., Maling T.J.B. 227
Carboxypeptidase activity in human urine from healthy subjects and renal disease patients
Hamai K., Ikeda R., Sumi H., Mihara H. 233
Inosine triphosphate pyrophosphohydrolase deficiency in a kindred with adenosine deaminase deficiency
Duley J.A., Simmonds H.A., Hopkinson D.A., Levinsky R.J. 243
Evaluation of a radioimmunoassay for somatomedin-C/insulin-like growth factor I (Sm-C/IGF-I) in human plasma
Giannella-Neto D., Cavaleiro A.M., Sadoyama R., Wajchenberg B.L., Spencer E.M. 253
The effect of incubation with Staphylococcus aureus Cowan I on subsequent determinations of IgG, IgM, and IgM rheumatoid factor by ELISA
Nesher G., Moore T.L. 261
Spectrophotometric estimation of erythrocyte ATP and deoxyadenosine triphosphate using hexokinase and glucokinase
Nakashima K., Nakashima H., Shimoyama M. 267
Succinic acidemia is not a new syndrome of organic acidemia
Elpeleg O.N., Hurvitz H. 271
Monoclonal gammopathies. Their identification and biological significance (I)
Radl J. 273

Vol. 189/1, 1990

Influence of human transferrin and group-specific protein on endotoxicity in vitro
Berger D., Winter M., Beger H.G. 1
Diagnosis of alcaptonuria: Rapid analysis of homogentisic acid by HPLC
Bory C., Boulieu R., Chantin C., Mathieu M. 7
65 Zinc absorption in patients with insulin-dependent diabetes mellitus assessed by whole-body counting technique
Kiilerich S., Hvid-Jacobsen K., Vaag A., Sorensen S.S. 13
Effect of diet on urinary excretion of unconjugated catecholamines and some of their metabolites in healthy control subjects and depressed patients
Moleman P. 19
The effect of n-3 polyunsaturated fatty acids on lipids, platelet function, coagulation, fibrinolysis and monocyte chemotaxis in patients with hypertension
Schmidt E.B., Nielsen L.K., Pederson J.O., Kornerup H.J., Dyerberg J. 25
Direct reading glucose electrodes detect the molality of glucose in plasma and whole blood
Fogh-Anderson N., Wimberley P.D., Thode J., Siggaard-Andersen O. 33
Use of a new simplified assay for phospholipase A2 to measure bacterial enzyme levels
Lumb M.R., Roseblade C.K., Helmig R., Uldbjerg N., Sullivan M.H.F., Elder M.G. 39
The acute phase protein response in mice does not show tolerance to recurrent sterile inflammation
Todd N.J., Whicher J.T., Westacott C., Gilbert A. 47
International study on the comparability of Apo A-1 and Apo B methods
Grafnetter D., Molinari E., Lonsky L. 55
High perfomance liquid chromatography methods for studying protein binding of aluminium in human serum in the absence and in the presence of desferrioxamine
Garcia Alonso J.I., Lopez Garcia A., Parajon J., Blanco Gonzalez E., Sanz Medel A., Cannata J.B. 69
S-Adenosylhomocysteine hydrolase and adenosine deaminase activities in human red cell ageing
Bozzi A., Furciniti-La Chiusa B., Strom R., Crifo C. 81
Changes in phospholipid composition of tracheal aspirates from newborns with hyaline membrane disease or transient tachypnoea
Bourbon J.R., Francoual J., Magny J.F., Lindenbaum A., Leluc R., Dehan M. 87

Vol. 189/2, 1990

Development of a double antibody radioimmunoassay for quantitation of 1α,25-dihydroxyvitamin D
Armbruster F.-P., Reichel H., Vogel G., Georgousis H., Schmidt-Gayk H. 97

Distribution of HDL_2 and HDL_3 in a random population of healthy French males and females — evaluation by a two-step precipitation procedure
Atger V., Wirbel E., Roche D., Apfelbaum M., Burstein M., Girard-Globa A. 111

Detection and estimation of the barley prolamin content of beer and malt to assess their suitability for patients with coeliac disease
Ellis H.J., Freedman A.R., Ciclitira P.J. 123

The levels of adenine nucleotides and pyridine coenzymes in red blood cells from the newborn, determined simultaneously by HPLC
Formato M., Masala B., De Luca G. 131

Human liver L-alanine–glyoxylate aminotransferase: Characteristics and activity in controls and hyperoxaluria type I patients using a simple spectrophotometric method
Wanders R.J.A., Ruiter J., Van Roermund C.W.T., Schutgens R.B.H., Ofman R., Jurriaans S., Tager J.M. 139

Hyperphosphatasemia related to three intestinal alkaline phosphatase isoforms: Biochemical study
Bentouati L., Samadi Baboli M., Hachem H., Hamza M., Canal P., Soula G. 145

Detection of apoliproprotein E polymorphisms using PCR/ASO probes and Southern transfer: Application for routine use
Houlston R.S., Wenham P.R., Humphries S.E. 153

Is basic red cell ferritin a more specific indicator than serum ferritin in the assessment of iron stores in the elderly?
Galan P., Sangare N., Preziosi P., Roudier M., Hercberg S. 159

Energy dependence of cytosolic enzyme efflux from rat skeletal muscle
West-Jordan J.A., Martin P.A., Abraham R.J., Edwards R.H.T., Jackson M.J. 163

Determination of α_1-antitrypsin in fecal extracts by enzyme immunoassay
Brouwer J., Smekens F. 173

Cutaneous interstitial fluid protein concentrations in the inflammatory syndrome: Pharmacological consequences
Clerc C., Pibouin M., Ruelland A., Legras B., Chevrant-Breton J., Cloarec L. 181

Combined diagnostic value of biochemical markers in acute pancreatitis
Viel J.F., Foucault P., Bureau F., Albert A., Drosdowsky M.A. 191

Rapid spectrophotometric assay of serum iron
Johnson D.J., Williams H.L. 199

Protease activities in gastric cancer tissues
Chung S.M., Kawai K. 205

Glomerular and urinary glycosaminoglycans in diabetic rats
Reddi A.S. 211

Human platelet thermostable phenol sulfotransferase: Assay of frozen samples and correlation between frozen and fresh activities
Anderson R.J., Liebentritt D.K. 221

Inhibition of lactate dehydrogenase by an EDTA · Ca^{2+} complex
Shina R., Condrea E. 231

Plasma polymorphonuclear leukocyte elastase-α_1-proteinase inhibitor complex in patients with gastric ulcer
Okajima K., Inoue M., Koga S., Okabe H., Takatsuki K. 237

Kinetic enzymatic assay for ethylene glycol
Hansson P. 243

Studies on the clinical significance of serum α_2HS-glycoprotein measurement in metabolic bone diseases
Scarnecchia L., Minisola S., Pacitti M.T., Carnevale V., Bigi F., Romagnoli E., Mazzuoli G.F. 245

Proper handling of blood samples is crucial for the accurate ion chromatographic determination of oxalate
Petraculo M., Marangella M. 249
Inhibition by a free-radical scavenger of ascorbate-induced hemoglobin denaturation in glucose-6-phosphate dehydrogenase deficient erythrocytes
Papandreou P.T., Rakitzis E.T. 253

Vol. 189/3, 1990

A time-resolved fluoroimmunoassay for the determination of prostaglandin F(2α)
Luke F.J., Schlegel W. 257
Immunochemical detection of human blood in feces
Uchida K., Matsuse R., Miyachi N., Okuda S., Tomita S., Miyoshi H., Hirata I., Tsumoto S., Ohshiba S. 267
Development of a radioimmunoassay for neurone specific enolase (NSE) and its application in the study of patients receiving intra hepatic arterial streptozotocin and floxuridine
Cunningham R.T., Johnston C.F., Irvine G.B., McIlrath E.M., McNeill A., Buchanan K.D. 275
Apolipoprotein B-48 analysis by high-performance liquid chromatography in VLDL: A sensitive and rapid method
Hidaka H., Kojima H., Nakajima Y., Aoki T., Nakamura T., Kawabata T., Nakano T., Harano Y., Shigeta Y. 287
Enzymatic diagnosis of 3-hydroxy-3-methylglutaryl-CoA lyase deficiency with high-performance liquid chromatography
Kikuchi M., Narisawa K., Tada K., Sweetman L. 297
Decreased proapoliprotein A-I processing in liver disease: Evidence for hepatic participation in proapolipoprotein A-I conversion
Isobe H., Sakai H., Satoh M., Sakamoto S., Koga S., Nawata H. 303
Lipoamidase activity in human serum is due to biotinidase
Garganta C.L., Wolf B. 313
Rapid diagnosis of 3-hydroxy-3-methylglutaryl-coenzyme A lyase deficiency via enzyme activity measurements in leukocytes or platelets using a simple spectrophotometric method
Wanders R.J.A., Zoeters P.H.M., Schutgens R.B.H., De Klerk J.B.C., Duran M., Wadman S.K., Van Sprang F.J., Hemmes A.M., Voorbrood B.S. 327
Evidence of acid hydrolysis as responsible for 2-(2-furoyl)-4(5)-(2-furanyl)-1H-imidazole (FFI) production
Gerhardinger C., Lapolla A., Crepaldi G., Fedele D., Ghezzo E., Seraglia R., Traldi P. 335

Vol. 190/1–2, 1990

The biology of transferrin
De Jong G., Van Dijk J.P., Van Eijk H.G. 1
Incidence and properties of aspartate aminotransferase-immunoglobulin complexes in patients with a high serum aspartate to alanine aminotransferase ratio
Moriyama T., Nobuoka M., Makino M. 47
A compensatory mechanism improving red cell membrane fluidity in hemodialysed patients
Peuchant E., Motta C., Salles C., Clerc M. 57
Liquid chromatographic studies of vitamin B6 metabolism in man
Edwards P., Liu P.K.S., Rose G.A. 67
A method for the rapid detection of urinary glycopeptides in α-N-acetylgalactosaminidase deficiency and other lysosomal storage diseases
Schindler D., Kanzaki T., Desnick R.J. 81
Heterogeneity of urinary albumin from diabetic patients
Hayashi Y., Morikawa A., Makino M. 93

Lack of correlation between human serum angiotensin converting enzyme activity and the renin-angiotensin-aldosterone system
Pedraza-Chaverri J., Herrero B., Ibarra-Rubio M.E., Cruz C., Tapia E., Pena J.C. 105
Seed oils containing fatty acids with conjugated double bonds as substrates for lipase assay
Anzai T., Kotake I., Komine S. 107
Serum immunoreactive prolyl 4-hydroxylase as a specific marker of hepatocellular necrosis
Nagai Y., Toda G. 109
Urinary adenosine deaminase binding protein, a marker of contrast media induced acute renal damage
Parvez Z., Patel N.B., Nelson J.E. 111
Glucose interferes with inulin determination by the resorcinol method: Overestimation of glomerular filtration rate in diabetic patients
Erman A., Van Dijk D.J., Pansky R., Wittenberg C., Boner G., Rosenfeld J.B. 115
Plasma fibronectin in patients affected by psoriasis: Relation with the extension of skin involvement and complications
Cecchi R., Brunetti L., Giomi A., Gironi A., De Giorgio L.A., Seghieri G. 119
Improved HPLC method for the determination of cephalosporins in human plasma and a new solid-phase extraction procedure for cefazolin and ceftizoxime
Moore C.M., Sato K., Hattori H., Katsumata Y. 121
Scientific Division. Committee on enzymes. IFCC methods for the measurement of catalytic concentration of enzymes. Part 7. IFCC method for creatine kinase (ATP: creatine N-phosphotransferase, EC 2.7.3.2). IFCC recommendation
Horder M., Elser R.C., Gerhardt W., Mathieu M., Sampson E.J. S4
Scientific Division. Committee on pH, blood gases and electrolytes. Guidelines for transcutaneous $p(O_2)$ and $p(CO_2)$ measurement
Wimberley P.D., Burnett R.W., Covington A.K., Maas A.H.J., Mueller-Plathe O., Siggaard-Andersen O., Weisberg H.F., Zijlstra W.G. S41
Education division expert panel on quantities and units. A protocol for the conversion of clinical laboratory data
Lehmann H.P., Worth H.G.J., Zinder O. S51

Vol. 190/3, 1990

Redox electrode for monitoring dehydrogenase-catalyzed reactions
Chen S.S.-H. 129
Enzyme immunoassay for flecainide
Freche J.P., Coudert G., Wellman M., Siest G. 139
Myoadenylate deaminase deficiency studies on normal and deaminase-deficient skeletal muscle
Kaletha K., Nowak G. 147
Elevation of blood lactate and pyruvate levels in acute intermittent porphyria — A reflection of haem deficiency?
Herrick A.L., Fisher B.M., Moore M.R., Cathcart S., McColl K.E.L., Goldberg A. 157
Comparison of the effects of purified human α_1-antichymotrypsin and α_1-proteinase inhibitor on NK cytotoxicity: Only α_1-proteinase inhibitor inhibits natural killing
Laine A., Leroy A., Hachulla E., Davril M., Dessaint J.-P. 163
Enzyme immunosorbant assay of oestradiol in unextracted plasma using penicillinase as label
Pandey P.K., Shrivastav T.G., Kumari G.L., Rao P.N., Grover P.K., Murthy H.G.K. 175
Immunoreactive vasopressin in end stage renal failure
Argent N.B., Baylis P.H., Wilkinson R. 185
Theoretical isoelectric points and electric charges of mutated human hemoglobin subunits
Meireles Ribeiro J., Ruiz A., Sillero M.A.G., Sillero A. 189
A sensitive radioimmunoassay using a monoclonal antibody that is equipotent for ercalcitriol and calcitriol (1,25-dihydroxy vitamin D_2 and D_3)
Mawer E.B., Berry J.L., Cundall J.P., Still P.E., White A. 199

The effect of oral contraceptive use and pregnancy on the daily rhythm of cortisol and cortisone
Meulenberg MN. P.M., Hofman J.A. 211
Does Tamm-Horsfall mucoprotein inhibit or promote calcium oxalate crystallization in human urine?
Grover P.K., Ryall R.L., Marshall V.R. 223
Two-site assay of intact parathyroid hormone in primary hyperparathyroidism: Studies in basal conditions, following adenoma removal and during calcium and EDTA infusion
Mazzuoli G., Minisola S., Scarnecchia L., Pacitti M.T., Carnevale V., Romagnoli E., Bigi F., Bianchi G. 239
Urinary proteins and red blood cell membrane negative charges in diabetes mellitus
Bernard A., Amor A.O., Goemare-Vanneste J., Antoine J.-L., Lauwerys R., Colin I., Vandeleene B., Lambert A. 249
Immunological study on hypophosphatasia
Goseki M., Oida S., Takagi Y., Okuyama T., Watanabe J., Sasaki S. 263
Dopamine-mediated effects induced by metoclopramide simultaneously on adrenocorticotropic hormone and prolactin in patients with pituitary and/or hypothalamic disorders
Nishida S., Matsuki M., Horino M., Kawai Y., Tsushima K., Yoneda M., Oyama H., Ishii R. 269
Different molecular forms of acid-stable trypsin inhibitor in normal and uremic plasma
Yoshida E., Sumi H., Mihara H. 277
The effect of non-insulin-dependent diabetes on serum concentrations of tumor-associated carbohydrate antigens of Ca19-9, CA-50, and sialyl SSEA-1 in association with the Lewis blood phenotype
Shimojo N., Naka K., Nakajima C., Ishizaki T., Okuda K., Murai J., Yamamoto M. 283
Serum alkaline phosphatase in sickle cell anaemia
Ogunleye I.O., Adekile A.D., Fakunle J.B. 291
Presence of thiosulfate in human plasma
Lundquist P. 293

Vol. 191/1–2, 1990

Concentrations of N-terminal ProANP in human plasma: evidence for ProANP (1-98) as the circulating form
Buckley M.G., Sagnella G.A., Markandu N.D., Singer D.R.J., MacGregor G.A. 1
Detection of aspartylglycosaminuria using urine specimens recovered from absorbent filter paper
Kaartinen V., Mononen I. 15
Free testosterone levels in plasma and saliva as determined by a direct solid-phase radioimmunoassay: a critical evaluation
Rey F., Chiodoni G., Braillard K., Berthod C., Lemarchand-Beraud T. 21
Determination of aluminium in bone in haemodialyzed patients, using inductively coupled argon plasma emission spectrometry
Leflon P., Plaquet R., Moriniere P., Fournier A. 31
Stable isotope dilution analysis of cystine in granulocyte suspensions as cysteine: a powerful method for the diagnosis, the follow up, and treatment of patients with cystinosis
Schierbeek H., Bijsterveld K., Chapman T.E., Van Luijk W.H.J., Reijngoud D.J., Berger R. 39
Studies on lipoamidase: characterization of the enzyme in human serum and breast milk
Backman-Gullers B., Hannestad U., Nilsson L., Sorbo B. 49
Presence of reverse transcriptase in human leukemias and lymphomas
Moyret C.F., Bernard D.J., Maurizis J.-C., Chassagne J., Plagne R., Chollet P. 61
Plasma oxalate and creatinine and oxalate/creatinine clearance ratios in normal subjects and in primary hyperoxaluria. Evidence for renal hyperoxaluria
Kasidas G.P., Nemat S., Rose G.A. 67
Longitudinal changes in serum fructosamine do not parallel those in glycated haemoglobin in young adults with insulin-dependent diabetes
Fisken R.A., Chan A.W., Hanlon A., MacFarlane I.A. 79

Lectin binding to serum 5′-nucleotidase and alkaline phosphatase in liver disease
Massoubre C., Verdier P., Meflah K., Masliah C., Perrin D., Orsonneau J.-L., Harb J. 87
Why serum F protein is a sensitive and specific marker of liver damage in man
Grewal G.K., Oliveira D.B.G. 93
Enzymatic determination of serum total and esterified cholesterol with the same reagents
Matsushita M., Irino T., Maehata E. 97
Ribonuclease activities in liver tissue of patients with chronic liver diseases
Tsuji H., Murai K., Akagi K., Fujishima M. 101
Labile HbA(1c) formation in fractionated erythrocytes in aerobic and anaerobic conditions
Yoshida Y., Nakashima K. 105
Storage and thawing influence plasma levels of immunoreactive atrial natriuretic peptide
Tan A.C.I.T.L., Kloppenborg P.W.C., Benraad Th. J. 111
Bayes' theorem should be considered when making decision limits
Ohman S. 115
On the role of thyroglobulin as an antigen of thyroid hormone autoantibodies
Sakata S. 119

Vol. 191/3, 1990

Clinical biochemistry of preeclampsia and related liver diseases of pregnancy: A review
Freund G., Arvan D.A. 123
Increased immunoreactivity of apolipoprotein B epitopes during prolonged storage of low density lipoproteins
La Belle M., Mok K.S., Karu A.E., Krauss R.M. 153
Histamine release with a solid-phase-coupled allergen
Mita H., Tadokoro K., Yasueda H., Shida T. 161
A simple enzyme-linked immunoassay of human carbonic anhydrase I for the study of developing erythroid cells
Kondo T., Murakami K., Isobe H., Taniguchi N., Kawakami Y. 169
Angiotensin I-converting enzyme activity in puromycin aminonucleoside-nephrotic syndrome
Arevalo A.E., Ibarra-Rubio M., Cruz C., Pena J.C., Pedraza-Chaverri J. 175
Comparative biochemical and immunological studies on gamma-glutamyltransferases from human kidney and renal cell carcinoma applying monoclonal antibodies
Fischer P., Scherberich J.E., Schoeppe W. 185
Amino acid profiles by HPLC after filter paper sampling: 'Appropriate technology' for monitoring of nutritional status
Lundsjo A., Hagelberg S., Palmer K., Lindblad B.S. 201
One-step sandwich enzyme immunoassay for human laminin using monoclonal antibodies
Iwata K. 211
Gamma-glutamyltransferase from human hepatoma cell lines: Purification and cell culture of HepG2 on microcarriers
Visvikis A., Goergen J.-L., Oster T., Bagrel D., Wellman M., Marc A., Engasser J.-M., Siest G. 221
A reversed-phase high-performance liquid chromatographic method for the determination of pyruvate dehydrogenase complex activity in human platelets and cultured skin fibroblasts
Maehara M., Kajita M., Kitoh M., Watanabe K. 233
Interference of abnormal hemoglobins on the measurement of hemoglobin A(1c) by ion-exchange chromatography
Engbaek F., Sorensen G.H., MacIntyre B., Clausen I., Lund H.T., Jastrup B. 239

Vol. 192/1, 1990

Ferritin and transferrin levels in human breast cyst fluids: Relationship with intracystic electrolyte concentrations
Troccoli R., Stella F., Biagioni S., Battistelli S., Cerroni L., Marcheggiani F., Artico M., Terzano C., Mannello F. 1

Cytochrome oxidase deficiency affecting the structure of the myofibre and the shape of mitochondrial cristae membrane
Sumegi B., Melegh B., Adamovich K., Trombitas K. 9
Interaction between endogenous circulating sulfated-glycosaminoglycans and plasma proteins
Pasquali F., Oldani C., Ruggiero M., Magnelli L., Chiarugi V., Vannucchi S. 19
Gunn rats: A reproducible experimental model to compare the different methods of measurements of bilirubin serum concentration and to evaluate the risk of bilirubin encephalopathy
Labrune P., Myara A., Trivin F., Odievre M. 29
Radioimmunoassay for sequence 38–54 of human progastrin: Increased diagnostic specificity of gastrin-cell diseases
Van Solinge W.W., Rehfeld J.F. 35
Radioimmunoassay for plasma neuropeptide-Y in physiological and physiopathological states and response to sympathetic activation
Miller M.A., Sagnella G.A., Markandu N.D., MacGregor G.A. 47
Evaluation of a simple method for the measurement of cytidine deaminase in serum and comparison with a reference method
Thompson P.W., James I.T., Rowell L., Jones D.D. 55
Urinary polyamine and metabolite excretion by children with Zellweger's syndrome
Govaerts L.C.P., Van den Berg G.A., Theeuwes A., Muskiet F.A.J., Monnens L.A.H. 61
The effect of excess daily methionine intake on plasma homocysteine after a methionine loading test in humans
Andersson A., Brattstrom L., Israelsson B., Isaksson A., Hultberg B. 69
One-step solvolysis of 3-, 7- and 12-sulfated free and conjugated bile acids
Princen H.M.G., Meijer P., Kuipers F. 77
A simple procedure to exclude false-positive galactose screening test results caused by antibiotics
Clemens P.C., Plettner C., Vellguth B. 85

Vol. 192/2, 1990

The relationship between erythrocyte transketolase activity and the 'TPP effect' in Wernicke's encephalopathy and other thiamine deficiency states
Nixon P.F., Price J., Norman-Hicks M., Williams G.M., Kerr R.A. 89
Measurement of β_2-microglobulin, retinol-binding protein, α_1-microglobulin and urine protein 1 in healthy children using enzyme-linked immunosorbent assay
Tomlinson P.A., Dalton R.N., Turner C., Chantler C. 99
Enzyme immunoassay of urinary β-hexosaminidase isoenzymes in patients with renal transplants
Hultberg B., Isaksson A., Sterner G., Weibull H. 107
A new, rapid, one-step method for the isolation of platelets from human blood
Ford T.C., Graham J., Rickwood D. 115
Erythrocyte nucleotide stability and plasma hypoxanthine concentrations: Improved ATP stability with short-term storage at room temperature
Simmonds H.A., Micheli V., Davies P.M., McBride M.B. 121
Stable isotope dilution analysis of very long chain fatty acids in plasma, urine and amniotic fluid by electron capture negative ion mass fragmentography
Stellaard F., Ten Brink H.J., Kok R.M., Van Den Heuvel L., Jakobs C. 133
Is dehydroepiandrosterone sulfate a natriuretic digitalis-like factor in human serum?
Goto A., Yamada K., Ishii M., Yoshioka M., Sugimoto T. 145
The antioxidant activity of human semen
Kocak T., Unal T., Toker N.K. 153

Vol. 192/3, 1990

Purines, lactate and phosphate release from child and adult heart during cardioplegic arrest
Smolenski R.T., Swierczynski J., Narkiewicz M., Zydowo M.M. 155

A new approach to the newborn screening for hyperphenylalaninemias: Use of L-phenylalanine dehydrogenase and microtiter plates
Wendel U., Koppelkamm M., Hummel W., Sander J., Langenbeck U. 165
Sensitive enzyme immunoassay for human Mn superoxide dismutase
Kurobe N., Inagaki T., Kato K. 171
Failure to conserve lactose and glucose polymers during frozen storage of fecal specimens: Methods for preservation
Murray R.D., Ailabouni A.H., Powers P.A., Juhling McClung H., Ulysses B., Li K., Heitlinger L.A., Sloan H.R. 181
One-step sandwich enzyme immunoassay for soluble human thrombomodulin using monoclonal antibodies
Kodama S., Uchijima E., Nagai M., Mikawatani K., Hayashi T., Suzuki K. 191
Excretion of epidermal growth factor (EGF) in diabetes
Lev-Ran A., Hwang D.L., Miller J.D., Josefsberg Z. 201
HPLC separation of alkaline phosphatase isoenzymes. Application to the amniotic fluid
Ragougy-Sengler C., Rolland H., Pileire B. 207

Vol. 193/1–2, 1990

Evidence for locally synthesized and clonally restricted immunoglobulin in the synovial fluid from rheumatoid arthritis patients
Carpenter A.B., Huczko E., Eisenbeis Jr. C.H., Kelly R.H. 1
Serum alpha-1-proteinase inhibitor with abnormal properties in ovarian cancer
Thompson S., Wong E., Cantwell B.M.J., Turner G.A. 13
Collagen of the normal and the varicose human saphenous vein: A biochemical study
Maurel E., Azema C., Deloly J., Bouissou H. 27
Serum baseline hyaluronate and disease activity in rheumatoid arthritis
Konttinen Y.T., Saari H., Honkanen V.E.A., Szocsik K., Mussalo-Rauhamaa H., Tulensalo R., Friman C. 39
Biliary excretion of iron in healthy man and in patients with alcoholic cirrhosis of the liver
Raedsch R., Stiehl A., Walker S., Theilmann L., Kommerell B., Waldherr R., Senn M. 49
Studies of circulating parathyroid hormone following parathyroidectomy in renal osteodystrophy
Straffen A.M., Carmichael D.J.S., Fairney A., Hulme B., Snell M.E. 55
The immunological evidence for a phenylalanine hydroxylase like immunoreactive protein in different human cells and tissues
Petruschka L., Rebrin I., Grimm U., Herrmann F.H. 65
Altered protein kinase C and protein kinase A activities in erythrocyte membrane, platelets and lymphocytes of Duchenne muscular dystrophy (DMD) patients
Jagadeesh G., Lavanya M., Anandaraj M.P.J.S., Anjaneyulu A. 79
Evaluation of the particle size distribution of plasma lipoprotein in a patient with Tangier disease by high performance liquid chromatography
Takizawa A., Komoda T., Hokari S., Sakagishi Y., Tanaka A., Hara I. 85
Decreased muscle enoyl-CoA hydratase activities in patients with exercise intolerance
Carroll J.E., McGuire B.S., Hartlage P.L., Howard J.C. 89
Differential diagnosis of an hyperamylasemia in an acute abdominal syndrome
Foucault P., Drosdowsky M.A. 91
Hyperphosphatasemia related to three intestinal alkaline phosphatase isoforms: Biochemical study
Bentouati L., Samadi Baboli M., Hachem H., Hamza M., Canal P., Soula G. 93

Vol. 193/3, 1990

Polymorphonuclear elastase in neonatal sepsis
Tsaka T., Herkner K.R. 103

Reduction of urinary bile alcohol excretion and serum cholestanol in patients with cerebrotendinous xanthomatosis after oral administration of deoxycholic acid
Wolthers B.G., Van der Molen J.C., Walrecht H., Hesselmans L.F.G.M. 113
Use of pralidoxime-Pd(II) complex for the spectrophotometric determination of the cholinesterase reactivator (PAM-2Cl) in the urine
Karljikovic-Rajic K.D., Stankovic B.S. 119
Mass concentration measurements of creatine kinase BB isoenzyme as an index of brain tissue damage
Delanghe J.R., De Winter H.A., De Buyzere M.L., Camaert J.J., Martens F.E., De Praeter C. 125
Proteolysis of human apolipoprotein B: Effect on quantitative immunoturbidimetry
De Bruin T.W.A., Vos M.C., Kortlandt W., Bouma B.N., Erkelens D.W. 137
Effect of fish-oil and vitamin E supplementation on lipid peroxidation and whole-blood aggregation in man
Brown J.E., Wahle K.W.J. 147
Radial immunodiffusion and immunoelectrophoresis compared for identifying autoantibodies to lactate dehydrogenase in human serum
Harff G.A., Backer E.T. 157
Lipoprotein particles in homozygous familial hypercholesterolemic patients treated with portacaval shunt and LDL apheresis
Gairin D., Monard F., Cachera C., Bard J.-M., Amouyel P., Duriez P., Tacquet A., Fruchart J.-C. 165
Effect of ethanol on metabolism of the hypouricemic agents allopurinol and benzbromarone
Kaneko K., Fujimori S., Ishizuka I., Akaoka I. 181
Redox electrode for monitoring oxidase-catalyzed reactions
Chen P.T., Chen S.S.-H. 187
Quantitation of tyrosine-0-sulfate in human urine by ion-pair reverse-phase high-performance liquid chromatography
Suiko M., Prasantha Fernando P.H., Arino Y., Terada M., Nakatsu S., Liu M.-C. 193
Spectrophotometric determination of oxalate in whole blood
Hatch M. 199

Vol. 194/1, 1990

The widening horizons of bioanalytical mass spectrometry
Hill R.E. 1
Analytical methodology for immunoassays and DNA hybridization assays — Current status and selected systems — Critical review
Diamandis E.P. 19
Ultrasensitive enzyme immunoassay
Ishikawa E., Hashida S., Kohno T., Hirota K. 51
Chemiluminescent DNA probes: A comparison of the acridinium ester and dioxetane detection systems and their use in clinical diagnostic assays
Nelson N.C., Kacian D.L. 73
Fluorescence spectroscopy and its application to a new generation of high sensitivity, multimicrospot, multianalyte, immunoassay
Ekins R., Chu F., Biggart E. 91

Vol. 194/2–3, 1990

Simple method for the measurement of antioxidants
Novak Z., Varga S.I., Pataki L., Matkovics B. 115
The effect of doxorubicin on the glutathione content and viability of cultured human lung cancer cell lines A549 and GLC_4 210
Al-Kabban M., Stewart M.J., Watson I.D., Reglinski J. 121

Serum concentrations of carotenoids, retinol, and α-tocopherol in healthy persons determined by high-performance liquid chromatography
Ito Y., Ochiai J., Sasaki R., Suzuki S., Kusuhara Y., Morimitsu Y., Otani M., Aoki K. 131
A delayed-addition enzyme immunoassay for the relative cholesteryl ester transfer protein mass in patients with deficient plasma cholesteryl ester transfer activity
Yamashita S., Sprecher D.L., Matsuzawa Y., Sakai N., Tarui S., Harmony J.A.K., Wetterau J.R. 145
The facile detection of the nt 1226 mutation of glucocerebrosidase by 'mismatched' PCR
Beutler E., Gelbart T., West C. 161
Inhibition of protein kinase C activation by low concentrations of aluminium
Cochran M., Elliott D.C., Brennan P., Chawtur V. 167
gamma-Glutamylglutamine identified in plasma and cerebrospinal fluid from hyperammonaemic patients
Hammond J.W., Potter M., Truscott R., Wilcken B. 173
Immunological quantification of human hexokinase type I
Magnani M., Serafini G., Chiarantini L., Corsi D., Stocchi V. 185
Further characterization of a heat-stable alkaline phosphatase with low sensitivity to L-phenylalanine
Onica D., Rosendahl K., Waldenlind L. 193
Creatine metabolism during metabolic perturbations in patients with organic acidurias
Davies S.E.C., Iles R.A., Stacey T.E., Chalmers R.A. 203
An HPLC-ECD procedure for measuring total phenolsulfotransferase (PST) activity in human liver, platelets and blood
Khoo B.Y., Sit K.H., Wong K.P. 219
The contribution of sex, electrophoretic phenotype, pregnancy and race to the variability of delta-aminolevulinate dehydrase (ALADH) levels in human erythrocytes. A study in Black mixed Brazilians
De Freitas Sousa M.G., Passos Bueno M.R., Olympio da Silva M.C.B., De Borja Goncalves Duarte A.F., Azevedo E.S. 229

Vol. 195/1,2, 1990

Activities of mitochondrial aspartate aminotransferase and creatine kinase isoenzyme MB in serum following coronary bypass surgery
Huseby N.-E., Grostad M., Almdahl S.M., Vaage J. 1
Short- and long-term effects of LDL-apheresis on lipoprotein (a) serum levels
Ritter M.M., Suhler K., Richter W., Schwandt P. 9
A new sandwich-ELISA method for the determination of keratan sulphate peptides in biological fluids employing a monoclonal antibody and labelled avidin biotin technique
Kongtawelert P., Ghosh P. 17
Interactions of pancreatic secretory trypsin inhibitor in small intestinal juice: Its hydrolysis and protection by intraluminal factors
Freeman T.C., Davies R., Calam J. 27
The action of myocrisin on membrane bound sulphhydryl groups
McGrae F., Reglinksi J., Smith W.E., Brzeski M., Sturrock R.D. 41
Atomic absorption and UV-VIS absorption spectrophotometric determination of oxalate in urine by ligand exchange extraction
Leyva J.A.M., Artiga M.P.H., Mendez M.M.A., Perez J.J.Q. 47
Binding of corticotropin-releasing hormone (CRH) in maternal and fetal plasma and in amniotic fluid
Salminen-Lappalainen K., Laatikainen T. 57
Characterisation of the main molecular forms of human fecal immunoglobulin A
Kapel N., Meillet D., Iscaki S., Bouvet J.P., Gobert J.G., Raichvarg D. 67
The effect of polycythemia and hypoxia on urinary N-acetyl-β-D-glucosaminidase activity in newborns
Csathy L., Pocsi I., Kiss L., Balla G., Modi N., Price R.G. 77

Immunoelectrophorectic pattern of α_2-macroglobulin and α_2-macroglobulin-protease complexes
Donnelly P.K., Trotter P., White M., Shenton B.K. 87
Letter to the Editor
Plasma lipids and apolipoproteins in end-stage liver disease
Lontie J.-F., Dubois D.Y., Malmendier C.L., Mathe D., Adam R., Gigou M., Bismuth H. 93

Vol. 195/3, 1991

Superiority of thyrotropin to thyroxine as a tool in the screening for congenital hypothyroidism by the filter paper spot technique
Bourdoux P.P., Van Thi H.V., Courtois P.A., Ermans A.M. 97
Total and ultrafiltrable plasma magnesium in hyper- and hypoparathyroidism, and in calcium-related metabolic disorders
Claeyssens S., Lavoinne A., Daragon A., Josse S., Godin M., Matray F., Kuhn J.M. 107
A rapid extraction method for serum calcitonin
Bucht E., Granberg B., Sjostedt U., Torring O. 115
Exopeptidases in human platelets: An indication for proteolytic modulation of biologically active peptides
Scharpe S.L., Vanhoof G.C., De Meester I.A., Hendriks D.F., Van Sande M.E., Muylle L.M., Yaron A. 125
Stability of β_2-microglobulin and retinol binding protein at different values of pH and temperature in normal and pathological urine
Blumsohn A., Morris B.W., Griffiths H., Ramsey C.F. 133
Effects of smoking on HDL subfractions in myocardial infarction patients: Effects on lecithin-cholesterol acyltransferase and hepatic lipase
Moriguchi E.H., Fusegawa Y., Tamachi H., Goto Y. 139

Vol. 196/1, 1991

A rapid method for determination of human C-peptide in plasma
Saelsen L., Tronier B., Madsbad S., Juel Christensen N. 1
Effects of selenium supplementation on blood and urine selenium levels and liver function in patients with primary biliary cirrhosis
Valimaki M., Alfthan G., Vuoristo M., Ylikahri R. 7
High-performance liquid chromatographic determination of plasma glycolic acid in healthy subjects and in cases of hyperoxaluria syndromes
Petrarulo M., Marangella M., Linari F. 17
Influx of urea during bronchoalveolar lavage depends on the permeability of the respiratory membrane
Van de Graaf E.A., Jansen H.M., Weber J.A., Koolen M.G.J., Out T.A. 27
Letters to the Editor
Erythrocyte sodium, potassium and sodium fluxes with cell and subject ageing
Heseltine D., Thomas T.H., Wilkinson R., James O.F.W., Potter J.F. 41
Disparity between apolipoprotein E phenotypes and genotypes (as determined by polymerase chain reaction and oligonucleotide probes) in patients with non-insulin-dependent diabetes mellitus
Snowden C., Houlston R.S., Arif M.H., Laker M.F., Humphries S.E., Alberti K.G.M.M. 49
In vivo effect of simvastatin on lipoprotein cholesteryl ester metabolism in normocholesterolemic volunteers
Magot T., Malmendier C.L., Ouguerram K., Lontie J.F., Lutton C. 59

Vol. 196/2–3, 1991

Modulative effects of cell adhesion peptide (Arg-Gly-Asp-Ser) on the aggregation of stimulated platelets from ophthalmic patients
Babizhayev M.A., Mucha A.I. 77

Aminopeptidase P and dipeptidyl peptidase IV activity in human leukocytes and in stimulated lymphocytes
Hendriks D., De Meester I., Umiel T., Vanhoof G., Van Sande M., Scharpe S., Yaron A. 87
Application of sialidase antibodies for the diagnosis of clostridial infections
Roggentin P., Hobrecht R., Tirpitz D., Rothe B., Schauer R. 97
Restriction fragment length polymorphisms in the Apo B gene in relation to coronary heart disease in a Southern Asian population
Mendis S., Shepherd J., Packard C.J., Gaffney D. 107
Lipid peroxidation and protective enzymes during myocardial infarction
Loeper J., Goy J., Rozensztajn L., Bedu O., Moisson P. 119
Hydrogen sulphide and total acid-volatile sulphide in faeces, determined with a direct spectrophotometric method
Florin T.H.J. 127
Characterization of dipeptidyl and tripeptidyl aminopeptidases in human kidney soluble fraction
Mantle D. 135
A simple method for determination of serum catalase activity and revision of reference range
Goth L. 143
Influence of splenectomy on the properties of erythrocytes with Hemoglobin Volga disease
Falcioni G., Grelloni F., Bertoli E., Felici L., Valentino Coppa G. 153
Assessment of the automated colorimetric and the high-performance liquid chromatographic methods for nicotine intake by urine samples of smokers' smoking low- and medium-yield cigarettes
Kolonen S.A., Puhakainen E.V.J. 159
Clinical relevance of the quantification of apolipoprotein E in cerebrospinal fluid
Carlsson J., Armstrong V.W., Reiber H., Felgenhauer K., Seidel D. 167
The association between plasma β-hexosaminidase and its isoenzyme patterns and retinopathy in Type 1 diabetes mellitus
Hultberg B., Isaksson A., Agardh E., Agardh C.-D. 177
Serum laminin P1 fragment concentration in renal diseases
Horikoshi S., Koide H. 185
Purification and properties of human serum carnosinase
Jackson M.C., Kucera C.M., Lenney J.F. 193
Aminopeptidase A in human placenta and pregnant serum
Okuyama T., Ishiura S., Nojima M., Tsukahara T., Yanagida M., Sugita H. 207
Erythrocyte sodium in chronic liver disease with ascites
Ramakrishna B.S., Kumaresan R., Balasubramanian K.A. 217
4-Hydroxybutyric aciduria: Application of a fluorometric assay to the determination of succinic semialdehyde dehydrogenase activity in extracts of cultured human lymphoblasts
Gibson K.M., Lee C.F., Chambliss K.L., Kamali V., Francois B., Jaeken J., Jakobs C. 219

Vol. 197/1, 1991

The nature of the proteins lost from isolated rat skeletal muscle during experimental damage
Jackson M.J., Page S., Edwards R.H.T. 1
An immunologic pathway for intravascular catabolism of creatine kinase subform MM3
Kanemitsu F., Okigaki T. 9
A sensitive assay of transthyretin (prealbumin) in human cerebrospinal fluid in nanogram amounts by ELISA
Vatassery G.T., Quach H.T., Smith W.E., Benson B.A., Eckfeldt J.H. 19
Carnitine in dried blood spots: A method suitable for neonatal screening
Barns R.J., Bowling F.G., Brown G., Clague A.E., Thompson A. 27
Purification and characterization of leucyl aminopeptidase and pyroglutamyl aminopeptidase from human skeletal muscle
Mantle D., Lauffart B., Gibson A. 35

Modified hemoglobin in hypoglycemia caused by hyperinsulinism
Niketic V., Maric S., Diklic A., Neskovic S., Tomasevic N. 47
Collagen peptidase and type III procollagen peptide serum levels in chronic liver diseases
Muller A., Krombholz B., Pott G., Machnik G., Vollandt R., Reinhardt M., Jorke D. 59
Low density lipoprotein receptor levels and the response to dietary and pharmacological intervention in polygenic hypercholesterolemia
Joven J., Vilella E., Masana L., Ansoleaga J. 67
Porphobilinogen deaminase in acute intermittent porphyria: Activity and concentration in erythrocytes and lymphocytes
Lilius L., Lannfelt L., Wetterberg L., Floderus Y., Henrichson A., Thunell S. 77

Vol. 197/2, 1991

Serum ferritin spectrotypes in patients with heterozygous β-thalassaemia
Lasne Y., Francina A., Benzerara O. 85
The relationship between plasma free fatty acids and experimentally induced hepatic encephalopathy in the rat
Smit J.J.H., Bosman D.K., Jorning G.G.A., De Haan J.G., Maas M.A.W., Chamuleau R.A.F.M. 95
Determination of gangliosides and sulfatide in human cerebrospinal fluid with a microimmunoaffinity technique
Davidsson P., Fredman P., Mansson J.-E., Svennerholm L. 105
Low serum creatine kinase in patients with infective endocarditis
De Scheerder I.K., Delanghe J.R., De Buyzere M.L., Hollanders G., Clement D.L., Leroux-Roels G.G. 117
Measurement of serum Apo A1 and Apo B: Comparison of immunoturbidimetric and rate nephelometric techniques
Glueck C.J., McCray C., Speirs J. 123
Multiple forms of γ-glutamyl transpeptidase in human submandibular gland
Sajjan A.R., Hinchigeri S.B., Datta K.S. 133
Determination of serum iron by aas: Deproteinization versus direct analysis
De Benzo Z.A., Fraile R., Gomez C., Carrion N. 141
Letter to the Editor
Reference intervals for retinol-binding protein in serum and urine
Berg B., Fex G., Tryding N., Nilsson J.-E., Ekman S. 149
Erratum: One-step sandwich enzyme immunoassay for human laminin using monoclonal antibodies (Clinica Chimica Acta (1990) 191 (211–220))
Fujimoto N., Sejima H., Kyotani M., Iawata K., Ichida T., Ooshima A. 157

Vol. 197/3, 1991

An enzyme immunoassay for determining total thyroxine in human serum using an ultramicroanalytical system
Gonzalez R.R., Robaina R., Rodriguez M.E., Blanca S. 159
The presence and measurement of secretory component in human bile and blood
Perez J.H., Van Schaik M., Mullock B.M., Bailyes E.M., Price C.P., Luzio J.P. 171
Stability of human blood serum aminoacids after storage at different pH and temperature conditions
Kornhuber M.E., Balabanova S., Heiligensetzer G.V., Kornhuber C., Zettlmeissl H., Kornhuber A.W. 189
Modulation of cholesterol 7α-hydroxylase activity by nonspecific lipid transfer protein in human liver — possibly altered regulation of its cytosolic level in patients with gallstones
Kawata S., Imai Y., Inada M., Inui Y., Kakimoto H., Fukuda K., Maeda Y., Tarui S. 201

A sensitive, direct colorimetric assay of serum zinc using nitro-PAPS and microwell plates
Makino T. 209
Hypogalactosylation of immunoglobulin G sugar chains and elevated serum interleukin 6 in Castleman's disease
Nakao H., Nishikawa A., Nishiura T., Kanayama Y., Tarui S., Taniguchi N. 221
Intact parathyroid hormone assay is superior to mid region assay in the EDTA-infusion test in hyperparathyroidism
Bergenfelz A., Norden N.E., Ahren B. 229
Characterization of human plasma sialyltransferase using a novel fluorometric assay
Gross H.J., Brossmer R. 237
Quantitative analysis of urinary pyroglutamic acid in patients with hyperammonemia
Nakanishi T., Shimizu A., Saiki K., Fujiwara F., Funahashi S., Hayashi A. 249
Isoforms of human serum biotinidase
Hart P.S., Hymes J., Wolf B. 257
Delay in cell separation, storage and anticoagulants induced inaccuracies in measuring plasma non-esterified fatty acids
McGann A.M., Hodson A.W. 265

Vol. 198/1–2, 1991

Analysis of immunoglobulin and T cell receptor genes. Part I: Basic and technical aspects
Van Dongen J.J.M., Wolvers-Tettero I.L.M. 1
Analysis of immunoglobulin and T cell receptor genes. Part II: Possibilities and limitations in the diagnosis and management of lymphoproliferative diseases and related disorders
Van Dongen J.J.M., Wolvers-Tettero I.L.M. 93

Vol. 198/3, 1991

Ionselective minielectrode determination of ionic and total calcium concentrations in mixed saliva
Lagerlof F., Matsuo S. 175
An enzyme-linked lectin assay for sialidase
Lambre C.R., Terzidis H., Greffard A., Webster R.G. 183
Detection of the common Hb F Sardinia [(A)γ(E19)Ile → Thr] variant by isoelectric focusing in normal newborns and in adults affected by elevated fetal hemoglobin syndromes
Masala B., Manca L. 195
Glyceroluria in healthy adults, mentally ill adults and children selected for metabolic screening
Kohlschutter A., Seitz H.J., Feldmann B., Lehnert W., Langenbeck U. 203
Facts and artefacts in mevalonic aciduria: Development of a stable isotope dilution GCMS assay for mevalonic acid and its application to physiological fluids, tissue samples, prenatal diagnosis and carrier detection
Hoffmann G.F., Sweetman L., Bremer H.J., Hunneman D.H., Hyanek J., Kozich V., Lehnert W., Nyhan W.L., Speidel I., Trefz F.K., Gibson K.M. 209
Significance of variation in turnover of glycated albumin on indices of diabetic control
Johnson R.N., Easdale R.W., Tatnell M., Baker J.R. 229
Decreased oxidant-induced proteolysis in erythrocytes with enhanced antioxidative defence enzymes due to Down's syndrome
Bartosz G., Kedziora J., Retelewska W. 239
Accelerated in vitro degradation of CCK-58 in blood and plasma of patients with acute pancreatitis
Springer C.J., Eberlein G.A., Eysselein V.E., Schaeffer M., Goebell H., Calam J. 245
Consequences of the presence of elongated variant of the major transmembrane protein (Band 3 protein) in the human erythrocyte
Retelewska W., Gaczynska M., Bartosz G., Judkiewicz L. 255

Serum aluminium and zinc and other variables in patients with and without cognitive impairment in the community
Haines A., Iliffe S., Morgan P., Dormandy T., Wood B. 261
Changes in proteolytic susceptibility of erythrocyte membrane proteins in hereditary spherocytosis
Gaczynska M., Bartosz G., Judkiewicz L. 267
The effect of n-3 polyunsaturated fatty acids on Lp(a)
Berg Schmidt E., Klausen I.C., Kristensen S.D., Lervang H.-H., Faergeman O., Dyerberg J. 271
Letter to the Editor
Nomenclature standardization on an international basis
Fuentes-Arderiu X. 279

Vol. 199/1, 1991

Lysosomal enzyme activity in human aqueous humor
Weinreb R.N., Jeng S., Miller A.L. 1
Leukocyte sonicates as a source for both enzyme assay and DNA amplification for mutational analysis of certain lysosomal disorders
Louie E., Rafi M.A., Wenger D.A. 7
Screening for disorders of pyruvate metabolism by measuring the ratio of the rates of lactate production and pyruvate decarboxylation in cultured skin fibroblasts
Kuroda Y., Kawakami I., Kobashi H., Naito E., Ito M., Saijo T., Yokota I., Takeda E. 17
Determination of α-amylase using a new blocked substrate (3-ketobutylidene β-2-chloro-4-nitrophenyl-maltopentaoside)
Teshima S., Hayashi Y., Emi S., Ishimaru K. 23
High-performance liquid chromatographic determination of urinary cysteine and cystine
Birwe H., Hesse A. 33
Clinical evaluation of high-performance affinity chromatography for the separation of bone and liver alkaline phosphatase isoenzymes
Gonchoroff D.G., Branum E.L., Cedel S.L., Riggs B.L., O'Brien J.F. 43
Serum fructosamine/haemoglobin A_1 ratio predicts the future changes in haemoglobin A_1 in type 2 (non-insulin dependent) diabetic patients
MacRury S.M., Kilpatrick E.S., Paterson K.R., Dominiczak M.H. 51
Determination of membrane cholesterol in normal and pathological red blood cells
Macchia T., Mancinelli R., Barbini D.A., Taggi F., Avico U., Cantafora A. 59
Investigation of glycosylation processes in mitochondria and microsomal membranes from human skeletal muscle
Gasnier F., Lerme F., Rousson R., Roussouly P., Vaganay E., Louisot P., Gateau-Roesch O. 69
Stable isotope dilution assay for 5β-cholestane-3α,7α,12α,25-tetrol and 5β-cholestane-3α,7α,12α,23,25-pentol in human serum using [26,27-D_6] labeled internal standards; a highly accurate approach to the serological diagnosis of cerebrotendinous xanthomatosis
Kosaka D., Hiraoka T., Kohoda T., Kajiyama G., Yamauchi T., Kihira K., Kuramoto T., Hoshita T. 83
Simultaneous analysis of plasma phenethylamine, phenylethanolamine, tyramine and octopamine in patients with hepatic encephalopathy
Yonekura T., Kamata S., Wasa M., Okada A., Kawata S., Tarui S. 91
Quantitative analysis of human erythrocyte spectrin by sodium dodecyl sulfate-polyacrylamide gel electrophoresis using pancreatic trypsin inhibitor (Kunitz) as an internal standard
Yoshida H., Takamatsu S., Satoh K., Kawamura Y., Imaizumi T., Teshiromori R., Kawauchi K. 99
Letters to the Editor
Elevated serum bile acids in hepatic vein occlusion and idiopathic portal hypertension
Imai Y., Kawata S., Tarui S. 109
The effective inhibition of the Maillard reaction by guanabenz acetate, and its relevance to the prevention of diabetic complications
Igaki N., Yamada H., Oimomi M., Kasuga M. 113

Effect of temperature on red cell ATP
Beutler E. 117

Vol. 199/2, 1991

An enzyme-linked immunosorbent assay (ELISA) for the detection and quantitation of the tumor marker 1-methylinosine in human urine
D'Ambrosio S.M., Gibson-D'Ambrosio, Trewyn R.W. 119
Lipoprotein-cyclodextrin interaction
Sharma A., Janis L.S. 129
Platelet activating factor acetylhydrolase activity in the urine of patients with renal disease
Kirschbaum B. 139
Quantitation of serum apolipoprotein A-I, A-II, B, C-II, C-III and E in healthy Japanese by turbidimetric immunoassay: Reference values, and age- and sex-related differences
Noma A., Hata Y., Goto Y. 147
Ecdysteroid-like compounds in human urine: They can occur in the absence of any parasitic infection
Gharib B., Baswaid S., Quilici M., De Reggi M. 159
Changes in ceruloplasmin activity and lactate concentration in patients at high risk of acute organ system failure
Dauberschmidt R., Mrochen H., Forster I., Stumpe C., Dressler C., Grajetzki H., Meyer M. 167
Determination of medium chain acyl-CoA dehydrogenase activity in cultured skin fibroblasts using mass spectrometry
Niezen-Koning K.E., Chapman T.E., Mulder I.E., Smit G.P.A., Reijngoud D.J., Berger R. 173
N-Acetyl-β-D-glucosaminidase (NAG) isoenzymes release from human monocyte-derived macrophages in response to zymosan and human recombinant interferon-γ
Bourbouze R., Raffi F., Dameron G., Hali-Miraftab H., Loko F., Vilde J.-L. 185
Response of urinary angiotensin to challenges of the renin-angiotensin system
Boer P., Vos P.F., Koomans H.A., Rabelink A.J., Beutler J.J., Dorhout Mees E.J. 195
A gas chromatographic method for the determination of neutral steroid profiles in urine, including studies on the effect of oxytetracycline amdinistration on these profiles in men
Hamalainen E., Fotsis T., Adlercreutz H. 205

Vol. 199/3, 1991

Immunological studies of plasma cholinesterase during pregnancy and the puerperium
Whittaker M., Jones J.W., Braven J. 223
Characteristics of hexosaminidase A in homogenates of white blood cells using methylumbelliferyl-N-acetyl-β-D-glucosaminide-6-sulphate as substrate
Den Tandt W.R., Scharpe S. 231
O-(2,3,4,5,6-Pentafluorobenzyl)oxime-trimethylsilyl ester derivatives for sensitive identification and quantitation of aldehydes, ketones, and oxoacids in biological fluids
Hoffmann G.F., Sweetman L. 237
Affinity electrophoresis of alkaline phosphatase using polyacrylamide gels
Ramasamy I. 243
Spectrofluorimetric determination of urinary p-aminobenzoic and p-aminosalicylic acids in the BT-PABA/PAS test of pancreatic function
Pemberton P.W., Gagjee P., Chaloner C., Braganza J.M., Lobley R.W. 253
Isotope dilution gas chromatographic-mass spectrometric method for the determination of lignans and isoflavonoids in human urine, including identification of genistein
Adlercreutz H., Fotsis T., Bannwart C., Wahala K., Brunow G., Hase T. 263
Agarose gel electrophoresis of glucose-6-phosphate-dehydrogenase isoenzymes
Forbes J., Steytler J.G., Van Heerden R. 279

Studies on the effects of aminopterin and methotrexate on the release of prostaglandin PGE_2, PGD_2, and PGI_2 from small intestine in the rat in vivo
Weiler H., Moser U., Weiler C., Gerok W. 283
Bile acid and phospholipid fatty acid composition in bile of patients with cholesterol and pigment gallstones
Van Erpecum K.J., Van Berge Henegouwen G.P., Stolk M.F.J. 295
A photometric assay for hydrogen peroxide production by polymorphonuclear leucocytes
Negri M., Bellavite P., Lauciello C., Guzzo P., Zatti M. 305
Elevated serum β-glucuronidase activity in acquired immunodeficiency syndrome
Saha A.K., Glew R.H., Kotler D.P., Omene J.A. 311

Vol. 200/1, 1991

Atherogenic risk factors in cerebrotendinous xanthomatosis
Fujiyama J., Kuriyama M., Arima S., Shibata Y., Nagata K., Takenaga S., Tanaka H., Osame M. 1
Measurement of plasma free steroids by direct radioimmunoassay of ultrafiltrate in association with the monitoring of free components with [^{14}C]glucose
Adachi K., Yasuda K., Fuwa Y., Goshima E., Yamakita N., Miura K. 13
Serum release of the creatine kinase tissue-specific isoforms MM3 and MB3 is simultaneous during myocardial reperfusion
Christenson R.H., Magnus Ohman E., Vollmer R.T., Clemmensen P., Grande P., Wagner G.S. 23
A new sensitive method for determining endotoxin in whole blood
Tamura H., Tanaka S., Obayashi T., Yoshida M., Kawai T. 35
A new mutant erythrocyte pyrimidine 5′-nucleotidase characterized by fast electrophoretic mobility in a Chinese boy with chronic hemolytic anemia
Li J.-Y., Wan S.-D., Ma Z.-M., Zhao Y.-H., Zhou G.-P. 43
Letters to the Editor
A universal eluent system for free amino acid analysis in biological fluids
Pospisil R., Valik D. 49
Macro creatine kinase type 2 in a patient with prostatic carcinoma
Arenas J., Bornstein B., Mayor J.J., Santos I., Leiva O., Martinez A. 53
Effects of dimethyl sulfoxide (DMSO) on carbon (CCL_4)-induced hepatotoxicity in mice
Achudume A.C. 57
A simplified analysis of fatty acids in serum phospholipids using Sep-Pak cartridges
Ellin A., Van den Berg S., Strandvik B. 59
Measurement of total fasting-state serum bile acids: Comparison of a solid-phase bioluminescence enzymatic assay, a homogeneous fluorescence enzymatic assay and isotope dilution gas chromatography-mass spectrometry
Rossi S.S., Angellotti M.A., Setchell K.D.R., Hofmann A.F. 63
Suppression/stimulation of chemiluminescence by human seminal plasma
Haq A., Sheth K., Abdullatif M., Jabbar F.A.-A., Hamilton C., Al-Sedairy S. 67

Vol. 200/2–3, 1991

β-N-Acetylhexosaminidases in human cerebrospinal fluid and serum of patients with multiple sclerosis
Datti A., Emiliani C., Capocchi G., Orlacchio A. 73
CA 15-3 and carcinoembryonic antigen in the clinical evaluation of breast cancer
Dnistrian A.M., Schwartz M.K., Greenberg E.J., Smith C.A., Schwartz D.C. 81
A sensitive fluorescence assay for quantitation of fludarabine and metabolites in biological fluids
Kemena A., Fernandez M., Bauman J., Keating M., Plunkett W. 95
Pyrophosphate inhibition of Proteus mirabilis-induced struvite crystallization in vitro
McLean R.J.C., Downey J., Claphan L., Wilson J.W.L., Nickel J.C. 107

Mechanized assay of plasma prekallikrein by activation with Pseudomonas aeruginosa elastase and amidolysis of chromogenic substrate
Shibuya Y., Yamamoto T., Tanaka H., Semba U., Kambara T., Okabe H. 119
Human aortic elastin from normal individuals and atherosclerotic patients: Lipid and cation contents; susceptibility to elastolysis
Saulnier J.M., Hauck M., Fulop Jr. T., Wallach J.M. 129
Development of enzyme-linked immunosorbent assay for free human pro-colipase activation peptide (APGPR)
Bowyer R.C., Jehanli A.M.T., Patel G., Hermon-Taylor J. 137
Biochemical properties of blood cells from histiocytosis X patients
Perlino E., Marra E., Petragallo V.A., Quagliariello E., De Terlizzi M., Santostasi T., Ceci A. 153
Stability of platelet activating factor (PAF) in human saliva. Quantitation by radioimmunoassay
Cooney S.J., Smal M.A., Baldo B.A. 161
Immunological assay of erythrocyte acetylcholinesterase
Jones J.W., Whittaker M., Braven J. 175
Absence of increased urinary excretion of adenosine-deaminase-binding protein by patients with chronic renal tubular malfunction
Flynn F.V., Lapsley M., Sansom P.A., Norden A.G.W. 183
Serum amyloid P component in chronic renal failure and dialysis
Nelson S.R., Tennent G.A., Sethi D., Gower P.E., Ballardie F.W., Amatayakul-Chantler S., Pepys M.B. 191
Membrane properties of the sarcolemma and sarcoplasmic reticulum of pigs susceptible to malignant hyperthermia. Action of halothane
Thomas M.A., Rock E., Viret J. 201
Increased elastase-α_1-antitrypsin complex in fulminant hepatic failure: Relationship to bacterial infection and activation of coagulation
Langley P.G., Hughes R.D., Rolando N., Williams R. 211
Letters to the Editor
Human liver vasopressin receptors
Howl J., Strain A.J., Wheatley M. 221
Usefulness of faecal water potassium/sodium ratios in the investigation of diarrhoea
Duncan A., Robertson C., Russell R.I. 223
Impossibility to study crystalluria directly with Coulter Counter
Hennequin C., Gaubil J., Bader C., Daudon M., Drueke T., Lacour B. 227
Glycohistones in diabetic human liver
Jobst K., Lakatos A., Horvath A. 231

Vol. 201/1–2, 1991

Neopterin in clinical practice
Muller M.M., Curtius H.-C., Herold M., Huber C.H. 1
Analysis of bismuth in serum and blood by electrothermal atomic absorption spectrometry using platinum as matrix modifier
Slikkerveer A., Helmich R.B., Edelbroek P.M., Van der Voet G.B., De Wolff F.A. 17
Bone-alkaline phosphatase as indicator of bone formation
Van Straalen J.P., Sanders E., Prummel M.F., Sanders G.T.B. 27
High-performance liquid chromatographic determination of bilirubin distribution on serum proteins
Soltes L., Sebille B., Fugedi A. 35
Clinical analysis in intact erythrocytes using ^{1}H spin echo NMR
Reglinski J., Smith W.E., Wilson R., Buchanan L.M., McKillop J.H., Thomson J.A., Brzeski M., Marabani M., Sturrock R.D. 45
Changes in fucose metabolism associated with heavy drinking and smoking: A preliminary report
Thompson S., Matta K.L., Turner G.A. 59

Differential low density lipoprotein hydrated density distribution in female and male patients with familial hypercholesterolemia
Tilly-Kiesi M.K., Tikkanen M.J. 65
Increased plasma cyclic guanosine monophosphate concentration in chronic obstructive pulmonary disease patients with increased circulating atrial natriuretic peptide
Rota M., Brulot N., Rezgui N., Labrousse J., Allard D. 75
Reference values for free gamma-aminobutyric acid determined by ion-exchange chromatography and fluorescence detection in the cerebrospinal fluid of children
Carchon H.A., Jaeken J., Jansen E., Eggermont E. 83
Pitfalls in the determination of diamine oxidase activity
Smith C.H., Maghsoudloo M., Himuna K., Murphy G.M. 89
Enzymatic phenylalanine estimation for the management of patients with phenylketonuria
Wendel U., Koppelkamm M., Hummel W. 95
HbA(1C) and serum fructosamine in diabetic patients: Relationship to age, clotting and fibrinolysis parameters and urinary microalbumin excretion
Van Wersch J.W.J., Westerhuis L.W.J.J.M., Venekamp W.J.R.R. 99
Detection of an eosinophil derived neurotoxin in the urine of a patient with idiopathic chronic eosinophilic pneumonia
Deviller P., Gruart V., Prin L., Tardy F., Bulle C., Capron M., Capron A., Cordier J.-F. 105
The common finding of covert hypothyroidism at initial clinical evaluation for hyperlipoproteinemia
Glueck C.J., Lang J., Tracy T., Speirs J. 113
Diurnal and postural variation in serum α_1 microglobulin in normal individuals
Itoh Y., Kawai T. 123
Hyperornithinemia, hyperammonemia and homocitrullinuria — A case report and study of ornithine metabolism using in vivo deuterium labelling
Tsujino S., Suzuki T., Azuma T., Higa S., Sakoda S., Kishimoto S. 129
Letters to the Editor
Nomenclature of hormones used as drugs for biochemical function tests
Fuentes-Arderiu X. 135
Instability of fecal lysozyme
Brouwer J., Smekens F. 137
Serum beta-2-microglobulin levels and cognitive function in chronic dialysis patients
Annunziata P., Biondi B., Aucone A.M., Poggi A., Moschini F. 139
Measurement of intact parathormone in hyperthyroidism
Roiter I., Da Rin G., Galliazzo S., Legovini P., Di Virgilio R., Foscolo G., Breda F., Conte N. 143
Diagnosis of organic acidemia in developing countries
Lin Y.Y., Wisniewski K.E., Inoue Y., Matsumoto I., Schmidt B.J., Raiman E., Cornejo V. 145
Aminopeptidases in the soluble fraction of human placenta
Mizutani S., Nomura S., Kurauchi O., Kasugai M., Ino K., Goto S., Tomoda Y. 151

Vol. 201/3, 1991

Improved synthesis of [1-^{14}C]acyl-sphingosine-galactose-3-sulfate (sulfatide) for diagnosis of metachromatic leukodystrophy: Usefulness of radioscanning
Masson M., Li W.X., Fluharty A.L., Beaucourt J.P., Turpin J.C., Baumann N. 157
Some aspects of the use of dextran sulfate-cellulose columns for the treatment of familial hypercholesterolemia
Erlich D., Aubert I., Neel D., Scrobohaci M.L., Loreau J., Bussel A., Rouffy J., Dreux C., Goussault Y., Weintraub H. 169
A radiometric assay for aspartoacylase activity in human fibroblasts: Application for the diagnosis of Canavan's disease
Barash V., Flhor D., Morag B., Boneh A., Elpeleg O.N., Gilon C. 175
Sensitive enzyme immunoassay for human 28 kDa calbinin-D
Zhu Y., Takashi M., Miyake K., Kato K. 183

Brain natriuretic peptides in human plasma, spinal cord and cerebrospinal fluid
Togashi K., Kameya T., Ando K., Marumo F., Kawakami M. 193
Anti-skeletal muscle and anti-acetylcholine receptor antibodies in patients with thymoma without Myasthenia gravis: Relation to the onset of Myasthenia gravis
Ohta M., Itoh M., Hara H., Itoh N., Nishitani H., Hayashi K., Ohta K. 201
Identification of human ventricular myosin heavy chain fragments with monoclonal antibody 2F4 in human sera after myocardial necrosis
Simeonova P.P., Kehayov I.R., Kyurkchiev S.D. 207
Immunonephelometric assay of myoglobin
Massoubre C., Chivot L., Mainard F., Bridji B., Madec Y. 223
Determination by latex immunoassay of protein 1 in normal and pathological urine
Bernard A., Lauwerys R., Noel A., Vandeleene B., Lambert A. 231
Examination of the role of oxygen free radicals in bronchial asthma in childhood
Novak Z., Nemeth I., Gyurkovits K., Varga S.I., Matkovics B. 247

Vol. 202/1–2, 1991

Quantitation of urobilinogen in feces, urine, bile and serum by direct spectrophotometry of zinc complex
Kotal P., Fevery J. 1
Investigation and validation of the affinity chromatography method for measuring glycated albumin in serum and urine
Silver A.C., Lamb E., Cattell W.R., Dawnay St. B.J. 11
A competitive radioimmunoassay for peptidoglycan monomer
Vranesic B., Ljevakovic D., Tomasic J., Ladesic B. 23
In thiamine deficiency, activation of erythrocyte transketolase by thiamine in vivo exceeds activation by cofactor in vitro
Price J., Clague A.E., Kerr R.A., Nixon P.F. 39
Comparison of normal subjects and asthma patients with respect to age-related elastase-binding activity of α_2-macroglobulin in plasma
Kilroe-Smith T.A., Becker P.J., Gaillard M.C. 47
α_2-Macroglobulin and generation of oxygen radicals by granulocytes: Potential role in prevention and treatment of reperfusion injury
Donnelly P.K., Booth H., White M., Shenton B.K. 55
Microheterogeneity of α1-acid glycoprotein in healthy elderly subjects: Patterns obtained by crossed affino-immunoelectrophoresis
Kawerk N., Succari-Aderschlag M., Foglietti M.-J. 65
Serum F-protein concentration following halothane or isoflurane anaesthesia
Cooper C.M., Foster G.R., Cooper A.E., Kenny M., Olivera D.B. 73
Comparison of sampling systems for the collection of $^{14}CO_2$ in ^{14}C breath tests
Peled Y., Gilat T. 83
Skeletal muscle cytochrome c in X-linked muscle dystrophies
Nigro G., Comi L.I., Politano L., Limongelli F.M. 89
Platelet-activating factor acetylhydrolase in plasma lipoproteins of healthy men and women
Satoh K., Imaizumi T.-A., Yoshida H., Kawamura Y., Takamatsu S., Takamatsu M., Mizuno S. 95
Carbon monoxide production from L-3,4-dihydroxyphenylalanine; a method for assessing the oxidant/antioxidant properties of drugs
Cooper M.J., Engel R.R. 105
Plasma lactoferrin levels after bone marrow transplantation monitored by a two-site enzyme immunoassay
Suzuki T., Takizawa-Mizuno M., Yazaki M., Wada Y., Asai K., Kato T. 111
Letter to the Editor
Lactate dehydrogenase isoenzymes in serum of patients with preeclampsia/eclampsia complicated by the HELLP syndrome
Beyer C. 119

IFCC Recommendation — Theory of reference values. Part 4. Control of analytical variation in the production, transfer and application of reference values
Solberg H.E., Stamm D. S5

IFCC Recommendation — Recommendation on sampling, trasnport and storage for the determination of the concentration of ionized calcium in whole blood, plasma and serum
Boink A.B.T.J., Buckley B.M., Christiansen T.F., Covington A.K., Maas A.H.J., Muller-Plathe O., Sachs Ch., Siggaard-Andersen O. S13

Guidelines (1990) for selection of safe laboratory centrifuges and for their safe use. With general purpose appendices concerning centrifuge nomenclature, quantities and units, and calculation of centrifugal acceleration
Uldall A., Damgaard P.T., Drachmann O., Jorgensen F., Kennedy D., Lauritzen M., Magnussen E., Rigg J.C., Voss P. S23

Vol. 202/3, 1991

Accumulation and impaired in vivo metabolism of di- and trihydroxycholestanoic acid in two patients
Wanders R.J.A., Casteels M., Mannaerts G.P., Van Roermund C.W.T., Schutgens R.B.H., Kozich V., Zeman J., Hyanek J. 123

Body fat distribution and serum lipoproteins in relation to age and body weight
Sonnichsen A.C., Richter W.O., Schwandt P. 133

Raised urinary excretion of inorganic pyrophosphate in asymptomatic members of a hypophosphatasia kindred
Macfarlane J.D., Poorthuis B.J.H.M., Mulivor R.A., Caswell A.M. 141

Arylsulfatases A and B in EBV-transformed lymphoid cell lines: Studies on their molecular forms in cells from patients with inborn sulfatase deficiencies. Comparative diagnostic value of enzymatic assays
Tempesta M.-C., Levade T., Salvayre R. 149

A rapid fluorometric enzyme immunoassay for the determination of neonatal TSH from blood spots
Tuuminen T., Rakkolainen A.E.M., Welin M.-G.I., Weber T.H., Nylander P.L., Kapyaho K.I. 167

Sandwich enzyme immunoassay for serum integrins using monoclonal antibodies
Katayama M., Kurome T., Yamamoto K., Uchida H., Hino F., Kato I. 179

A new immunochemical assay for biotin
Thuy L.P., Sweetman L., Nyhan W.L. 191

Isolation of amniotic fluid proteins of non-maternal serum origin by negative immuno-affinity chromatography
Sorensen S. 199

Effect of alcohol on δ-aminolevulinic acid dehydratase and porphyrin metabolism in man
Sieg I., Doss M.O., Kandels H., Schneider J. 211

ELISA using monoclonal antibody to human serum arylesterase
Kawai H., Yomoda S., Inoue Y. 219

A new enzymatic assay of urinary guanidinoacetic acid
Shirokane Y., Nakajima M., Mizusawa K. 227

Role of iron in the photosensitization by uroporphyrin
Menon I.A., Becker M.A.C., Haberman H.F. 237

Radioimmunoassay of growth hormone-releasing hormone (GHRH) with a polyclonal antibody against synthetic GHRH(1-29)-Gly_4-Cys-NH_2: Method and clinical studies
Zhang C., Yamasaki R., Mitsuhashi S., Takahashi H., Bando H., Saito S. 243

Vol. 203/1, 1991

Isotransferrins and pregnancy: A study in the guinea pig
Van Dijk J.P., Van Noort W.L., Kroos M.J., Starreveld J.S., Van Eijk H.G. 1

Marked variation in blood beta-hexosaminidase in Gaucher disease
Natowicz M.R., Prence E.M., Cajolet A. 17
Specific diagnosis of medium-chain acyl-CoA dehydrogenase (MCAD) deficiency in dried blood spots by a polymerase chain reaction (PCR) assay detecting a point-mutation (G985) in the MCAD gene
Gregersen N., Blakemore A.I.F., Winter V., Andresen B., Kolvraa S., Bolund L., Curtis D., Engel P.C. 23
Interest of zinc determination in leucocyte fractions for the assessment of marginal zinc status
Peretz A., Neve J., Jeghers O., Leclercq N., Praet J.-P., Vertongen F., Famaey J.-P. 35
Aminoterminal propeptide of type III procollagen in cerebrospinal fluid. Variation with age and in childhood leukemia
Vainionpaa L., Risteli L., Lanning M., Myllyla V., Risteli J. 47
Increased concentrations of endogenous 13-cis- and all-trans-retinoic acids in diffuse idiopathic skeletal hyperostosis, as demonstrated by HPLC
Periquet B., Lambert W., Garcia J., Lecomte G., De Leenheer A.P., Mazieres B., Thouvenot J.P., Arlet J. 57
Flow injection system based on the sandwich technique for saving expensive reagents
Araujo A.N., Lima J.L.F.C., Alonso-Chamarro J., Bartroli J., Poch M. 67
Synthesis of ^{13}C-labeled chenodeoxycholic, hyodeoxycholic, and ursodeoxycholic acids for the study of bile acid metabolism in liver disease
Matern S., Marschall H.-U., Schill A., Schumacher B., Lehnert W., Sjovall J., Matern H. 77
Letters to the Editor
Refinement of a specific assay for hydroxyproline in urine
Sela B.-A., Doolman R. 91
Rapid quantitation of serum myoglobin by latex agglutination turbidimetry
Abe J., Yamaguchi T., Isshiki T., Naka H., Kawada Y., Katayama Y. 95
A discrepancy between in vivo and in vitro lipoprotein lipase activity in a patient with Tangier disease
Takizawa A., Komoda T., Hokari S., Sakagishi Y., Tanaka A. 97
Wheat germ agglutinin binding to serum 5′-nucleotidase and alkaline phosphatase
Novo F.J., Tutor J.C. 101
Further data on an RIA kit for the direct measurement of free testosterone in plasma
Rivera-Coll A., Navarro M.A. 103
Micromethod for the assay of glutamate: Glyoxylate aminotransferase and modifications of a micromethod for the assay of alanine: Glyoxylate aminotransferase. Implications for the prenatal diagnosis of Type I hyperoxaluria by fetal liver biopsy
Toone J.R., Applegarth D.A. 105

Vol. 203/2–3, 1991

Effect of simvastatin on the apparent size of LDL particles in patients with type IIB hyperlipoproteinemia
Zhao S.P., Hollaar L., Van 'T Hooft F.M., Smelt A.H.M., Leuven J.A.G., Van Der Laarse A. 109
Energy metabolism during damaging contractile activity in isolated skeletal muscle: A ^{31}P-NMR study
West-Jordan J.A., Martin P.A., Abraham R.J., Edwards R.H.T., Jackson M.J. 119
Abnormal serum immunoglobulin concentrations in patients with diabetes mellitus
Rodriguez-Segade S., Camina M.F., Paz J.M., Del Rio R. 135
Phosphoribosylpyrophosphate synthetase in human erythrocytes: Assay and kinetic studies using high-performance liquid chromatography
Sakuma R., Nishina T., Yamanaka H., Kamatani N., Nishioka K., Maeda M., Tsuji A. 143
Antibodies to dietary antigens in rheumatoid arthritis — Possible molecular mimicry mechanism
Perez-Maceda B., Lopez-Bote J.P., Langa C., Bernabeu C. 153

Changes in α_1-acid glycoprotein serum concentrations and glycoforms in the developing human fetus
Seta N., Tissot B., Forestier F., Feger J., Daffos F., Durand G. 167
Determination of apolipoprotein E phenotypes from stored or postmortem serum samples
Lehtimaki T. 177
Detection of the most common mutation of adenine phosphoribosyltransferase deficiency among Japanese by a non-radioactive method
Kawaguchi R., Higashimoto H., Hikiji K., Hakoda M., Kamatani N. 183
Rapid automated quantitation of isoleucine, leucine, tyrosine and phenylalanine from dried blood filter paper specimens
Qu Y., Miller J.B., Slocum R.H., Shapira E. 191
Urinary oligosaccharide excretion and severity of galactosialidosis and sialidosis
Takahashi Y., Nakamura Y., Yamaguchi S., Orii T. 199
High-performance liquid chromatography of urinary oligosaccharides in the diagnosis of glycoprotein degradation disorders
Hommes F.A., Varghese M. 211
Assay for type III collagenolytic activity in lung cancer tissue
Kawamura M., Kato R., Kikuchi K., Kobayashi K., Ishihara T., Shibata T., Inayama S. 225
In vitro stability of human atrial natriuretic peptide (h-ANP)
Artner-Dworzak E., Lindner H., Puschendorf B. 235
The glycoconjugate content of human faeces and ileostomy effluents
Clamp J.R., Gough M. 243
Phospholipase A_2 is associated with albumin in patients with acute pancreatitis
Langton S.R., Dench J. 249
A re-evaluation of the lipid-bound sialic acid determination
Vegh Z., Kremmer T., Boldizsar M., Gesztesi K.A., Szajani B. 259
Deoxycholic acid is not reconverted to cholic acid in humans — A study by isotope ratio mass spectrometry
Fischer S., Beuers U., Berr F., Paumgartner G. 269
Assay of plasma glycine by HPLC with electrochemical detection in patients undergoing glycine irrigation during gynaecological surgery
Sherwood R.A., Bayliss E.M., Chappatte O. 275
Urinary hydroxyproline and serum alkaline phosphatase in sickle cell disease
Mohammed S.M., Suleiman S.A., Addae S.K., Annobil S.H., Adzaku F.K., Kadoummi O.F., Richards J.T. 285
Rapid and sensitive discriminating determination of acetylcholinesterase activity in amniotic fluid with a choline sensor
Morelis R.M., Coulet P.R., Simplot A., Boisson C., Guibaud G. 295
Serum levels of human alkaline phosphatase isozymes in relation to blood groups
Domar U., Hirano K., Stigbrand T. 305
Glycosylation of α_1-acid glycoprotein in relation to duration of disease in acute and chronic infection and inflammation
Fassbender K., Zimmerli W., Kissling R., Sobieska M., Aeschlimann A., Kellner M., Muller W. 315
Effect of acute hyperinsulinemia on fatty acid composition of serum lipids in non-insulin-dependent diabetics and healthy men
Pelikanova T., Kohout M., Base J., Stefka Z., Kovar J., Kazdova L., Valek J. 329
Azopigments from serum biliprotein and accuracy of total bilirubin measurement
Franzini C., Cattozzo G. 339
Lysosomal enzymes and severity of saphen varicosis
Hayer M., Caporiccio A., Pages A., Caporiccio G., Toueri M.F., Magnan de Bornier B. 343
The identification of type 1 Gaucher disease patients, asymptomatic cases and carriers in The Netherlands using urine samples: An evaluation
Aerts J.M.F.G., Sa Miranda M.C., Wanzeller De Lacerda L., Van Weely S., Donker-Koopman W., Brouwer-Kelder B., Jansen D.C., Van Leeuwen M., Schram A.W., Tsiapara A., Tager J.M. 349

Elevated plasma histamine in sickle cell anaemia
Enwonwu C.O., Lu M. 363
Prenatal diagnosis of isovaleric acidemia by fast atom bombardment and tandem mass spectrometry
Shigematsu Y., Kikawa Y., Sudo M., Kanaoka H., Fujioka M., Dan M. 369
Serum gamma glutamyl transpeptidase in head and neck cancer
Singh J., Sharma A., Yadev S.P.S., Lal H. 375
Variations in relative concentrations of variants of human α_1-acid glycoprotein after acute-phase conditions
Eap C.B., Fischer J.-F., Baumann P. 379
Identification of renal calculi by their Raman spectra
Sudlow K., Woolf A. 387
Measurement of ornithine carbamyl transferase (OCT) in plasma by means of enzymatic determination of ammonia
Peltenburg H.G., Janssen M.A., Soeters P.B., Flendrig J.G., Hermens W.T. 395
Letters to the Editor
Isoelectric focusing of neutrophil alkaline phosphatase in trisomy 21 pregnancies
Brisson-Lougarre A., Vergness H., Grozdea J., Bourrouillou G., Colombies P. 403
A practical new assay for determining N-acetyl-β-D-glucosaminidase activity in urine
Jung K., Priem F. 405
Biochemical properties of γ-glutamyl transpeptidase in human submandibular gland
Sajjan A.R., Hinchigeri S.B., Datta K.S. 409
Improved method for the determination of the magnesium concentration of mononuclear blood cells
Nozue T., Kodama T., Kobayashi A. 411
Level is not a synonym of concentration
Fuentes-Arderiu X. 417

The effects of common matrices for assay standards on performance of 'ultra sensitive' immunometric assays for TSH. Report of a joint WHO/IFCC collaborative study
Gaines-Das R.E., Brettschneider H., Bristow A.F. S5

Vol. 204/1–3, 1991

Alterations of the glycan moiety of human α1-acid glycoprotein in late-term pregnancy
Biou D., Bauvy C., N'Guyen H., Codogno P., Durand G., Aubery M. 1
Strenuous physical activity, aspirin and heat stress increase urinary dolichols: Evidence for lysosomal origin of urinary dolichols
Roine R., Heinonen T., Salmela K., Heikkonon E., Suokas A., Luurila O.J., Koskinen P., Palo J., Salaspuro M. 13
Effectiveness of palladium matrix modification for the determination of thallium by graphite furnace atomic absorption spectrometry
Yang Q., Smeyers-Verbeke J. 23
In vitro glycation of human apoliprotein A(I) reduces its efficiency in lecithin:cholesterol acyltransferase activation
Gugliucci A., Stahl A.J.C. 37
Determination of protein-bound urinary gamma-carboxyglutamic acid in calcium nephrolithiasis
Colette C., Benmbarek A., Boniface H., Astre C., Pares-Herbute N., Monnier L., Guitter J. 43
The adrenergic control of lipolysis: Development of tolerance to beta-2 receptor stimulation with terbutaline
Kendall M.J., Hafner C.A., Hughes B., Florkowski C. 51
Predisposition to LDL oxidation during copper-catalyzed oxidative modification and its relation to α-cotopherol content in humans
Cominacini L., Garbin U., Cenci B., Davoli A., Pasini C., Ratti E., Gaviraghi G., Lo Cascio V., Pastorino A.M. 57

Arylsulfatase A in serum from patients with cancer of various organs
Laidler P., Kowalski D., Silberring J. 69
Gas chromatography method for the separation of amino acids enantiomers in plasma and urine. Application in a case of short bowel syndrome
Ketting D., Wadman S.K., Spaapen L.J.M., Van Der Meer S.B., Duran M. 79
Urine thiocyanate:creatinine ratio as a reliable indicator of cigarette smoking
Pre J., Vassy R. 87
Cellular and humoral responses in coeliac disease. 1. Wheat protein fractions
Penttila I.A., Devery J.M., Gibson C.E., LaBrooy J.T., Skerritt J.H. 95
Cellular and humoral responses in coeliac disease. 2. Protein extracts from different cereals
Skerritt J.H., Devery J.M., Penttila I.A., Labrooy J.T. 109
Influence of serum proteins on fructosamine concentration in multiple myeloma
Montagna M.P., Laghi F., Cremona G., Zuppi C., Barbaresi G., Castellana M.L. 123
3-Hydroxy-3-methylglutaric aciduria: A possible pitfall in diagnosis
Mills G.A., Hill M.A.W., Buchanan R., Corina D.L., Walker V. 131
Optimisation of non-denaturing polyacrylamide gel electrophoretic analysis of glyoxalase I phenotypes in clinical blood samples
McLellan A.C., Thornalley P.J. 137
The nature of small bowel luminal fluid lactase
Quak S.H., Brown G.A., Booth I.W., McNeish A.S. 145
Composition of very low density lipoproteins and in vitro effect of lipoprotein lipase
Saheki S., Takahashi I., Murase M., Takeuchi N., Uchida K. 155
Fatty acid composition of hepatic triglycerides in Reye's syndrome: Implications for hepatic desaturase abnormalities
Kang E.S., Johnson S.B., Holman R.T. 167
Purification and structural study of two albumin variants in an Irish population
Sakamoto Y., Davis E., Madison J., Watkins S., McLaughlin H., Leahy D.T., Putnam F.W. 179
Capillary plasma lipid profiles
Mace R., May B., McCray C., Tracy T., Speirs J., Glueck C.J. 189
Comparison of nonparametric recursive partitioning to parametric discriminant analyses in laboratory differentiation of hypercalcemia
Lacher D.A. 199
Ethinyl estradiol and norethindrone radioimmunoassay following Sephadex LH-20 column chromatography
Zacur H.A., Linkins S., Chang V., Smith B., Kimball A.W., Burkman R. 209
Assay of inulin in biological fluids by high-performance liquid chromatography with pulsed amperometric detection
Ruo T.I., Wang Z., Dordal M.S., Atkinson Jr. A.J. 217
Creatinine deiminase (EC 3.5.4.21) from bacterium BN11: Purification, properties and applicability in a serum/urine creatinine assay
Gottschalk E.M., Hippe H., Patzke F. 223
Betaine:homocysteine methyltransferase — A new assay for the liver enzyme and its absence from human skin fibroblasts and peripheral blood lymphocytes
Wang J., Dudman N.P.B., Lynch J., Wilcken D.E.L. 239
Platelet aggregation and fatty acid composition of platelets in type 1 diabetes mellitus
Myrup B., Bregengaard C., Petersen L.R., Winther K. 251
Carnitine therapy and metabolism in the disorders of propionyl-CoA metabolism studies using ^{1}H-NMR spectroscopy
Davies S.E.C., Iles R.A., Stacey T.E., De Sousa C., Chalmers R.A. 263
Increased susceptibility to peroxide-induced haemolysis with normal vitamin E concentrations in cystic fibrosis
James D.R., Alfaham M., Goodchild M.C. 279
Serum Lp(a) concentrations are unaffected by treatment with the HMG-CoA reductase inhibitor Pravastatin: Results of a 2-year investigation
Fieseler H.-G., Armstrong V.W., Wieland E., Thiery J., Schutz E., Walli A.K., Seidel D. 291

Plasma bromine concentrations and lipid profiles
Corrigan F.M., Skinner E.R., MacIntyre F., Ijomah G., Holliday J., Horrobin D.F., Ward N.I. 301
Letter to the Editor
Isoelectric focusing of neutrophil alkaline phosphatase in trisomy 21 pregnancies
Brisson-Lougarre A., Vergnes H., Grozdea J., Bourrouillou G., Colombies P. 305

Vol. 205/1–2, 1992

Generation of analytic plasma lipoprotein profiles using two prepacked superose 6B columns
Gerdes L.U., Gerdes C., Klausen I.C., Faergeman O. 1
High-performance liquid chromatographic determination of free polyamines in human seminal plasma
Oefner P.J., Wongyai S., Bonn G. 11
Discriminant analysis of urinary calculi by near-infrared reflectance spectroscopy
Peuchant E., Heches X., Sess D., Clerc M. 19
Tubular secretion of Tamm-Horsfall protein in type 1 (insulin-dependent) diabetes mellitus using a simplified enzyme linked immunoassay
Torffvit O., Agardh C.-D., Kjellsson B., Wieslander J. 31
Influences of diet and surgical trauma on serum alkaline DNase activity levels
Avall Lundqvist E.H., Sjovall K., Eneroth P.H.E. 43
Limitations of the triolein breast test
Duncan A., Cameron A., Stewart M.J., Russell R.I. 51
An enzyme-linked immunosorbent assay for heparin cofactor II (HCII). Application to the measurement of HCII in clinical materials
Toulon P., Costa J.M., Amiral J. 65
Selenium and glutathione peroxidase variations induced by polyunsaturated fatty acids oral supplementation in humans
Bellisola G., Galassini S., Moschini G., Poli G., Perona G., Guidi G. 75
Characterization of 6-hexadecanoylamino-4-methylumbelliferyl-β-D-galactopyranoside as fluorogenic substrate of galactocerebrosidase for the diagnosis of Krabbe disease
Wiederschain G., Raghavan S., Kolodny E. 87
Effects of unsaturated and saturated dietary plant sterols on their serum contents
Vanhanen H.T., Miettinen T.A. 97
Homocarnosinosis patients and great apes have a serum protein that cross-reacts with human serum carnosinase
Jackson M.C., Lenney J.F. 109
Mechanized assay of serum cholinesterase by specific colorimetric detection of released acid
Takeuchi T., Kabasawa Y., Horikawa R., Tanimura T. 117
A coupled assay detecting defects in fibroblast isoleucine degradation distal to enoyl-CoA hydratase: Application to 3-oxothiolase deficiency
Gibson K.M., Lee C.F., Kamali V., Sovik O. 127
Separation of plasma lipoproteins with a tabletop ultracentrifuge
Sykes E., Meany M., Schulz V., Kessel D. 137
Letters to the Editor
More on monoamine oxidase activity in whole blood
Konings C.H., Kuiper M.A., Wolters E.C. 145
Solving one of the problems of in vitro cytokine production and endotoxin measurement in blood
Redl H., Bahrami S., Leichtfried G., Schlag G. 149
Intestinal permeability: Update on the enzymatic assay of mannitol
Lunn P.G., Northrop-Clewes C.A. 151
The precipitation of histone-proteins with trichloroacetic acid
Lakatos A., Jobst K. 153
Erratum: Purification and structural study of two albumin variants in an Irish population (Clinica Chimica Acta, 204 (1991) 179–188)
Sakamoto Y., Davis E., Madison J., Watkins S., McLaughlin H., Leahy D.T., Putnam F.W. 155

Vol. 205/3, 1992

Cholinesterases exhibiting aryl acylamidase activity in human amniotic fluid
Jayanthi L.D., Balasubramanian N., Balasubramanian A.S. 157
Activity of cholesterol in human gallbladder bile in relation to nucleation of cholesterol monohydrate crystals
Noshiro H., Chijiwa K., Hirota I. 167
Increased serum concentration of type IV collagen peptide and type III collagen peptide in hyperthyroidism
Inui T., Ochi Y., Chen W., Nakajima Y., Kajita Y. 181
The determination of asialoglycoforms of serum glycoproteins by lectin blotting with Ricinus communis agglutinin
Lundy F.T., Wisdom G.B. 187
Clearance of purified human liver γ-glutamyltransferase after intravenous injection in the rat
Huseby N.-E., Kindberg G.M., Grostad M., Berg T. 197
Increase in plasma total and lipoprotein cholesterol during incubation of whole blood samples at 37°C — Influence of LCAT inhibitors
Ruhling K., Lang A., Holtz H., Winkler L., Schlag B., Till U. 205
A competitive ELISA for lipoprotein(a)
Yeo K.H.J., Walmsley T.A., Owen M.C., George P.M. 213
Serum phospholipase A_2 and lysolecithin changes following myocardial infarction
Langton S.R., Jarnicki A. 223
Direct gas chromatographic assay of urinary medium-chain fatty acylcarnitines by their thermal decomposition
Farquharson J., Jamieson E.C., Muir J., Cockburn F., Logan R.W. 233
Letters to the Editor
The influence of antioxidants on the oxidative stress of red blood cells
Varga S.I., Novak Z., Pataki L., Patocskai M., Matkovics B. 241
Competitive enzyme inhibition immunoassay of apolipoprotein B based on monoclonal antibody
Rojas A., Zayas M., Falero G., Rodriguez I. 245
Use of homocysteine to stabilise ascorbic acid, or to reduced dehydroascorbic acid, during HPLC separation of large volumes of tissue extracts
Bates C.J. 249

Quantities and units for electrophoresis in the clinical laboratory
Ferard G. S17
Quantities and units for centrifugation in the clinical laboratory
Lauritzen M. S5

Vol. 206/1–2, 1992

International Week of Laboratory Science — Rome, Italy, November 12–17, 1990: Introduction
Arvan D.A., Burlina A. 1
Tumor cell heterogeneity: An overview
Arvan D.A. 3
Phenotypic heterogeneity in acute leukemia
Ryan D.H. 9
Drug-transporter proteins in clinical multidrug resistance
Scheper R.J., Broxterman H.J., Scheffer G.L., Meijer C.J.L.M., Pinedo H.M. 25
Evaluation of tumor cell heterogeneity based on response to chemotherapy
Asselin B.L. 33
Enzymes as agents for the treatment of disease
Goldberg D.M. 45
Enzymes as prognostic markers and therapeutic indicators in patients with cancer
Schwartz M.K. 77

Application of clinical laboratory measurements to issues of environmental health
Rej R., Silkworth J.B., DeCaprio A.P. 83
Glutathione, protein sulfhydryls and cysteine proteases in gastric mucosal injury and protection
Szabo S., Nagy L., Plebani M. 95
Measurement of tryptase in endoscopic gastroduodenal biopsies: Distribution and relationship with ulcer disease
Plebani M., Di Mario F., Battistel M., Basso D., Mantovani G., Giacomini A., Burlina A. 107
Magnetic resonance spectroscopy of the brain
Prichard J.W. 115
Carbon-13 magnetic resonance in biology and medicine
Seelig J., Burlina A.P. 125
Quantitative analysis of rat synaptosomes and cerebrum using high-resolution ^1H magnetic resonance spectroscopy
Petroff O.A.C., Burlina A.P., Black J., Prichard J.W. 137
ATP-stimulated glutamate-dependent calcium uptake by rat synaptosomes
Galzigna L., Bianchi M., Battistin T., Scarpa M., Rigo A. 147

Vol. 206/3, 1992

A sequential and simple determination of zinc, copper and aluminium in blood samples by inductively coupled plasma atomic emission spectrometry
Chappuis P., Poupon J., Rousselet F. 155
Characteristics of the interactions between acid glycosaminoglycans and protein in normal human plasma as revealed by the behaviour of the protein-polysaccharide complexes in ultrafiltration and chromatographic procedures
Calatroni A., Vinci R., Ferlazzo A.M. 167
Relationship of urinary pseudouridine and 1-methyladenosine to activity of leukemia and lymphoma
Itoh K., Konno T., Sasaki T., Ishiwata S., Ishida N., Misugaki M. 181
Creatine kinase MB isoforms for early diagnosis and monitoring of acute myocardial infarction
Kanemitsu F., Okigaki T. 191
Organic acids in post-mortem cerebrospinal fluid
Coude M., Kamoun P. 201
Effects of storage, iron and time of day on indices of lipid peroxidation in plasma from healthy volunteers
Duthie G.G., Morrice P.C., Ventresca P.G., McLay J.S. 207
Determination of fetal hemoglobin by cation exchange liquid chromatography
Koskinen L., Sillanaukee P. 215
Serum α_1-antitrypsin in patients with hepatocellular carcinoma
Lee H.B., Yoo O.J., Ham J.S., Lee M.H. 225
β-N-Acetylhexosaminidase in the urine, kidney and serum of bromobenzene-intoxicated mice
Tassi C., Beccari T., Casini A., Orlacchio A. 231
Letters to the Editor
Use of toxi-lab thin layer chromatography in plasma very long chain fatty acid analysis
Smith N.B., Brooks K.E. 241
Reference intervals for α_1-microglobulin in urine
Jung K., Pergande M., Schreiber G., Thierfelder W. 245
Improved isolation of mononuclear cells from stored blood
Ford T.C., Rickwood D. 249

Vol. 207/1–2, 1992

Assay of γ-glutamyltransferase with amino acid dehydrogenases from Bacillus stearothermophilus as auxiliary enzymes
Kondo H., Hashimoto M., Nagata K., Tomita K., Tsubota H. 1

Enzyme-linked immunosorbent assay for serum pseudocholinesterase
Itoh Y., Fukamachi I., Hashimoto H., Hachisu T., Kawai T. 11
A specific method for determination of peroxisomal β-oxidation activity in cultured human skin fibroblasts using a specific substrate, C_9: A possible application for screening of peroxisomal disorders
Suzuki H., Yamada J., Watanabe T., Suga T. 19
Serum analysis for potassium ions using a fibre optic sensor
Pereiro Garcia R., Alava Moreno F., Diaz Garcia M.E., Sanz-Medel A., Narayanaswamy R. 31
Aluminum-binding serum proteins: Desferrioxamine alters serum aluminum speciation
Favarato M., Mizzen C.A., Sutherland M.K., Krishnan B., Kruck T.P.A., Crapper McLachlan D.R. 41
Insulin binding to erythrocytes of nonpregnant women: A reevaluation, underlining the importance of body weights even in nonobese subjects
Desoye G., Schmon B., Gmoser G., Friedl H., Urdl W., Weiss P.A.M. 57
An indirect sandwich ELISA for LP(a) in serum and dried blood spots
Wang X.L., Wilcken D.E.L., Dudman N.P.B. 73
Distribution of SP_1, immunoreactivity among different plasma proteins: Real molecular heterogeneity or adsorption of SP_1-beta to other plasma proteins?
Pala A., Di Ruzza A., Rossetto G., Moro M., Strom R., Padula F., Cosmi E.V. 87
Influence of triglycerides and urate on methods for determination of fructosamine
Kallner A. 99
Comparison of soluble aminopeptidases in human cerebral cortex, skeletal muscle and kidney tissues
Mantle D. 107
The effect of blood sample aging and food consumption on plasma total homocysteine levels
Ubbink J.B., Hayward Vermaak W.J., Van der Merwe A., Becker P.J. 119
A case of Hashimoto's thyroiditis with antibodies to thyroid hormone analogues
Sakata S., Maekawa H., Ogawa T., Sarui H., Matsuda M., Endo J., Wada S., Nagai K. 129
Selenium, glutathione peroxidase (GSH-Px) and lipid peroxidation products before and after selenium supplementation
Wilke B.C., Vidailhet M., Favier A., Guillemin C., Ducros V., Arnaud J., Richard M.J. 137
Critical evaluation of the serum kappa/lambda light-chain ratio in the detection of M proteins
Laine S.T., Soppi E.T., Morsky P.J. 143
Letter to the Editor
3-Methylglutaconic aciduria in two adults
Kuhara T., Matsumoto I., Saiki K., Takabayashi H., Kuwabara S. 151

Vol. 207/3, 1992

Increased prostaglandin PGE_2 and 6-keto-PGF(1α) immunoactivity in incubates of gastric mucosa and forestomach of nephrectomized rats
Weiler H., Esser D., Weiler C., Gerok W. 161
Routine use of hair root or buccal swab specimens for PCR analysis: Advantages over using blood
Thomson D.M., Brown N.N., Clague A.E. 169
Determination of enterokinase activity by measuring the disappearance of trypsinogen
Carchon H.A., Jansen E., Eggermont E. 175
The measurement of carnitine and acyl-carnitines: Application to the investigation of patients with suspected inherited disorders of mitochondrial fatty acid oxidation
Bhuiyan A.K.M.J., Jackson S., Turnbull D.M., Aynsley-Green A., Leonard J.V., Bartlett K. 185
A flow-through system for the determination of salicylate in blood
Bertocchi P., D'Ottavio D., Evangelisti M.E., Mascini M., Palleschi G. 205
A simple, sensitive technique for classification of apolipoprotein(a) isoforms by sodium dodecyl sulphate-polyacrylamide gel electrophoresis
Farrer M., Game F.L., Adams P.C., Laker M.F., Alberti K.G.M.M. 215
Purification anhd characterisation of antigenic gliadins in coeliac disease
Sjostrom H., Friis S.U., Noren O., Anthonsen D. 227

Human urinary protein 1: Evidence for identity with the Clara cell protein and occurrence in respiratory tract and urogenital secretions
Bernard A., Roels H., Lauwerys R., Witters R., Gielens C., Soumillion A., Van Damme J., De Ley M. 239
Need for both angiotensin converting enzyme and angiotensinase specific inhibition for a reliable assessment of angiotensin II levels in human plasma
Boddi M., Coppo M., Rostagno C., Gensini G.F., Neri Serneri G.G. 251
Letters to the Editor
Within-subject biological variation of pituitary-ovarian axis hormones and desirable imprecision
Fuentes-Arderiu X., Albert P. 257
Expression of CuZn-superoxide dismutase and glutathione peroxidase in erythrocytes from diabetic and non-diabetic subjects
Strange R.C., Jones P., Bicknell J., Scarpello J. 261

Structure and genetic engineering of antigens and antibodies: Applications in immunoassays
Vihko P., Wagener C. S5
Testing urine for drugs
Catlin D., Cowan D., Donike M., Fraisse D., Oftebro H., Rendic S. S13

Vol. 208/1–2, 1992

Possible role for chondroitin sulfate in urolithiasis: In vivo studies in an experimental model
Michelacci Y.M., Boim M.A., Bergamaschi C.T., Rovigatti R.M., Schor N. 1
Serum immunoreactive β-glucuronidase determined by an enzyme-linked immunosorbent assay in patients with hepatic diseases
Ohta H., Ono M., Sekiya C., Namiki M. 9
Procedure for development of an enzyme-linked immunosorbent assay. Development of an assay for human apolipoprotein A-I
Gaver A., Weedy N.K., Maldonado B.R., Huang S., Wong L. 23
^{13}C and ^{31}P NMR studies of glucose and 2-deoxyglucose metabolism in normal and enzyme-deficient human erythrocytes
Ferretti A., Bozzi A., Di Vito M., Podo F., Strom R. 39
Optimized determination of malondialdehyde in plasma lipid extracts using 1,3-diethyl-2-thiobarbituric acid: Influence of detection method and relations with lipids and fatty acids in plasma from healthy adults
Hoving E.B., Laing C., Rutgers H.M., Teggeler M., Van Doormaal J.J., Muskiet F.A.J. 63
The glycosaminoglycan composition of the lung with acute and chronic pathology and in senescence
Grundboeck-Jusco J., Binette J.P., Kimura A., Talarico L., Kaelin R., Schmid K. 77
Determination of (S)- and (R)-2-oxo-3-methylvaleric acid in plasma of patients with maple syrup urine disease
Wendel U., Even G., Langenbeck U., Schadewaldt P., Hummel W. 85
An immunoreactive xanthine oxidase protein-possessing xanthinuria and her family
Yamamoto T., Moriwaki Y., Suda M., Takahashi S., Hada T., Nanahoshi M., Agbedana E.O., Higashino K. 93
A receptor assay for serum low density lipoprotein-apolipoprotein B-100: Preliminary report
Pham Q., Soldin S.J., Rifai N. 99
Amniotic fluid granulocyte colony-stimulating factor in preterm and term labor
Saito S., Kato Y., Ishihara Y., Ichijo M. 105
Abnormal lysosomal cathepsin activities in leukocytes and cultured skin fibroblasts in late infantile, but not in juvenile neuronal ceroid-lipofuscinosis (Batten disease)
Bennett M.J., Chern L., Carpenter K.H., Sladky J.T. 111
Liquid chromatography determination of α-tocopherol in erythrocytes
Sierra C., Pastor M.C., De Ramon M. 119

Letters to the Editor

Effect of anaesthesia and acute intestinal ischemia on serum β-N-acetylhexosaminidase activity in rabbit as biological model
Risueno J.C., Gomez-Alonso A., Garcia-Criado F.J., Garcia-Sastre A., Corral J., Cabezas J.A., Villar E. 127

Fetal and amniotic α_1-acid glycoprotein
Johnson A.M. 133

A method for quantitative collection of stool and urine for nitrogen balance studies in infants
Kandil H.M., Switzer B.R., Heizer W.D. 135

Rapid and sensitive colloidal silver staining on cellulose acetate membranes
Kovarik A., Ondrkalova M., Prachar J., Stofko J. 137

The effect of reducing and non-reducing sugars on the bicinchoninic acid reaction for protein determination
Milton J.D., Mullen P.J. 141

Erratum: The influence of antioxidants on the oxidative stress of red blood cells (Clinica Chimica Acta (1992) 205 (241–244))
Varga S.I., Novak Z., Pataki L., Patocskai M., Matkovics B. 147

Vol. 208/3, 1992

N-glycosylation of serum proteins in disease and its investigation using lectins
Turner G.A. 149

Prenatal diagnosis of Niemann-Pick disease type C
De Winter J.M., Janse H.C., Van Diggelen O.P., Loss F.J., Beemer F.A., Kleijer W.J. 173

High-performance liquid chromatographic microassay for L-alanine:glyoxylate aminotransferase activity in human liver
Petrarulo M., Pellegrino S., Marangella M., Cosseddu D., Linari F. 183

Serum triglyceride and insulin levels are associated with erythrocyte sodium-lithium countertransport activity in normoglycaemic individuals
Winocour P.H., Thomas T.H., Brown L., Laker M.F., Wilkinson R., Alberti K.G.M.M. 193

Liver transplantation modifies serum cholestanol, cholesterol precursor and plant sterol levels
Nikkila K., Hockerstedt K., Miettinen T.A. 205

A simple electrophoretic method for separating elevated liver and bone alkaline phosphatase isoenzymes in plasma after neuraminidase treatment
Chamberlain B.R., Buttery J.E., Pannall P.R. 219

2,3-Butanediol: A potential interfering substance in the assay of ethylene glycol by an enzymatic method
Nilsson L., Jones A.W. 225

Letters to the Editor

Oligosaccharides in (galacto)sialidosis urine
Van Pelt J. 231

Enzyme immunoassay for serum cortisol
Candito M., Krebs B., Chambon P. 233

Effect of long-term storage on human plasma osteocalcin level
Tarallo P., Balland M., Henny J. 237

Vol. 209/1–2, 1992

Decrease of pyruvate dehydrogenase phosphatase activity in patients with congenital lactic acidemia
Ito M., Kobashi H., Naito E., Saijo T., Takeda E., Mahbubul Huq A.H.M., Kuroda Y. 1

Clinical test of renal guanidinoacetic acid metabolism by oral citrulline and creatine loading
Kadono K., Yamaguchi T., Tetsutani T., Yasunaga K. 9

Plasmodium falciparum induced perturbations of the erythrocyte antioxidant system
Dubey M.K., Dubey M.L., Ganguly N.K., Mahajan R.C. 19
Characterization of the carbohydrate moiety of human γ-glutamyltransferase using lectin-blotting and glycosidase treatment
Evjen G., Huseby N.-E. 27
Use of an azo derivative immobilized on an anionic resin for the determination of aluminium in water and dialysis fluids
Hernandez Torres O., Jimenez Moreno F., Jimenez Abizanda A., Arias Leon J.J. 35
Investigation of urea cycle enzyme disorders by ^{1}H-NMR spectroscopy
Burns S.P., Woolf D.A., Leonard J.V., Iles R.A. 47
Diagnosis of medium chain acyl CoA dehydrogenase deficiency by measurement of cis-4-decenoic acid in dried blood spots
Heales S.J.R., Leonard J.V. 61
Increased urinary excretion of coproporphyrin isomers after transcatheter arterial embolization of hepatocellular carcinoma
Horie Y., Yamada S., Kawasaki H. 67
Diagnostic value of adenosine deaminase and lysozyme in tuberculous pleurisy
San Jose E., Valdes L., Sarandeses A., Alvarez D., Chomon B. 73
Evaluation of solubilizing agents for 25-hydroxyvitamin D_3 immunoassays
Kobayashi N., Ueda K., Shimada K. 83
Raised monophosphatase activity in schizophrenic patients
Zilberman-Kaufman M., Agam G., Moscowits L., Livne A.A., Belmaker R.H. 89
Increased superoxide dismutase activity in erythrocytes of children with pulmonary hypertension
Soave M.C., Moulsma M., Chevalier P., Pillot R., Guidollet J. 95
Letters to the Editor
Screening of glycerol kinase deficiency in patients affected by Duchenne and Becker muscular dystrophy
Schreiber R., Passos-Bueno M.R., Rapaport D., De Cassia Pavanello R., Vainzof M., Fernandes M.E.S., Zatz M. 103
A colorometric determination of 5-hydroxyindole acetic acid in urine
Mukerjee H., Pincus M.R. 105
9th European Congress of Clinical Chemistry
Enzymes as reporters of pathophysiological events: Introduction
Moss D.W., Naskalski J. 107
Enzyme induction by drugs and toxins
Batt A.-M., Siest G., Magdalou J., Galteau M.-M. 109
Alkaline phosphatase as a reporter of cancerous transformation
Millan J.L. 123
Changes in enzyme expression related to differentiation and regulatory factors: The acid phosphatase of osteoclasts and other macrophages
Moss D.W. 131

Vol. 209/3, 1992

Alterations of purine metabolism in mononuclear cell populations of first degree relatives of insulin-dependent diabetic individuals with disturbed glucose tolerance
Schafer J.H., Jira T.B., Kohler E., Michaelis D., Rjasanowski I.M. 141
Melatonin and 6-sulfatoxymelatonin circadian rhythms in serum and urine of primary prostate cancer patients: Evidence for reduced pineal activity and relevance of urinary determinations
Bartsch C., Bartsch H., Schmidt A., Ilg S., Bichler K.-H., Fluchter S.-H. 153
Lipid and apolipoprotein changes after orthotopic liver transplantation for end-stage liver diseases
Malmendier C.L., Lontie J.-F., Mathe D., Adam R., Bismuth H. 169
Relationships between androgen and estrogen sulfates in breast cyst fluid
Nahoul K., Kottler M.-L. 179
Increased membrane activity of glyceraldehyde 3-phosphate dehydrogenase in erythrocytes of patients with homozygous sickle cell anaemia
Mohamed A.O., Ronquist G., Al Bayoumi R. 189

An ELISA method to measure human myoglobin in urine
Hodson A.W., Skillen A.W., Argent N.B. 197
Comparison of three techniques for the determination of protein content in human tears
Meijer F., Van Haeringen N.J. 209
Letter to the Editor
Consequences of applying parametric methods to non-gaussian distributions in the validation of analytical results
Machuca I., Cortes J.D., Fuentes-Arderiu X. 215

Diagnostic applications of repetitive DNA sequences
Epplen J.T. S5
Genes in the diagnosis of malignant disease
Wagener C. S15

Vol. 210/1–2, 1992

Finally! Automated microanalysis of ammonia in arterial blood
Conn H.O. 1
Arginase, a new marker of mammary carcinoma
Straus B., Cepelak I., Festa G. 5
Cellulose acetate electrophoretic determination of bone alkaline phosphatase activity in healthy subjects and diabetic patients with and without osteopenia
Nishikawa Y., Kanda T., Yoshihara H., Fukumoto K., Uematsu I. 13
Characterization of dipeptidyl peptidase IV (CD26) from human lymphocytes
De Meester I., Vanhoof G., Hendriks D., Demuth H.-U., Yaron A., Scharpe S. 23
Differences in the enzymatic nature and the sugar-chain structure of γ-glutamyl transferase between normal and carcinomatous human kidney and prostate
Arai K., Yoshida K.-I., Komoda T., Kobayashi N., Sakagishi Y. 35
Comparison of urinary excretion of four lysosomal hydrolases in healthy elderly and young adults
VanderJagt D.J., Steinberg B.R., Glew R.H. 47
Identification of urinary metabolites of (±)-2-(p-isobutylphenyl)propionic acid (Ibuprofen) by routine organic acid screening
Bennett M.J., Sherwood W.G., Bhala A., Hale D.E. 55
Intestinal type alkaline phosphatase hyperphosphatasemia associated with liver cirrhosis
Saheki S., Ochi M., Horiuchi T., Sakagishi Y., Fujimori-Arai Y., Koyama I., Komoda T., Takeuchi N. 63
Purification of electron transfer flavoprotein from pig liver mitochondria and its application to the diagnosis of deficiencies of acyl-CoA dehydrogenases in human fibroblasts
Bertrand C., Dumoulin R., Divry P., Mathieu M., Vianey-Saban C. 75
Interaction of oxidized low density lipoproteins with both apo B,E and scavenger receptors. A model for its production in vitro
Vilella E., Joven J., Bargallo T., Turner P.R., Masana L. 93
Concentration of serum laminin and type IV collagen in liver diseases assayed by a sandwich enzyme-immunoassay using monoclonal antibodies
Yokoya Y., Iwata K., Muragaki Y., Shiota C., Morimoto Y., Inoue M., Itoh H., Nishioka S., Ooshima A. 109
A hyaluronan-loading test applied to patients with liver and joint diseases
Lindqvist U., Groth T., Loof L., Hellsing K. 119
Pregnancy-associated α_2-glycoprotein in children with acute lymphocytic leukemia, Hodgkin's disease and non-Hodgkin's lymphomas
Zalazar F.E., Chiabrando G.A., De Aldao N.A., Ojeda F., Vides M.A., Aldao M.A.J. 133
PAF acetylhydrolase activities in human systemic lupus erythematosus and lupus-prone mice
Matsuzaki M., Ishiguro Y., Tetsumoto T., Tsukada Y., Kayahara H., Tadasa K. 139
CA 19-9 in non-neoplastic liver diseases. A clinical and laboratory study
Collazos J., Genolla J., Ruibal A. 145

Letter to the Editor
A rapid method for blood ammonia determination using the new blood ammonia checker (BAC) II
Huizenga J.R., Tangerman A., Gips C.H. 153

Vol. 210/3, 1992

The role of citrate complexes in preventing urolithiasis
Ashby R.A., Sleet R.J. 157
Immunochemiluminometric assays (ICMA) specific for growth hormone releasing hormone 1-44 NH_2 and 1-40 OH
Page M.D., Edwards C.A., Brown R., Weeks I., Woodhead J.S., Scanlon M.F. 167
The inter-relationships between albuminuria, plasma albumin concentrtion and indices of glycaemic control in non-insulin-dependent diabetes mellitus
Chan J.C.N., Yeung V.T.F., Cheung C.K., Swaminathan R., Cockram C.S. 179
IL6 and acute phase plasma proteins in peritoneal fluid of women with endometriosis
Boutten A., Dehoux M., Edelman P., Seta N., Menard A., Madelenat P., Durand G. 187
Development and validation of a robust specific enzyme mediated assay for phenylalanine in serum
Campbell R.S., Brearley G.M., Varsani H., Morris H.C., Milligan T.P., Hall S.K., Hammond P.M., Price C.P. 197
Evidence that the presence of arginine can lead to overestimation of glucagon levels measured by radioimmunoassay
Lacey R.J., Scarpello J.H.B., Morgan N.G. 211
Evaluation of rat intestinal lactase in vivo with 4-galactosylxylose
Aragon J.J., Fernandez-Mayoralas A., Jimenez-Barbero J., Martin-Lomas M., Rivera-Sagredo A., Villanueva D. 221
Letters to the Editor
Impaired synthesis of taurine in a patient with tyrosinemia type I during the oral L-methionine loading test
Valik D., Klusacek D., Pintera J. 227
Lipoprotein(a) and the acute phase response
Craig W.Y., Ledue T.B. 231
Activity of cathepsin H,B and metalloproteinase in the serum of patients with acute myocardial infarction
Laszlo A., Sohar I., Sagi I., Kovacs J., Kovacs A. 233
Disease-coupled specificity in the glycan structures of orosomucoid
Treuheit M.J., Halsall H.B. 237

Vol. 211/1–2, 1992

Diagnosis of HIV infection
Schochetman G. 1
Differential isoform profiles of α_2-macroglobulin from plasma of patients with chronic-progressive or relapsing-remitting multiple sclerosis
Back S.A., Alhadeff J.A. 27
High levels of nervous system-specific proteins in cerebrospinal fluid in patients with early stage Creutzfeldt-Jakob disease
Jimi T., Wakayama Y., Shibuya S., Nakata H., Tomaru T., Takahashi Y., Kosaka K., Asano T., Kato K. 37
Renin assay using a fluorogenic substrate and high performance liquid chromatography
Nakamura-Imajo N., Satomura S., Matsuura S., Murakami K. 47
A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 3 (stromelysin-1) using monoclonal antibodies
Obata K., Iwata K., Okada Y., Kohrin Y., Ohuchi E., Yoshida S., Shinmei M., Hayakawa T. 59

Isolation of acylcarnitines from urine: A comparison of methods and application to long-chain acyl-CoA dehydrogenase deficiency
Morrow R.J., Rose M.E. 73
Glycation of human tissue and serum creatine kinase
Langlois M.R., Delanghe J.R., De Buyzere M.L., Leroux-Roels G.G. 83
Serum apolipoproteins in heterozygous familial hypercholesterolemia
Kajinami K., Mabuchi H., Koizumi J., Takeda R. 93
Lipid peroxides in blood from patients with atherosclerosis of coronary and peripheral arteries
Plachta H., Bartnikowska E., Obara A. 101
Photographic chemiluminescent ELISA for detection of anti-human T-cell leukemia virus type-I antibodies by using synthetic peptides as antigens
Kuroda N., Nakashima K., Akiyama S., Shiraki H., Maeda Y. 113
Letters to the Editor
Transferrin, cholesterol and aluminium in Alzheimer's disease
Corrigan F.M. 121
Correlations between the serum pancreatic enzyme activities and lipid peroxidation in acute pancreatitis
Matkovics B. 125
Effect of long-term storage on human plasma cholinesterase activity
Henny J. 129

Vol. 211/3, 1992

Bone metal content in patients with chronic renal failure
Navarro J.A., Granadillo V.A., Salgado O., Rodriguez-Iturbe B., Garcia R., Delling G., Romero R.A. 133
High-performance liquid chromatographic assay for L-glyceric acid in body fluids. Application in primary hyperoxaluria type 2
Petrarulo M., Marangella M., Cosseddu D., Linari F. 143
Red blood cells and platelet membrane fatty acids in non-dialyzed and dialyzed uremics
Girelli D., Azzini M., Olivieri O., Guarini P., Trevisan M.T., Lupo A., Bernich P., Panzetta G., Corrocher R. 155
Distribution of lipoprotein species (LpA-I, LpA-I:A-II) in serum and HDL subfractions of untrained and trained normolipemic men
Frey I., Kamps H., Baumstark M.W., Berg A., Keul J. 167
A screening method for cystine and homocystine in urine
Thuy L.P., Nyham W.L. 175
Increase in alkaline phosphodiesterase I activity of tumor-derived cultured fibroblasts from neurofibromatosis patients
Maruyama E., Takashima S., Arima M. 181
Letters to the Editor
Selenium in Graves' disease
Reglinski J., Smith W.E., Wilson R., Halls D.J., McKillop J.H., Thomson J.A. 189
Standardization of Lp(a) assays
Kostner G.M. 191

Quantities and units for metabolic processes as a function of time. Recommendations 1992
Ferard G. S5
Flow cytometry in the clinical laboratory. Principles, applications and problems
Ruiz-Arguelles A. S13

Vol. 212/1–2, 1992

Measurement of glycated albumin by the nitroblue tetrazolium colorimetric method
Mashiba S., Uchida K., Okuda S., Tomita S. 3

A new quantitative assay for glycosaminoglycans
Thuy L.P., Nyhan W.L. 17
Adaptation of transferrin protein and glycan synthesis
De Jong G., Van Noort W.L., Feelders R.A., De Jeu-Jaspars C.M.H., Van Eijk H.G. 27
Effects of methylprednisolone and uremia on renal and intestinal calbindin-D in the rat
Egfjord M., Staun M., Olgaard K. 47
Determination of free carnitine and 'total' carnitine in human urine: Derivatization with 4′-bromophenacyl trifluoromethanesulfonate and high performance liquid chromatography
Minkler P.E., Hoppel C.L. 55
Biliary lipid composition and metastability of biliary cholesterol in patients with acalculous cholesterolosis
Chijiiwa K., Nakano K., Makino I., Noshiro H., Yamasaki T. 65
Heterogeneity of antigens related to 7S domain of type IV collagen in serum: Application to patients with chronic liver disease
Yamada S., Kishimoto Y., Suou T., Kawasaki H., Horie Y. 73
An acquired acute hepatic porphyria: A novel type of δ-aminolevulinate dehydratase inhibition
Akagi R., Prchal J.T., Eberhart C.E., Sassa S. 79
Letter to the Editor
Seasonal variation in serum total concentrations of cholesterol and protein in elderly subjects
Fujimara A., Ohashi K., Ebihara A., Hirose T. 85

Vol. 212/3, 1992

Quantitative analysis of extracellular-superoxide dismutase in serum and urine by ELISA with monoclonal antibody
Adachi T., Ohta H., Yamada H., Futenma A., Kato K., Hirano K. 89
The BT-PABA/PAS test in tropical diabetes
Gagee P., Pemberton P., Lobley R., Chaloner C., Snehalatha C., Mohan V., Braganza J.M. 103
Detection of erythrocyte membrane components in hemoglobin-based blood substitutes
Biessels P.T., Berbers G.A., Broeders G.C., Landsvater R., Huisman H.G., Bleeker W.K., Bakker J.C. 113
Serum neurone-specific enolase levels in patients with neuroendocrine and carcinoid tumours
Cunningham R.T., Johnston C.F., Irvine G.B., Buchanan K.D. 123
A colorimetric microassay for glycated collagen based on the thiobarbituric acid method
Ahmed N., Malik N.S., Furth A.J. 133
Development of a method for the purification of human trypsin by single step affinity chromatography suitable for human isotope incorporation studies
Ogden J.M., O'Keefe S.J.D., Marks I.N. 141
Letters to the Editor
The variation of serum lipoprotein (a) during surgical operations
Magnani B., Brega Massone P.P., Meriggi F., Di Jeso F. 149
Antibodies to thyroid hormone analogues
John R. 153
Kinetic fluorometric micromethod for screening of newborns for galactosemia — A way to improve assay specificity?
Tuuminen T., Akerlind G. 155

Vol. 213/1–3, 1992

Platelet hyperreactivity in hyperlipidaemia with special reference to platelet lipids and fatty acid composition
Chetty N., Naran N.H. 1
Measurement of ascorbic acid in platelets and its relationship to polymorphonuclear leukocyte levels
McCulloch R.K., Vandongen R. 15

Polyclonal immunoglobulin M: Location of glycation sites
Menini T., Gugliucci A., Stahl A.J.C. 23
Characterization of the gliadin-derived peptides which are biologically active in coeliac disease
Cornell H., Wieser H., Belitz H.-D. 37
Copper metabolism — A factor in gestational diabetes?
Loven A., Romem Y., Pelly I.Z., Holcberg G., Agam G. 51
Differences in erythrocyte sodium transport between human plasma and artificial medium: The role and character of sodium efflux and influx stimulating plasma factors
Tepper T., Jilderda J.F., Huisman R.M., Van der Hem G.K., De Zeeuw D. 61
Protection of vascular basement membrane and microcirculation from elastase-induced damage with a fluorinated β-lactam derivative
Maillard J.-L., Favreau C., Vergely I., Reboud-Ravaux M., Joyeau R., Kobaiter R., Wakselman M. 75
A pulmonary origin of amniotic fluid thermolabile alkaline phosphatase near term
Ragoucy-Sengler C., Bredent-Bangou J., Pileire B. 87
Neonatal screening for familial hypercholesterolaemia by immunoturbidimetric assay of apolipoprotein B in dried blood spots
Bangert S.K., Eldridge P.H., Peters T.J. 95
A two-phase colorimetric method for the measurement of pepsin activity
Heinova D., Blahovec J., Kasafirek E., Bartik M., Rosival I. 103
Letter to the Editor
Improved immunoassay for perphenazine
Uebel R., Van Jaarsveld P., Wium C., Hawtrey A.O. 111

Vol. 214/1, 1993

Establishment of assay kits for the determination of microheterogeneities of alpha-fetoprotein using lectin-affinity electrophoresis
Shimizu K., Taniichi T., Satomura S., Matsuura S., Taga H., Taketa K. 3
Selective and sensitive determination of urinary total proline and hydroxyproline by gas chromatography with flame photometric detection
Kataoka H., Nabeshima N., Nagao K., Makita M. 13
A rapid GLC method for the direct analysis of plasma medium chain fatty acids
Mingrone G., De Gaetano A., Greco A.V., Capristo E., Raguso C., Tataranni P.A., Castagneto M. 21
A clinical evaluation of a direct radioimmunoassay of testosterone
Holownia P., Conway G.S., Shrivastava U., Round J., Honour J.W. 31
Lipoprotein(a) in diabetes mellitus
Ritter M.M., Loscar M., Richter W.O., Schwandt P. 45
Use of seronegative samples for the calibration and reporting of serological enzyme immunoassay results
Sorensen K., Viljanen M.K. 55
Enzymes of lysosomal origin in plasma of twin neonates
Goi G., Burlina A.B., Bairati C., Bordugo A., Zanardo V., Zacchello F., Tettamanti G., Lombardo A. 61
Measurement of insulin-like growth factor binding protein-1 in cervical/vaginal secretions: Comparison with the ROM-check Membrane Immunoassay in the diagnosis of ruptured fetal membranes
Rutanen E.-M., Pekonen F., Karkkainen T. 73
Characterization of γ-glutamyltranspeptidase from human hepatocellular carcinoma, compared with enzymes from normal liver and cirrhotic liver
Ohta H., Sawabu N., Kawakami H., Watanabe H., Ozaki K., Toya D., Hattori N. 83
Preparation of microsome fractions of human myometrium. A comparison of three methods
Lamminen S., Rantala I., Tuimala R. 93
Red blood cell adenosine deaminase activity in multiple sclerosis
Kopff M., Zakrzewska I., Czernicki J., Klem J., Strzelczyk M. 97

Letters to the Editor

A methodological approach to superoxide dismutase (SOD) activity assay based on inhibition of nitroblue tetrazolium (NBT) reduction
Durak I., Yurtarslanl Z., Canbolat O., Akyol O. 103

Activity of phospholipase A in serum from patients with cystic fibrosis
Laszlo A., Nemeth M., Gyurkovits K. 105

Basal concentration of the isoenzyme BB of the glycogen phosphorylase b in human blood
Rabitzsch G., Noll F., Hofmann U., Krause E.-G., Armbruster F.P. 109

Impact of circadian rhythms on the interpretation of stat measurements
Fuentes-Arderiu X., Rivera-Coll A. 113

The role of lipids in the increased mortality following bereavement
Coulter F., Armstrong K.M.D., Armstrong E.M., Davies A., Merry E., Skinner E.R., Corrigan F.M. 119

Vol. 214/2, 1993

Circulating proalbumin associated with a second case of antitrypsin Pittsburgh
Brennan S.O., Sheat J.M., Aiach M. 123

Gravimetric determination of the water concentration in whole blood, plasma and erythrocytes and correlations with hematological and clinicochemical parameters
Lijnema T.H., Huizenga J.R., Jager J., Mackor A.J., Gips C.H. 129

Circulating tumour-associated mucin concentrations, determined by the CASA assay, in healthy women
McGuckin M.A., Ramm L.E., Joy G.J., Devine P.L., Ward B.G. 139

Determination of lipoprotein(a): Enzyme immunoassay and immunoradiometric assay compared
Marz W., Siekmeier R., Gross E., Gross W. 153

Variability of kidney tubular interstitial distortions in glomerulonephritis as measured by ^{1}H-NMR urinalysis
Lundina T.A., Knubovets T.L., Sedov K.R., Markova S.A., Sibeldin L.A. 165

Enzymatic determinations in acute rejection after liver transplantation: Preliminary report on necrosis index
Gozzo M.L., Avolio A., Forni F., Agnes S., Colacicco L., Barbaresi G., Castagneto M. 175

Antioxidant defenses and its modulation by iron in carrageenan-induced inflammation in rats
Muntane J., Puig-Parellada P., Fernandez Y., Mitjavila S., Mitjavila M.T. 185

High performance liquid chromatographic determination of free and conjugated bile acids in serum, liver biopsies, bile, gastric juice and feces by fluorescence labeling
Guldutuna S., You T., Kurts W., Leuschner U. 195

Trace element concentrations (Cu, Zn, Fe) in sera from patients with dilated cardiomyopathy
Oster O. 209

Cholinesterases in cerebrospinal fluid in patients with meningitis and hydrocephaly
Tornel P.L., Campoy F.J., Vidal C.J. 219

Plasma malondialdehyde in type 1 and type 2 diabetic patients
Gallou G., Ruelland A., Legras B., Maugendre D., Allannic H., Cloarec L. 227

Correction factors for single radial immunodiffusion and electroimmunodiffusion assays of IgA in secretory fluids
Meillet D., Kapel N., Favennec L., Raichvarg D., Savel J., Gobert J.G. 235

Letters to the Editor

Contamination of biological samples in metal determination
Villalba A. 243

Biological variability of twenty-five serum analytes during normal pregnancy
Alberty R. 245

Vol. 215/1, 1993

Relationship between alcohol consumption and the activity of GTP-binding regulatory proteins in human erythrocyte membranes
Nakamura J., Harada S., Tanaka N. 1
Further applications of blood gas analysis to gastric acidity determination
Niv Y., Abu-Avid S., Neumann G. 9
Magnesium, zinc and copper in plasma and blood cellular components in children with IDDM
Rohn R.D., Pleban P., Jenkins L.L. 21
Serum hyaluronic acid in healthy infants and children and its value as a marker of progressive hepatobiliary disease starting in infancy
Trivedi P., Cheeseman P., Mowat A.P. 29
Changes of lipid peroxide levels in blood and liver tissue of patients with obstructive jaundice
Tsai L.Y., Lee K.T., Tsai S.M., Lee S.C., Yu H.S. 41
Epidermal growth factor in human breast cancer, endometrial carcinoma and lung cancer. Its relationship to epidermal growth factor receptor, estradiol receptor and tumor TNM
Bolufer P., Lluch A., Molina R., Alberola V., Vazquez C., Padilla J., Garcia-Conde J., Llopis F., Guillem V. 51
Biocytin-specific 110-kDa biotinidase from human serum
Oizumi J., Hayakawa K. 63
Increased platelet activating factor in the tracheal aspirates from neonates with patent ductus arteriosus
Koyama N., Ogawa Y., Kamiya K., Eguchi H., Tanaka T., Takasaki J., Ohama Y., Itakura Y., Kaneko K. 73
Detection of heterozygotes for fructose-1,6-diphosphatase deficiency by measuring fructose-1,6-diphosphatase activity in monocytes cultured with calcitriol
Kikawa Y., Takano T., Nakai A., Shigematsu Y., Sudo M. 81
Frequency of the XbaI, EcoRI, PvuII and MspI polymorphisms of the apolipoprotein B gene in relation to hypercholesterolaemia in the general population
Series J.J., Gaffney D., Packard C.J., Shepherd J. 89
A new highly sensitive radioimmunoassay for human calcitonin useful for physiological studies
Schifter S. 99
Acceleration of hemoglobin glycation with aging
Nakashima K., Nishizaki O., Andoh Y. 111
Letters to the Editor
Ethanediol stabilised quality control serum for measurement of zinc
Swaminathan S., Seshadri M.S., Srikrishna G., Selvakumar R., Kanagasabapathy A.S. 119
Serum trehalase activity in patients with rheumatoid arthritis
Yoshida K., Mizukawa H., Haruki E. 123
Erratum: Distribution of lipoprotein species (LpA-I, LpA-I:AII) in serum and HDL subfractions of untrained and trained normolipemic men (Clinica Chimica Acta, 211 (1992) 167–173)
Frey I., Kamps H., Baumstark M.W., Berg A., Keul J. 125

Vol. 215/2, 1993

Molecular weight and substrate characteristics of human serum arylesterase following purification by immuno-affinity chromatography
Tanaka-Kawai H., Yomoda S. 127
A method for the assay of phosphatidylinositol-specific phospholipase D activity in serum
Raymond F.D., Fortunato G., Moss D.W. 139
Survey of methods for measuring human anti-mouse antibodies
Kricka L.J., Schmerfeld-Pruss D., Kaladas P., Colcher D., Scassellati G.A., Seccamani E., Courtenay-Luck N.S., Kosmas C., Beatty B.G., Beatty J.D., Chatenoud L., Zimmer A.M., Auerbach B. 153

Enzymatic diagnosis of steroid sulfatase deficiency by high performance liquid chromatography
Ohsaki M., Matsumoto T., Sakura N., Ueda K. 165
Glycan microheterogeneity for α_1-antitrypsin in serum and meconium from normal and cystic fibrosis patients by crossed immunoaffinoelectrophoresis with different lectins (Con A, LCA, WGA)
Duthel S., Revol A. 173
Phosphorylation of insulin-like growth factor-binding protein-1 increases in human amniotic fluid and decidua from early to late pregnancy
Koistinen R., Angervo M., Leinonen P., Hakala T., Seppala M. 189
Vitamin E attenuates induction of elastase-like activity by tumor necrosis factor-α, cholestan-3β,5α,6β-triol and linoleic acid in cultured endothelial cells
Toborek M., Hennig B. 201
Changes in immunoreactive manganese-superoxide dismutase concentration in human serum after 93 h strenuous physical exercise
Ohno H., Kayashima S., Nagata N., Yamashita H., Ookawara T., Taniguchi N. 213
Cigarette-smoking, birthweight, thiocyanate and fluorescent lipid-peroxidation products in maternal and cord plasma
Pre J., Colau J.C., Vassy R. 221
Measurement of histidase activity in human fingernail by reversed-phase HPLC
Fujitaka M., Eguchi T., Sakura N., Ueda K. 227
Letter to the Editor
Effects of ethanol on the assay of acyl-CoA:cholesterol acyltransferase (ACAT)
Smith J.L., Madden L.J., De Jersey J. 233

Vol. 216/1–2, 1993

Deregulated production of interleukin-4 (IL4) in autoimmune thyroid disease assayed with a new radioimmunoassay
Hirooka Y., Kayama M., Ohga S., Kimura M., Hasegawa M., Shin K., Nogimori T., Ishizuki Y., Mitsuma T. 1
Variable expression of leukocyte cytosolic broad-specificity β-glucosidase activity
Forsyth G.W., Romero K.M., Alverson J., VanderJagt D.J., Glew R.H. 11
A novel lectin-based enzyme-linked immunosorbent assay for the measurement of IgA1 in serum and secretory IgA1 in secretions
Cullina M.J., Greally J.F. 23
Intrathecal synthesis of IgM and IgA in neurological diseases: Comparison of two formulae with isoelectric focusing
Kaiser R., Lucking C.H. 39
Determination of acylcarnitines in urine of patients with inborn errors of metabolism using high-performance liquid chromatography after derivatization with 4′-bromophenacylbromide
Poorthuis B.J.H.M., Jille-Vlckova T., Onkenhout W. 53
Hyperphenylalaninemia and pterin metabolism in serum and erythrocytes
Ponzone A., Guardamagna O., Spada M., Ponzone R., Sartore M., Kierat L., Heizmann C.W., Blau N. 63
Release of β-hexosaminidase isoenzymes in cultured human fibroblasts
Hultberg B., Isaksson A., Nordstrom M., Kjellstrom T. 73
A ^{125}I-radioimmunoassay for diethylstilbestrol in serum of patients with prostatic cancer treated with stilphostrol
Economou E.V., Livaniou E., Evangelatos G.P., Ithakissios D.S. 81
Fetal intestinal and renal origins of trehalase activity in human amniotic fluid
Elsliger M.-A., Dallaire L., Potier M. 91
Determination of proteins by a reverse biuret method combined with the copper-bathocuproine chelate reaction
Matsushita M., Irino T., Komoda T., Sakagishi Y. 103

Lipoprotein lipase mass and activity in severe hypertriglyceridemia
Kobayashi J., Hashimoto H., Fukamachi I., Tashiro J., Shirai K., Saito Y., Yoshida S. 113
Decreased renal excretion of β-hexosaminidase in adults with insulin-dependent diabetes mellitus and normal renal function
Romero K.M., Butcher B.A., Boyle P.J., Fry D.E., Glew R.H. 125
Enzyme reactor for urinary acylcarnitines assay by reversed-phase high-performance liquid chromatography
Matsumoto K., Takahashi M., Takiyama N., Misaki H., Matsuo N., Murano S., Yuki H. 135
Erythrocyte galactokinase assay with high performance liquid chromatography
Mizoguchi N., Eguchi T., Sakura N., Ueda K. 145
Cerebrospinal fluid ferritin in brain necrosis: Evidence for local synthesis
Keir G., Tasdemir N., Thompson E.J. 153
Mono-, di- and triacylglycerol concentrations in human plasma: Effects of heparin injection and of a high-fat meal
Fielding B.A., Humphreys S.M., Allmann R.F.C., Frayn K.N. 167
Characteristics of membrane-bound 5′-nucleotidase on human prostasomes
Fabiani R., Ronquist G. 175
Hormone-sensitive lipase: Quantitation of enzyme activity and mRNA level in small biopsies of human adipose tissue
Frayn K.N., Langin D., Holm C., Belfrage P. 183
Apolipoprotein E phenotypes and genotypes as determined by polymerase chain reaction using allele-specific oligonucleotide probes and the amplification refractory mutation system in children with insulin-dependent diabetes mellitus
Stavljenic-Rukavina A., Sertic J., Salzer B., Dumic M., Radica A., Fumic K., Krajina A. 191
Letters to the Editor
Prostatic specific antigen and prostatic acid phosphatase measured by radioimmunoassay in vaginal washings from cases of suspected sexual assault
Roach B.A., Vladutiu A.O. 199

The application of molecular biology techniques to the diagnosis of hyperlipidaemia and other risk factors for cardiovascular disease
Humphries S.E. S5

Vol. 217/1, 1993

Clinical impact of genomic analysis in the diagnosis and management of genetic disease and cancer: Introduction
Werner M., Garrett C.T. 1
The diagnosis of inborn errors of metabolism by examination of the genotype
Hill R.E. 3
Genotype-phenotype correlations in phenylketonuria
Trefz F.K., Burgard P., Konig T., Goebel-Schreiner B., Lichter-Konecki U., Konecki D., Schmidt E., Schmidt H., Bickel H. 15
Cystic fibrosis: The impact of analytical technology for genotype-phenotype studies
Tummler B., Dork T., Kubesch P., Fislage R., Kalin N., Neumann T., Wulbrand U., Wulf B., Steinkamp G., Von der Hardt H. 23
Variable phenotypic expression of genotypic abnormalities in the porphyrias
Hindmarsh J.T. 29
Comparative clinical evaluation of biochemical and genomic tumor markers
Werner M., Ballo M.S., Gallagher J.V. 39
myc Amplification: Regulation of Myc function
Koskinen P.J., Makela T.P., Vastrik I., Alitalo K. 57
Genetic and molecular heterogeneity of breast cancer cells
Callahan R., Cropp C., Merlo G.R., Diella F., Venesio T., Lidereau R., Cappa A.P.M., Lisicia D.S. 63

Detection of minimal residual leukemia by the polymerase chain reaction: Potential implications for therapy
Bartram C.R. 75
Molecular diagnostics: Issues of utilization, regulation and organization
Garrett C.T., Ferreira-Centeno A., Nasim S. 85

Vol. 217/2, 1993

In vitro degradation of very low density lipoprotein from diabetic patients by lipoprotein lipase
Saheki S., Hitsumoto Y., Murase M., Takeuchi N., Uchida K. 105
Increased fragmentation of urinary fibronectin in cancer patients detected by immunoenzymometric assay using domain-specific monoclonal antibodies
Katayama M., Kamihagi K., Nakagawa K., Akiyama T., Sano Y., Ouchi R., Nagata S., Hino F., Kato I. 115
Demonstration of immunoglobulin G with affinity for dopamine in cerebrospinal fluid from psychotic patients
Bergquist J., Bergquist S., Axelsson R., Ekman R. 129
Comparison of a mechanized version of the 'Konig' reaction and a fluorescence polarization immunoassay for the determination of nicotine metabolites in urine
Pickert A., Lingenfelser T., Pickert C., Birbaumer N., Overkamp D., Eggstein M. 143
Value of faecal alpha-1-antitrypsin, haemoglobin and a chemical occult blood test in the detection of gastrointestinal disease
Moran A., Lawson N., Morrow R., Jones A., Asquith P. 153
Ascorbate concentration in human cerebrospinal fluid (CSF) and serum. Intrathecal accumulation and CSF flow rate
Reiber H., Ruff M., Uhr M. 163
Semi-automatic method for determination of different isoforms of carbohydrate-deficient transferrin
Lof K., Koivula T., Seppa K., Fukunaga T., Sillanaukee P. 175
Inactivation of glutathione peroxidase following entrapment of purified α or β hemoglobin chains in human erythrocyte
Grelloni F., Gabbianelli R., Santroni A.M., Falcioni G. 187
The preferential site of non-enzymatic glycation of human apolipoprotein A-I in vivo
Calvo C., Ulloa N., Campos M., Verdugo C., Ayrault-Jarrier M. 193
A study of growth factors in human renal cysts with or without renal cell carcinoma
Taide M., Kanda S., Eguchi J., Igawa T., Kanetake H., Saito Y. 199
Cyclooxygenase pathway in dermal fibroblasts from patients with metabolic disorders of peroxisomal origin
Malle E., Leis H.J., Steinmetz A., Paschke E., Hoefler G. 205
Studies on zinc and angiotensin-converting enzyme
Buttery J.E., Stuart S. 213
Letters to the Editor
Bile acid metabolism in three patients with mevalonic aciduria due to mevalonate kinase deficiency
Gibson K.M., Stellaard F., Hoffmann G.F., Rating D., Hrebicek M., Jakobs C. 217
Urinary excretion patterns of modified nucleosides, pseudouridine and 1-methyladenosine, in healthy individuals
Itoh K., Aida S., Ishiwata S., Sasaki S., Ishida N., Mizugaki M. 221
Fab treatment in acute digitalis intoxication: Reliability of serum digoxin determination with the Stratus system
Bizzaro N., Finco B., Milani L. 225
A rare case of IgD gammopathy associated with free light-chain cryoglobulinemia and a gelling Bence Jones protein
Myara I., Llirbat B., Rossert J., Cosson C., Tenenhaus D., Moatti N. 229
Abnormal serum lipid levels in subjects with gastric xanthoma
Chang F.-Y., Shih C.-Y., Lee S.-D. 233

Vol. 218/1, 1993

High concentrations of immunoreactive gliostatin/platelet-derived endothelial cell growth factor in synovial fluid and serum of rheumatoid arthritis
Asai K., Hirano T., Matsukawa K., Kusada J., Takeuchi M., Otsuka T., Matsui N., Kato T. 1

Human seminal deoxyribonuclease I (DNase I): Purification, enzymological and immunological characterization and origin
Yasuda T., Sawazaki K., Nadano D., Takeshita H., Nakanaga M., Kishi K. 5

A radioimmunoassay for pancreatic elastase 1: Use of α_1-antitrypsin-elastase 1 conjugate as both standard and tracer to eliminate the influence of anti-eslastase 1 autoantibodies
Asada H., Shibata K., Uchida K., Takeyama Y., Saitoh Y., Ohyanagi H. 17

Repertoires of natural autoantibodies against muscle tissue are independent of age or gender in normal human adults. A Western blot study
Zimmermann C.W. 29

Malondialdehyde adducts to, and fragmentation of, apolipoprotein B from human plasma
Lecomte E., Artur Y., Chancerelle Y., Herbeth B., Galteau M.-M., Jeandel C., Siest G. 39

Serum tissue inhibitor of metalloproteinases in patients with chronic liver disease and with hepatocellular carcinoma
Murawaki Y., Yamamoto H., Kawasaki H., Shima H. 47

A rate turbidimetric immunoassay for theophylline using biotin-avidin system
Oh C., Kim J., Kearns B., Cheng A., Dobashi T. 59

5,6-Dihydrouridine: A marker ribonucleoside for determining whole body degradation rates of transfer RNA in man and rats
Topp H., Duden R., Schoch G. 73

Comparison of lipoprotein(a) assay methods in serum and in a plasminogen-free fraction
Baldo-Enzi G., Baiocchi M., Crepaldi G. 83

Quantitation of plasma oxidatively modified low density lipoprotein by sandwich enzyme linked immunosorbent assay
Wang H., Chen S., Kong X., Wang X., Chang G., Xu S., Luo Z., Xie Y. 97

Biochemical and immunological characteristics of neutrophil alkaline phosphatase in Down's syndrome
Vergnes H.A., Denier C.J., Brisson-Lougarre A., Grozdea J.G., Bourrouillou G., Colombies P., Klaebe A.G., Perie J.J. 105

Vigabatrin — Interference with urinary amino acid analysis
Preece M.A., Sewell I.J., Taylor J.A., Green A. 113

Letter to the Editor

Oxidative metabolism and severe asthma in children
Malvy J.-M.D., Lebranchu Y., Richard M.-J., Arnaud J., Favier A. 117

Vol. 218/2, 1993

Rapid enzyme immunoassay of anti-streptokinase antibodies in human plasma
Bom V.J.J., Brugemann J., Van Der Schaaf W., Van Wijk R.T., Van Der Meer J. 121

The ins/del polymorphism in the signal sequence of apolipoprotein B has no effect on lipid parameters
Gaffney D., Freeman D.J., Shepherd J., Packard C.J. 131

Fluorescence-based diagnosis of lipid storage diseases by analysis of the culture medium of skin fibroblasts
Vered A., Eugenio M., Arie D., Augusto P., Sergio M., Shimon G. 139

Two-dimensional gel electrophoresis of four serum samples from patients with IgD myeloma
Stulik J., Tichy M., Kovarova H. 149

Rapid determination of glutamine in biological samples by high-performance liquid chromatography
Teerlink T., Hennekes M.W.T., Van Leeuwen P.A.M., Houdijk A. 159

Urinary excretion of modifoed nucleosides as biological marker of RNA turnover in patients with cancer and AIDS
Nakano K., Nakao T., Schram K.H., Hammargren W.M., McClure T.D., Katz M., Petersen E. 169
Increased excretion of endogenous urinary leukotriene E_4 in extrahepatic cholestasis
Mayatepek E., Pecher G. 185
High-performance liquid chromatographic microassay for L-glutamate:glyoxylate aminotransferase activity in human liver. Application in primary hyperoxaluria type 1
Petrarulo M., Pellegrino S., Marangella M., Cosseddu D., Linari F. 193
Simultaneous determination of lead, bismuth and thallium in plasma and urine by inductively coupled plasma mass spectrometry
Mauras Y., Premel-Cabic A., Berre S., Allain P. 201
Rapid automated enzymatic measurement of phenylalanine in plasma and blood spots
Taylor R.P., Smith I.C., Standing S.J. 207
Compact gas chromatography used with thin-layer chromatography for assessment of abnormalities of organic acids
Lee B.Y., Thurmon T.F. 215
Effects of gemfibrozil on the activities of plasma lipolytic enzymes in normolipidemic subjects
Horsmans Y., Desager J.P., Harvengt C. 223
Letter to the Editor
Variation in the amount of stimulatory GTP-binding protein (Gs) in human erythrocyte membranes in relation to alcohol consumption
Nakamura J. 229

Vol. 219/1–2, 1993

A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 1 (interstitial collagenase) using monoclonal antibodies
Zhang J., Fujimoto N., Iwata K., Sakai T., Okada Y., Hayakawa T. 1
Development and application of an enzyme immunoassay for tenascin
Washizu K., Kimura S., Hiraiwa H., Matsunaga K., Kuwabara M., Ariyoshi Y., Kato K., Takeuchi K. 15
Evaluation of thyroid function in patients with thyroid hormone autoantibodies
Sakata S., Komaki T., Ogawa T., Takuno H., Matsui I., Sarui H., Kojima N., Takamatsu J., Miura K. 23
Blood and liver lipid peroxide status after chronic ethanol administration in rats
Coudray C., Richard M.J., Faure H., Favier A. 35
Evidence for a multifactorial control of serum erythropoietin concentration in liver disease
Pirisi M., Fabris C., Falleti E., Soardo G., Toniutto P., Gonano F., Bartoli E. 47
Levels and physiochemical properties of lipoprotein subclasses in moderate hypertriglyceridemia
Manzato E., Zambon S., Zambon A., Cortella A., Sartore G., Crepaldi G. 57
Postoperative studies on parathyroid hormone secretion in patients operated on for primary hyperparathyroidism
Bergenfelz A., Norden N.E., Ahren B. 67
Bismuth in human serum: Reference interval and concentrations after intake of a therapeutic dose of colloidal bismuth subcitrate
Vanhoe H., Versieck J., Vanballenberghe L., Dams R. 79
Studies on the extraction of phospholipids from erythrocyte membranes in the rat
Eder K., Reichlmayr-Lais A.M., Kirchgessner M. 93
Neonatal screening for maple syrup urine disease by an enzyme-mediated colorimetric method
Wendel U., Gonzales J., Hummel W. 105
Dissipation of the calcium gradient in human erythrocytes results in increased heat production
Engstrom I., Waldenstrom A., Nilsson-Ehle P., Ronquist G. 113
Determination of changes in serum lathosterol during treatment with simvastatin to evaluate the role of lathosterol as a parameter for whole body cholesterol synthesis
De Cuyper I., Wolthers B.G., Van Doormaal J.J., Wijnandts P.N. 123

Serum sialic acid and acute phase proteins in type 1 and type 2 diabetes mellitus
Crook M.A., Tutt P., Simpson H., Pickup J.C. 131
Radioimmunoassay of glutathione peroxidase in human serum
Huang W., Akesson B. 139
Lipoprotein(a) phenotyping using a computerized micro scale and phenotype frequencies in a healthy Japanese population
Abe A., Noma A., Itakura H. 149
Changes in glutathione metabolic enzymes in erythrocytes of Plasmodium vivax infected patients
Mathews S.T., Selvam R. 159
The vitamin A fat-loading test in young normolipidemic subjects
Heller F.R., Vandenplas C., Desager J.P., Harvengt C. 167
Abnormally fucosylated haptoglobin as a marker for alcoholic liver disease but not excessive alcohol consumption or non-alcoholic liver disease
Chambers W., Thompson S., Skillen A.W., Record C.O., Turner G.A. 177
Differences in the time course of creatine kinase-MB activity and mass concentration after acute myocardial infection
Baum H.H.E., Schwab I., Boekstegers P., Steinbeck G., Neumeier D. 183
Simple mixing technique for the preparation of blood samples of known oxygen saturation
Clark D.J. 189
Letter to the Editor
Effect of ethanol on prolidase I and prolinase activity in the human hepatoma cell line Hep G_2
Myara I., Cosson C., Pourci M.L., Moatti N., Lemonnier A. 195

Vol. 220/1, 1993

HPLC determination of oxidized and reduced pyridine coenzymes in human erythrocytes
Micheli V., Simmonds H.A., Bari M., Pompucci G. 1
Urinary growth hormone measurements as a marker of renal tubular function in diabetes mellitus
Turner G., Coates P., Porter S., Peters J.R., Woodhead J.S. 19
A one-step sandwich enzyme immunoassay for tissue inhibitor of metalloproteinases-2 using monoclonal antibodies
Fujimoto N., Zhang J., Iwata K., Shinya T., Okada Y., Hayakawa T. 31
Regulation of transferrin receptor expression and distribution in vitro cultured human cytotrophoblasts
Starreveld J.S., Van Dijk H.P., Kroos M.J., Van Eijk H.G. 47
Scintillation proximity assay for calcitriol in serum without high pressure liquid chromatography
Wildermuth S., Dittrich K., Schmidt-Gayk H., Zahn I., O'Riordan J.L.H. 61
A fibroblast glutaryl-CoA dehydrogenase assay using detritiation of ^{3}H-labelled glutaryl-CoA: Application in the genotyping of the glutaryl-CoA dehydrogenase locus
Christensen E. 71
The hepatobiliary disease marker serum lanine aminopeptidase predominantly comprises an isoform of the haematological myeloid differentiation antigen and leukaemia marker CD-13/gp150
Favaloro E.J., Browning T., Nandurkar H. 81
An improved Bradford protein assay for collagen proteins
Lopez J.M., Imperial S., Valderramad R., Navarro S. 91
False rise in C-reactive protein in a pateint with monoclonal IgM immunoglobulin
Ponge T.D., Le Carrer D.L., Sagniez M.M., Pontet M., Cottin S.L. 101
The glycosylation of haptoglobin in rheumatoid arthritis
Thompson S., Dargan E., Griffiths I.D., Kelly C.A., Turner G.A. 107
Letters to the Editor
Senile cataractic lenses do not accumulate galactitol in either lactose tolerant or intolerant subjects
Wacker H., Soldati L., Simonelli F., Richter C., Gazzaniga A., Auricchio S., Semenza G. 115
Non-enzymatic glycosylation of mouse monoclonal antibody reduces its binding activity to antigen
Sasaki Y., Mori T., Shiiki H., Dohi K., Ishikawa H. 119

Measurement of seminal plasma α-amylase that is not inhibited by monoclonal antibodies against the salivary isoenzyme
Huguet J., Cortes J.D., Arranz B., Fuentes-Arderiu X. 123

Vol. 220/2, 1993

Urinary macromolecules are promoters of calcium oxalate nucleation in human urine: Turbidimetric studies
Rodgers A.L., Ball D., Harper W. 125
Increase in plasma concentration of ubiquitin in dialysis patients: Possible involvement in β_2-microglobulin amyloidosis
Okada M., Miyazaki S., Hirasawa Y. 135
Determination of human immunoglobulin A and secretory immunoglobulin A in bronchoalveolar lavage fluids by solid phase enzyme immunoassay
Sato T., Kitajima A., Ohmoto S., Chikuma M., Kado M., Nagai S., Izumi T., Takeyama M., Masada M. 145
Quantification of 3-methylglutaconic acid in urine, plasma, and amniotic fluid by isotope-dilution gas chromatography/mass spectrometry
Kelley R.I. 157
Improved PCR/NcoI method for the molecular diagnosis of medium chain acyl-CoA dehydrogenase deficiency using dried blood samples: Two-stage amplification using two different sets of primers improves accuracy and sensitivity
Nagao M., Raymond D., Kim J., Tanaka K. 165
Alterations in the concentration of an apolipoprotein E-containing subfraction of plasma high density lipoprotein in coronary heart disease
Wilson H.M., Patel J.C., Russell D., Skinner E.R. 175
2,4-Diamino-7-hydroxy-pteridines in biological fluids of patients on high-dose methotrexate
Dhondt J.-L., Millot F., Hayte J.-M., Mazingue F., Farriaux J.-P. 189
Sensitive chemiluminescent immunoassay for human granulocyte colony-stimulating factor (G-CSF) in clinical applications
Kiriyama R., Chichibu K., Matsuno T., Ohsawa N. 201
Anion exchange protein (band 3) in human erythrocytes characterized by different abnormalities
Giuliani A., Marini S., Ferroni L., Condo S.G., Ramacci M.T., Giardina B. 211
Investigation of β-oxidation in human liver biopsies
Eaton S., Record C.O., Bartlett K. 219
Letters to the Editor
Enzyme-linked immunoglobulin which binds both lactate dehydrogenase and hydroxybutyric dehydrogenase
Taguchi H., Nishiya K., Yamanaka S., Saika K., Morishita N., Sasaki M., Miyoshi I. 225
A simple coloured-paper thin-layer chromatography test as a backup procedure for vanillylmandelic acid and homovanillic acid evaluation in a neuroblastoma screening program
Mathieu P., Philip T., Greffe J., Montegue A., Lacroix C. 229
A comparative study of thyroid function in African neonates: A reference thyroid profile
Das S.C., Isichei U.P. 233

Vol. 221/1–2, 1993

An investigation of argininosuccinic acid anhydrides in argininosuccinic acid lyase deficiency by ^{1}H-NMR spectroscopy
Burns S.P., Iles R.A. 1
Is an increase of deoxyribonucleases activity in uraemic lymphocytes caused by their augmented synthesis in these cells?
Adamiec R., Szewczyk Z., Szmidzinski R., Szopa J. 15

Microwave oven digestion procedure for atomic absorption spectrometry analysis of bone and teeth
Gil F., Perez M.L., Facio A., Villanueva E., Tojo R., Gil A. 23
Insulin receptor binding to erythrocytes in the first half of pregnancy is increased in healthy pregnant women as compared with non-pregnant or gestational diabetic women
Schmon B., Desoye G., Friedl H., Hofmann H., Weiss P.A.M., Hagmuller K. 33
The lysosomal N-acetyl-β-D-glucosaminidase (NAG) isoenzymes in plasma: Study of distribution in a general population by a simple routine chromatofocusing procedure
Goi G., Bairati C., Roggi C., Maccarini L., Tettamanti G., Meloni Lombardo C.A. 47
Changes in the glycoforms of rat alpha-1-acid glycoprotein during experimental polyarthritis
Venembre P.C., Cong H.N., Biou D.R., Durand G.M. 59
Cholesteryl ester transfer activity in lipoprotein lipase deficiency and other primary hypertriglyceridemias
Iglesias A., Contreras J.A., Martinez-Pardo M., Entrala A., Herrera E., Lasuncion M.A. 73
A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 2 (72-kDa gelatinase/type IV collagenase) using monoclonal antibodies
Fujimoto N., Mouri N., Iwata K., Ohuchi E., Okada Y., Hayakawa T. 91
Expression and nature of the alkaline phosphatase gene in cultured osteosarcoma cells
Stinson R.A., Thacker J.D., Lin C.C. 105
A novel HPLC method for the simultaneous quantification of monosaccharides and disaccharides used in tests of intestinal function and permeability
Sorensen S.H., Proud F.J., Adam A., Rutgers H.C., Batt R.M. 115
LDL sialic acid content in patients with coronary artery disease
Ruelland A., Gallou G., Legras B., Paillard F., Cloarec L. 127
The investigation of UDPGlucose and UDPGalactose concentration in red blood cells of patients with classical galactosaemia
Keevill N.J., Holton J.B., Allen J.T. 135
High performance liquid chromatographic profiling of tryptophan and related indoles in body fluids and tissues of carcinoid patients
Kema I.P., Schellings A.M.J., Hoppenbrouwers C.J.M., Rutgers H.M., De Vries E.G.E., Muskiet F.A.J. 143
Characterization of apo(a) polymorphism by a modified immunoblotting technique in an Italian population sample
Geroldi D., Bellotti V., Buscaglia P., Bonetti G., Gazzaruso C., Caprioli A., Fratino P. 159
Diagnosis of mucopolysaccharidoses in a clinically selected population by urinary glycosaminoglycan analsyis: A study of 2,000 urine samples
Piraud M., Boyer S., Mathieu M., Maire I. 171
Serum bile acid concentrations in mild liver cirrhosis
Greco A.V., Mingrone G. 183
An increase of serum remnant-like particles in non-insulin-dependent diabetic patients with microalbuminuria
Shimizu H., Mori M., Saito T. 191
Study of chronic granulomatous disease by a nitroblue tetrazolium densitometric kinetic test: A new research method
Iacobini M., Torre A., Palumbo G., Del Principe D., Digilio G., Balducci L. 197
β-N-Acetylhexosaminidase in spinal muscular atrophy fibroblasts
Tassi C., Beccari T., Costanzi E., Orlacchio A. 203
Enzyme immunoassay of serum type IV collagen in anti-HCV positive chronic liver diseases
Matsuzaki Y., Yoshiga S., Sugitani T., Shoda J., Tanaka N., Osuga T. 209
The validity of the ethanol precipitation method for the measurement of mucin content in human gastric juices and its possible relationship to gastroduodenal diseases
Azuumi Y., Ichikawa T., Ishihara K., Hotta K. 219

Vol. 222/1–2, 1993

Introduction — The OpenLabs Project
O'Moore R. 1
Object-oriented methods
Solberg H.E. 3
Practical experience with graphical user interfaces and object-oriented design in the clinical laboratory
Wells I.G., Cartwright R.Y., Farnan L.P. 13
RefVal for Windows — migrating to the Windows environment with the help of Visual Basic
Kenny D., Solberg H.E. 19
Supporting interoperability between medical knowledge-based systems: Experience from pilot implementations
Boran G.P., Given P.G., Grimson J.B., O'Moore R.R. 23
Interpretative reporting and alarming based on laboratory data
Nykanen P., Boran G., Pince H., Clarke K., Yearworth M., Willems J.L., O'Moore R. 37
To test or not to test, that is the question
Hasman A., Pop P., Winkens R.A.G., Blom J.L. 49
Problem-oriented management of laboratory work through dynamic test scheduling
Brender J., Magdal U., Wiegell B., Schioler T., McNair P. 57
Knowledge based lipid management system for general practitioners
Sinnott M.M., Carr B., Markey J., Brosnan P., Boran G., Grimson J., McGill T., Feely J., O'Moore R. 71
Knowledge-based approach to clinical decision-support system, with an application in tetanus serology
Schroder J.P., Kuhlmann W.D., Trendelenburg C. 78
Competence reasoning — Handling ambiguous and imprecise data in the Pro.M.D. expert system shell
Pohl B. 85
Integrating knowledge-based systems and databases
Grimson J.B. 101
Clinical utility of an international thyroid database
Nolan J., Schioler T., Brosnan P., Irjala K., Nuutila P., McNair P., O'Moore R. 117
Arden Syntax as a standard for knowledge bases in the clinical chemistry laboratory
Johansson B., Bergvin L. 123
Analytical quality goals and assessment to ensure transferability of laboratory results
Groth T., De Verdier C.-H. 129
Distribution of laboratory test results to primary health care centres with the EDIFACT standard
Bergvin L., Johansson B., Borjesson U. 141
Guidelines for hospital laboratory computer systems
Kenny D. 147
The Ninth International Conference on Computing in Clinical Laboratories: Foreword
Kenny D., Solberg H.E. vii

Vol. 223/1–2, 1993

Comparative drug influence on peroxide mediated increase of cytosolic calcium in human endothelial cells
David M., Horvath G., Schimke I., Nagy I., Mueller M.M. 1
Metabolites of dietary (soya) isoflavones in human urine
Kelly G.E., Nelson C., Waring M.A., Joannou G.E., Reeder A.Y. 9
Excretion of 3,6-epoxydicarboxylic acids in peroxisomal disorders
Pitt J.J., Poulos A. 23
Immunonephelometric/turbidimetric apolipoprotein B assays for the clinical laboratory
Levinson S.S., Wagner S.G. 31

Effect of tamoxifen on serum cholesterol and lipoproteins during chemohormonal therapy
Dnistrian A.M., Schwartz M.K., Greenberg E.J., Smith C.A., Schwartz D.C. 43
Cholesterol in remnant-like lipoproteins in human serum using monoclonal anti apo B-100 and anti apo A-I immunoaffinity mixed gels
Nakajima K., Saito T., Tamura A., Suzuki M., Nakano T., Adachi M., Tanaka A., Tada N., Nakamura H., Campos E., Havel R.J. 53
The relationship between serum levels of lipoprotein(a) and proteins associated with the acute phase response
Ledue T.B., Neveux L.M., Palomaki G.E., Ritchie R.F., Craig W.Y. 73
Lactate dehydrogenase isoenzyme patterns in blood cells from histiocytosis X children
Perlino E., Marra E., Maenza S., De Terlizzi M., Coppola B.C., Santostasi T., Quagliariello E. 83
High adenosine content in human uterine smooth muscle compared with striated skeletal muscle
Ronquist G., Wedenberg K., Waldenstrom A., Ericson A., Ulmsten U. 93
Bilirubin inhibits protein kinase C activity and protein kinase C-mediated phosphorylation of endogenous substrates in human skin fibroblasts
Amit Y., Boneh A. 103
Lipoprotein composition in the insulin-deficient non-acidotic phase of type I diabetic patients and early evolution after the start of insulin therapy
Reverter J.L., Senti M., Rubies-Prat J., Lucas A., Salinas I., Pizarro E., Pedro-Botet J., Sanmarti A. 113
Lipoprotein(a) serum concentration in familial combined hyperlipidemia
Fonda M., Da Col P.G., La Verde R., Battello C., Fisicaro M., Tonizzo M., Cattin L. 121
Effect of bone resorbing factors on u-PA and its specific receptor in osteosarcoma cell line
Nonaka T., Matsumoto H., Shimada W., Okada K., Fukao H., Ueshima S., Kikuchi H., Tanaka S., Matsuo O. 129
Extremely high HDL levels in a patient with multiple symmetric lipomatosis
Deiana L., Pes G.M., Carru C., Campus G.V., Tidore M.G.B., Cherchi G.M. 143
An enzymatic assay of inorganic phosphate in serum using nucleoside phosphorylase and xanthine oxidase
Ungerer J.P.J., Oosthuizen H.M., Bissbort S.H. 149
Free thyroxine measured in undiluted serum by dialysis and ultrafiltration: Effects of non-thyroidal illness, and an acute load of salicylate or heparin
Faber J., Waetjen I., Siersbaek-Nielsen K. 159
Total pancreatic and salivary serum iso-amylase activities in alcohol misusers in relapse and remission and in alcoholic liver disease
Deenmamode J.M.J.M., Sherwood R.A., Sherman D.I.N., Peters T.J. 169
Prenatal screening for Down's syndrome and human chorionic gonadotropin isoforms
Defasque P., Pigny P., Racadot A., Farriaux J.-P., Boersma A. 173
Stability of human plasma atrial natriuretic peptide during storage at −80°C
Bhaggoe U.M., Boomsma F., Admiraal P.J.J., Man in 't Veld A.J., Schalekamp M.A.D.H. 179
Letters to the Editor
Pedigree of serum extracellular-superoxide dismutase level
Adachi T., Nakamura M., Yamada H., Kitano M., Futenma A., Kato K., Hirano K. 185
The molecular mass and concentrations of protein 1 or Clara cell protein in biological fluids: A reappraisal
Bernard A., Dumont X., Roels H., Lauwerys R., Dierynck I., De Ley M., Stroobant V., De Hoffmann E. 189

Vol. 224/1, 1994

Alteration of asparagine-linked glycosylation in serum transferrin of patients with hepatocellular carcinoma
Matsumoto K., Maeda Y., Kato S., Yuki H. 1
Non-specific steroidal esterase activity and distribution in human and other mammalian tissues
Lund-Pero M., Jeppson B., Arneklo-Nobin B., Sjogren H.-O., Holmgren K., Pero R.W. 9

Very-fast ultracentrifugation of human plasma lipoproteins: Influence of the centrifugal field on lipoprotein composition
Leonhardt W., Pietzsch J., Nitzsche S. 21
Glomerular filtration rate and creatinine production in adult icteric patients
Delanghe J.R., Louagie H.K., De Buyzere M.L., Leroux-Roels G.G. 33
The allopurinol loading test for identification of carriers for ornithine carbamoyl transferase deficiency: Studies in a healthy control population and females at risk
Sebesta I., Fairbanks L.D., Davies P.M., Simmonds H.A., Leonard J.V. 45
A simplified method for the determination of phosphoribosylpyrophosphate synthetase activity in hemolysates
Torres R.J., Mateos F.A., Puig J.G., Becker M.A. 55
Effects of alpha-1 acid glycoprotein on human polymorphonuclear neutrophils: Influence of glycan microheterogeneity
Vasson M.P., Roch-Arveiller M., Couderc R., Baguet J.C., Raichvarg D. 65
Detection of viral DNA in hybridization reactions using two chemiluminescent substrates
Girotti S., Ferri E., Musiani M., Gibellini D., Zerbini M., Roda A., Gentilomi G., Venturoli S. 73
Application of ^{1}H-NMR spectroscopy for qualitative measurement of muscle carnitine levels
Nakajima H., Sugie H., Tsurui S., Ito M. 81
Relationship of plasma malondialdehyde, vitamin E and antioxidant micronutrients to human immunodeficiency virus-1 seropositivity
Malvy D.J.-M., Richard M.-J., Arnaud J., Favier A., Amedee-Manesme O. 89
Development of a radioimmunoassay for fetal antigen 2
Price K.M., Silman R., Armstrong P., Grudzinskas J.G. 95
Letters to the Editor
Improvement of ELISA sensitivity by allogeneric adsorption of polyclonal antibodies — A technical note for non-experts
Abe Y., Kimura K., Horiuchi A., Miyake M., Kimura S. 103
Effect of short-term and long-term alcohol intake on plasma cholesterol in non-alcoholic humans
Nakamura J. 107

Vol. 224/2, 1994

A new case of apo C-II deficiency with a nonsense mutation in the apo C-II gene
Zanelli T., Catapano A.L., Averna M.R., Barbagallo C.M., Liotta A., Giardina F.C., Notarbartolo A. 111
Analytical goals for rectilinear calibration functions
Fuentes-Arderiu X., Alia P., Machuca I. 119
Relationship between lipoprotein(a) levels in serum and some indices of protein synthesis in liver cirrhosis
Alessandri C., Basili S., Maurelli M., Andreozzi P., Violi F., Cordova C. 125
Genotyping compared with protein phenotyping the common apolipoprotein E polymorphism
Hansen P.S., Gerdes L.U., Klausen I.C., Gregersen N., Faergeman O. 131
Serum octadeca-9,11 dienoic acid — An assay of free radical activity or a result of bacterial production?
Jack C.I.A., Ridgway E., Jackson M.J., Hind C.R.K. 139
The esters of p-hydroxy-benzoate (parabens) inhibit the release of lysosomal enzymes by mitogen-stimulated peripheral human lymphocytes in culture
Bairati C., Goi G., Lombardo A., Tettamanti G. 147
Therapeutic whole-blood levels of cyclosporin A in heart and lung transplantation
Andert S.E., Grimm M., Schreiner W., Muller M.M. 159
Erythrocyte Na^{+},K^{+}-ATPase properties and adenylate energy charge in normotensives and in essential hypertensives
Canestrari F., Galli F., Boschi S., Albertini M.C., Gheller G., De Crescentini S., Bossu M. 167
HPLC assay of conjugated bile acids in human fluids using on-line sample pretreatment on a standard isocratic chromatograph

Scalia S., Pazzi P. 181
Light chains of immunoglobulins in human secretions
Mole C.M., Bene M.C., Montagne P.M., Seilles E., Faure G.C. 191
Acute phase proteins and lipoprotein(a) in patients with severe hypercholesterolaemia and normal subjects
Crook M.A., Haq S., Chusney G., Haq M., Tutt P. 199
Creatine kinase isoenzyme MB mass concentrations in patients undergoing aortocoronary bypass surgery
Mair P., Mair J., Seibtb I., Antretter H., Balogh D., Puschendorf B. 203
Tissue polypeptide specific antigen (TPS) in liver disease
Moraglio D., Pagano M., Galeazzi D., Arnelli A., Bertero L. 209
Letters to the Editor
The effect of silicon as silicic acid on superoxide dismutase activity
Bellia J.P., Birchall J.D., Roberts N.B. 215
Concanavalin A affinity pattern of CA 19-9: An in vitro and in vivo study
Basso D., Panozzo M.P., Del Giudice G., Carbone M.D., Meggiato T., Del Favero G., Plebani M. 219

The clinical laboratory is in the information business
Weilert M. S1
Interpreting the results is the expertise of the laboratory
Winkel P. S9
The role of suppliers and laboratories in the new information environment
Korpman R.A. S23

Vol. 225/1, 1994

Glycosylation of gamma-glutamyltransferase is modified by ethanol in H5-6 hepatoma cell line
Odoul M., Bagrel D., Peyrieras N., Bodaud H., Siest G. 1
A solid phase enzyme linked immunofiltration assay for secretory leucocyte-proteinase inhibitor
Melrose J., Ghosh P., Jacoby A. 17
GC-MS determination of ratios of stable-isotope labelled to natural urea using [$^{13}C^{15}N_2$]urea for studying urea kinetics in serum and as a means to validate routine methods for the quantitative assay of urea in dialysate
Wolthers B.G., Tepper T., Withag A., Nagel G.T., de Haan T.H.Y., van Leeuwen J.J., Stegeman C.A., Huisman R.M. 29
ELISA for the core protein of the cartilage large aggregating proteoglycan, aggrecan: Comparison with the concentrations of immunogenic keratan sulphate in synovial fluid, serum and urine
Moller H.J., Larsen F.S., Ingemann-Hansen T., Poulsen J.H. 43
Serum carnosinase activities in central nervous system disorders
Wassif W.S., Sherwood R.A., Amir A., Idowu B., Summers B., Leigh N., Peters T.J. 57
Increased proteolysis of oxidatively damaged hemoglobin in erythrocyte lysates in diabetes mellitus
Raghothama C., Rao P. 65
Increases in serum β3 integrin in patients with chronic liver disease
Yamauchi M., Mizuhara Y., Takahashi A., Toda G. 71
Phenotyping is an accurate means of analysing the principal apolipoprotein E isoforms
James R.W., Ruiz J., Blanche H., Pometta D., Passa P., Froguel P. 77
Letters to the Editor
The topography of thiol distribution on sperm surface: A sensitive marker for human fertility diagnosis
Sinha S., Kumar G.P., Laloraya M. 83
Polymorphism at the glutathione s-transferase GSTMl locus: A study of the frequencies of the GSTMl a, b, A/B and null phenotypes in nigerians
Zhao L., Alldersea J., Fryer A., Tighe A., Ollier B., Thomson W., Jones P., Strange R. 85

Enzyme reference materials: Their place in diagnostic enzymology
Moss D.W. S5
Systematic top-down approach to clinical chemistry
Pardue H.L. S13
Molecular biology techniques in the diagnosis of monogenic diseases
Wagener C., Epplen J.T., Erlich H., Peretz H., Vihko P. S35

Vol. 225/2, 1994

Elevated serum and urinary thrombomodulin levels in patients with non-insulin-dependent diabetes mellitus
Shimano H., Takahashi K., Kawakami M., Gotoda T., Harada K., Shimada M., Yazaki Y., Yamada N. 89
Vitamin E and the hypercoagulability of neonatal blood
Jain S.K., McCoy B., Wise R. 97
Fully mechanized latex immunoassay for serum lipoprotein(a)
Abe A., Yoshimura Y., Sekine T., Maeda S., Yamashita S., Noma A. 105
Sex hormone binding globulin phenotypes: Their detection and distribution in healthy adults and in different cinical conditions
Cornelisse M.M., Bennett P.E., Christiansen M., Blaakaer J., Gluud C., Andersen J.R., Homann C., Garred P. 115
Anti-thyroid peroxidase (anti-TPO) antibodies in thyroid diseases, non-thyroidal illness and controls. Clinical validity of a new commercial method for detection of anti-TPO (thyroid microsomal) autoantibodies
Engler H., Riesen W.F., Keller B. 123
Monitoring of osteoblast activity with an immunoradiometric assay for determination of bone alkaline phosphatase mass concentration in patients receiving renal transplants
Withold W., Degenhardt S., Castelli D., Heins M., Grabensee B. 137
Changes in purine nucleotide content in the lymphocyte subpopulations of patients infected with HIV
Tabucchi A., Carlucci F., Consolmagno E., Monari P., Re M.C., Marinello E., Pizzichini M., Pagani R., Lai C.Y. 147
A rapid and sensitive sandwich enzyme immunoassay for low concentration of albumin in human urine
Chiabrando G.A., Zalazar F.E., Aldao M.A.J., Vides M.A. 155
Feto-maternal plasma phenylalanine concentration gradient from 19 weeks gestation to term
Schoonheyt W.E., Clarke J.T.R., Hanley A.B., Johnson J.M., Lehotay D.C. 165
Stability of human atrial natriuretic peptide in blood samples
Tsuji T., Masuda H., Imagawa K., Haraikawa M., Shibata K., Kono M., Inouye K., Uchida K. 171
The glycosaminoglycan composition of human tracheas and the changes observed during aging and in disease
Binette J.P., Burgi W., Ohishi H., Grundboeck-Jusko J., Burki R., Maekawa Y., Tschopp T.A., Kimura A., Schmid K. 179
1,25-Dihydroxyvitamin D radioreceptor assay using bovine mammary gland receptor non-high performance liquid chromatographic purification
Watanabe Y., Kubota T., Suzumura E., Suzuki T., Yonezawa M., Ishigamia T., Ichikawa M., Seino Y. 187
Effect of storage of sera and apolipoprotein(a) phenotypes on detection of lipoprotein(a) by a new agarose gel
Roberta B., Grandhomme M., Mainard F., Madec Y. 195

Vol. 226/1, 1994

Rapid and accurate random urinary porphyrin quantitation
Boynton S.B., Roth K.S. 1

Serum levels of steroid sulfatase protein in gynecologic carcinomas
Sugawara T., Honke K., Gasa S., Tanaka T., Fujimoto S., Makita A. 13
Quantitative analysis of Lewis antigens on erythrocytes by flow cytometry
Murai J., Naka K., Shimojo N., Katakami T., Nakagishi M., Kuroki T., Okuda K. 21
Glycosaminoglycan content is increased in dissecting aneurysms of human thoracic aorta
Cattell M.A., Hasleton P.S., Anderson J.C. 29
Clinical evaluation of biotin-binding immunoglobulin in patients with Graves' disease
Nagamine T., Takehara K., Fukui T., Mori M. 47
Demonstration and some properties of N-acetyl-β,D-hexosaminidase (HEX) C isoenzyme in human renal tissues: Relative increase of HEX C activity in renal cell carcinoma
Yoshida K.-I. 55
Sensitive colorimetric assay of serum diamine oxidase
Takagi K., Nakao M., Ogura Y., Nabeshima T., Kunii A. 67
Rapid detection of common metachromatic leukodystrophy mutations by restriction analysis of arylsulfatase A gene amplimers
Ben-Yoseph Y., Mitchell D.A. 77
Isolation of fetal antigen 2 and assay standardisation
Price K.M., Silman R., Grudzinskas J.G. 83
Rapid assay for furancarboxylic acid accumulated in uremic serum using high-performance liquid chromatography and on-line mass spectrometry
Niwa T., Kawagishi I., Ohya N. 89
A rapid semi-micro method for the separation of lipoprotein fractions that uses a benchtop ultracentrifuge
Fletcher C.D., Barnes J.F., Farish E. 95
Letters to the Editor
Suitability of copper chloride as a reaction terminator for superoxide dismutase activity assay
Sun Y., Oberley L.W. 101
Lipoprotein(a) concentration in various hyperlipidemias
Alberty R. 105
Automated kinetic assay for endotoxin and (1 → 3)-β-D-glucan in human blood
Tamura H., Arimoto Y., Tanaka S., Yoshida M., Obayashi T., Kawai T. 109
Increased levels of plasma endothelin-1 in patients with psoriasis
Cecchi R., Giomi A., Ciuti M., Gironi A., Seghieri G. 113

Increasing the biosafety of analytical systems in the clinical laboratory
Truchaud A., Schnipelsky P., Pardue H.L., Place J., Ozawa K. S5

Vol. 226/2, 1994

Human prostatic acid phosphatase: Selected properties and practical applications
Ostrowski W.S., Kuciel R. 121
Release of membrane-bound enzymes from cells and the generation of isoforms
Moss D.W. 131
Bone-origin alkaline phosphatase in plasma by wheat-germ lectin methods in bone disease
Rosalki S.B. 143
Monitoring skeletal cancer metastases with the bone isoenzyme of tissue unspecific alkaline phosphatase
Burlina A., Rubin D., Secchiero S., Sciacovelli L., Zaninotto M., Plebani M. 151
Antenatal diagnosis of tetrahydrobiopterin deficiency by quantification of pterins in amniotic fluid and enzyme activity in fetal and extrafetal tissue
Blau N., Kierat L., Matasovic A., Leimbacher W., Heizmann C.W., Guardamagna O., Ponzone A. 159
Drug metabolizing enzymes related to laboratory medicine: Cytochromes P-450 and UDP-glucuronosyltransferases
Batt A.M., Magdalou J., Vincent-Viry M., Ouzzine M., Fournel-Gigleux S., Galteau M.M., Siest G. 171

Enzymes and circulating proteins as markers of alcohol abuse
Goldberg D.M., Kapur B.M. 191
The enzymes of mitochondrial fatty acid oxidation
Bennett M.J. 211
Detection of point mutations in human genes by the solid-phase minisequencing method
Syvanen A.-C. 225
Viral diagnosis using hybridization assays with digoxigenin labeled probes
Musiani M., Zerbini M., Gibellini D., Venturoli S., Gentilomi G., Gallinella G., La Placa M. 237
Screening for cystic fibrosis mutations in Spanish patients
Molano J., Ezquieta B., Granell R. 247

Vol. 227/1–2, 1994

Inflammation-associated changes in the cellular availability of tryptophan and kynurenine in renal transplant recipients
Holmes E.W., Russell P.M., Kinzler G.J., Bermes Jr. E.W. 1
Initiation codon mutation of the tyrosinase gene as a cause of human albinism
Breimer L.H., Winder A.F., Jay B., Jay M. 17
Neurotrophin-3 (NT-3) levels in the developing rat nervous system and in human samples
Murase K., Igarashi K., Hayashi K. 23
Role of the measurement of serum procollagen type III N-terminal peptide in the evaluation of liver diseases
Collazos J., Diaz F. 37
Factors affecting the thiobarbituric acid test as index of red blood cell susceptibility to lipid peroxidation: A multivariate analysis
Girelli D., Olivieri O., Stanzial A.M., Guarini P., Trevisan M.T., Bassi A., Corocher R. 45
Serum beta-hexosaminidase isoenzymes are related to risk factors for atherosclerosis in a large population of postmenopausal women
Hultberg B., Isaksson A., Nilsson J.A., Lindgarde F. 59
Monosaccharide composition of haptoglobin in liver diseases and alcohol abuse: Large changes in glycosylation associated with alcoholic liver disease
Mann A.C., Record C.O., Self C.H., Turner G.A. 69
An HPLC-linked assay of phosphoribosylpyrophosphate synthetase activity in the erythrocytes of adults and children with neurological disorders
Micheli V., Rocchigiani M., Pompucci G. 79
Biological markers and ovarian carcinomas: Galactosyltransferase, CA 125, isoenzymes of amylase and alkaline phosphatase
Sichel F., Salaun V., Bara E., Gauduchon P., Malas J.-P., Goussard J., Le Talaer J.-Y. 87
Acidification reveals a greater number of epidermal growth factor receptors in human placental and breast cancer membranes
Hatzoglou A., Romain S., Dalapascha H., Castanas E. 97
Separation of alkaline phosphatase isoforms with and without intact glycan-phosphatidylinositol anchors in aqueous polymer phase systems
Raymond F.D., Moss D.W., Fisher D. 111
[^{14}C]Methylamine accumulation in cultured human skin fibroblasts a biochemical test for lysosomal storage and lysosomal diseases
Kopitz J., Gerhard C., Hofler P., Cantz M. 121
Plasma fibronectin and angiotensin-converting enzyme: Markers of primary pulmonary injury in burn patients
Plassart F., Cynober L., Baudin B., Lioret N., Beneteau-Burnat B., Nolland X.B., Saizy R., Giboudeau J. 135
Optimized determination of angiotensin I-converting enzyme activity with hippuryl-L-histidyl-L-leucine as substrate
Schnaith E., Beyrau R., Buckner B., Klein R.M., Rick W. 145

A rapid and simple method for measuring the susceptibility of low-density-lipoprotein and very-low-density-lipoprotein to copper-catalyzed oxidation
Zhang A., Vertommen J., van Gaal L., De Leeuw I. 159
Comparison of enhanced chemiluminescence and colorimetric techniques for the immune-detection of α-antitrypsin
Venembre P., Seta N., Boutten A., Dehoux M., Aubier M., Durand G. 175
Use of HPLC to measure circulating adenosine levels in migrainous patients
Guieu R., Sampieri F., Bechis G., Rochat H. 185
Relationship between ulinastatin and alpha-1-microglobulin in human urine
Shikimi T., Himeno Y., Shigeno K., Gonda T., Ishibe T., Hattori K., Takaori S. 195
Measurement of the oxidisable polyamine content of seminal plasma
Fernandez C., Sharrard R.M., Monks N., Barrett C.L.R. 201
Letters to the Editor
Detection of modified β_2-microglobulin (β_2m) from amyloid deposits in tenosynovial tissue of carpal tunnel syndrome (CTS)
Odani H., Mizutani A., Hidaka H., Fujisawa K., Shinzato T., Maeda K. 209
Examination of bronchoalveolar wash fluid for ras, myc and fos oncoproteins and glutathione S-transferase isoenzymes: An attempt to improve the accuracy of lung cancer diagnosis
Pinkham J.M., Strange R.C., Pantin C. 211

Vol. 228/1, 1994

ClinBio 92' and 93': Introduction
Salvatore F., Pane F. 1
Greater differences in forensic DNA profile frequencies estimated from racial groups than from ethnic subgroups
Budowle B., Monson K.L. 3
Magnetic resonance imaging and spectroscopy of MCF7 human breast cancer: Pathophysiology and monitoring of treatment
Degani H., Furman E., Fields S. 19
Biochemical and molecular investigations in respiratory chain deficiencies
Rustin P., Chretien D., Bourgeron T., Gerard B., Rotig A., Saudubray J.M., Munnich A. 35
Reference charts for respiratory chain activities in human tissues
Chretien D., Rustin P., Bourgeron T., Rotig A., Saudubray J.M., Munnich A. 53

Vol. 228/2, 1994

Estimation of pS2 protein level in human body fluids by a sensitive two-site enzyme immunoassay
Miyashita S., Nomoto H., Konishi H., Hayashi K. 71
A detection method for point mutation in the precore region of human hepatitis B virus (HBV)-DNA using mutation-site-specific assay
Kinoshita M., Seno T., Fukui T., Shin S., Tsubota A., Kumada H. 83
Elevation of serum group II phospholipase A_2 levels in patients with advanced cancer
Yamashita S., Ogawa M., Sakamoto K., Abe T., Arakawa H., Yamashita J. 91
Evaluation of tumor-associated antigen (2H6 antigen) in detecting early stages of gastric cancer
Harada H., Tsukada Y., Karasawa Y. 101
Thyroid hormone autoantibodies in patients with Graves' disease: Effect of anti-thyroid drug treatment
Ogawa T., Sakata S., Nakamura S., Takuno H., Matsui I., Sarui H., Yasuda K. 113
Tubular nephrotoxicity after cardiac surgery utilising cardiopulmonary bypass
Westhuyzen J., McGiffin D.C., McCarthy J., Fleming S.J. 123
A new precipitation method with magnetic separation for high-density-lipoprotein cholesterol assay
Iida S., Osawa S., Yonemitsu H. 133

Cathepsin B and N-acetyl-β-D-glucosaminidase in human synovial cells in culture: Effects of interleukin-1
Lecomte V., Knott I., Burton M., Remacle J., Raes M. 143
Modulation defect of sodium pump evidenced in diabetic patients by a microcalorimetric study
Issautier T., Kovacic H., Gallice P., Raccah D., Vague P., Crevat A. 161
Lipoprotein profiling by high performance gel chromatography
Tallis G.A., Shephard M.D.S., Whiting M.J. 171
Interaction of β-cyclodextrin with bile acids and their competition with vitamins A and D3 as determined by ^{1}H-NMR spectrometry
Comini S., Olivier P., Riottot M., Duhamel D. 181
Characterization of tumor-associated antigen (2H6 Antigen) defined by monoclonal antibody 2H6
Harada H., Kubota T., Tsukada Y., Karasawa Y. 195
Serum type IV collagen and prolyl hydroxylase levels: Effect of gender and age
Tamaro G., Moretti M., Marchi P., Mangiarotti M., Pozzato G. 205

Vol. 229/1–2, 1994

The correlation coefficient in the interpretation of laboratory data
Chinchilli V.M., Gruemer H.-D. 1
Early changes of serum N-acetyl-β-glucosaminidase, tissue plasminogen activator and erythrocyte superoxide dismutase in relation to retinopathy in type 1 diabetes mellitus
Skrha J., Hodinar A., Kvasnicka J., Stibor V., Sperl M., Stolba P., Hilgertova J. 5
Nicotinic acid phosphoribosyltransferase activity in human erythrocytes: Studies using a new HPLC method
Pescaglini M., Micheli V., Simmonds H.A., Rocchigiani M., Pompucci G. 15
The utilization of intravenously infused pyridoxine in humans
Zempleni J., Kubler W. 27
Determination of trace amounts of albumin in human bronchoalveolar lavage fluids by fluorometry with chromazurol S
Sato T., Oh'ue M., Nakarai T., Ichikawa K., Kitagishi K., Murakami N., Chikuma M., Saito Y., Kado M., Nagai S., Izumi T. 37
A processing-independent assay for human procholecystokinin and its products
Paloheimo L.I., Rehfeld J.F. 49
Quantitative computer image analysis of a human chorionic gonadotropin colloidal carbon dipstick assay
Van Amerongen A., Van Loon D., Berendsen L.B.J.M., Wichers J.H. 67
Precipitation procedures used to isolate high density lipoprotein with particular reference to effects on apo A-I-only particles and lipoprotein(a)
Vaillance D.T., Byrne D.J., Winder A.F. 77
Differential processing of type I and type III procollagens in the tumour cysts and peritoneal ascitic fluid of patients with benign and malignant ovarian tumours
Zhu G.-G., Melkko J., Risteli J., Kauppila A., Risteli L. 87
Measurement of short-chain acyl-CoA dehydrogenase (SCAD) in cultured skin fibroblasts with hexanoyl-CoA as a competitive inhibitor to eliminate the contribution of medium-chain acyl-CoA dehydrogenase
Niezen-Koning K.E., Wanders R.J.A., Nagel G.T., Sewell A.C., Heymans H.S.A. 99
Estradiol receptors in combination with neu or myc oncogene amplifications might define new subtypes of breast cancer
Bolufer P., Molina R., Ruiz A., Hemandez M., Vazquez C., Lluch A. 107
Quantitative and qualitative changes of extracellular-superoxide dismutase in patients with various diseases
Adachia T., Nakamura M., Yamada H., Futenma A., Kato K., Hirano K. 123
L-Homoserine hydroxamic acid as an antitumor agent
Natelson S., Pantazis P., Natelson E.A. 133

Semi-quantitative latex method for lipoprotein(a) assay
Lippi G., Ruzzenente O., Facchinetti R., Guidi G. 147
Cardiac myosin light chain-I release in acute myocardial infarction is associated with scintigraphic estimates of myocardial scar
Mair J., Wagner I., Fridrich L., Lechleitner P., Diestl F., Puschendorf B., Michel G. 153
A combination assay of MB and MM isoforms of serum creatine kinase in acute myocardial infarction
Kanemitsu F., Okigaki T. 161
Determination of amyloid type by ELISA using milligram amounts of tissue
Kaplan B., German G., Ravid M., Pras M. 171
Simultaneous detection of ΔF508, G542X, N1303K and 1717-1G→A mutations in cystic fibrosis by capillary electrophoresis in polymer networks
Gelfi C., Righetti P.G., Magnani C., Cremonesi L., Ferrari M. 181
Serum hyaluronate levels predict responsiveness to interferon alpha therapy in patients with chronic hepatitis C
Matsunaga T., Kojima J., Inoue A., Matsumiya K. 191
A simple non-radioactive assay for hemoglobin E gene in prenatal diagnosis
Fucharoen S., Fucharoen G., Ratanasiri T., Jetsrisuparb A., Fukumaki Y. 197
Letter to the Editor
Serum serotonin, lactate and pyruvate levels in infantile autostic children
Laszlo A., Horvath E., Eck E., Fekete M. 205

Vol. 230/1, 1994

The detection of 3-methylglutarylcarnitine and a new dicarboxylic conjugate, 3-methylglutaconylcaritine, in 3-methylglutaconic aciduria
Jooste S., Erasmus E., Mienie L.J., De Wet W.J., Gibson K.M. 1
Carbohydrate-mediated recognition of a circulating placental alkaline phosphatase-immunoglobulin M complex
Koyama I., Sato M., Sakai H., Nagata A., Miura M., Sakagishi Y., Komoda T. 9
Determinants of serum 1,25-dihydroxyvitamin D concentration in healthy premenopausal subjects
Ho S.C., Mac Donald D., Chan C., Fan Y.K., Chan S.S.G., Swaminathan R. 21
Vitamin, trace element and peroxide status in HIV seropositive patients: Asymptomatic patients present a severe β-carotene deficiency
Sappey C., Leclercq P., Coudray C., Faure P., Micoud M., Favier A. 35
A screening method for abnormally high lipoprotein(a) concentrations by agarose lipoprotein electrophoresis
Campos E., Fievet P., Caces E., Fruchart J.-C., Fievet C. 43
Copper,zinc-superoxide dismutase and hydrogen peroxide: A hydroxyl radical generating system
Koningsberger J.C., Van Asbeck B.S., Van Faassen E., Wiegman L.J.J.M., Van Hattum J., Van Berge Henegouwen G.P., Marx J.J.M. 51
The use of HPLC to evaluate the variations of blood coronary adenosine levels during percutaneous transluminal angioplasty
Guieu R., Paganelli F., Sampieri F., Bechis G., Levy S., Rochat H. 63
Abnormal glycine betaine content of the blood and urine of diabetic and renal patients
Lever M., Sizeland P.C.B., Bason L.M., Hayman C.M., Robson R.A., Chambers S.T. 69
A competitive dual-label time-resolved immunofluorometric assay for simultaneous detection of carbonic anhydrose I and II in cerebrospinal fluid
Parkkila A.-K., Parkkila S., Serlo W., Reunanen M., Vierjoki T., Rajeniemi H. 81
Urinary excretion of bomocitric acid and methylhomocitric acid in propionic acidaemia: Minor metabolic products of the citrate synthase aldol condensation reaction
Van Rooyen J.P.G., Mienie L.J., Erasmus E., De Wet W.J., Duran M., Wadman S.K. 91
Incidence of the apolopoprotein B-3500 mutation in Denmark
Hansen P.S., Norgaard-Petersen B., Meinertz H., Jensen H.K., Hansen A.B.B., Klausen I.C., Gerdes L.U., Faergeman O. 101

Letter to the Editor
Effect of a 6-month storage on human serum fructosamine concentration [1]
Balland M., Schiele F., Henny J. 105

Vol. 230/2, 1994

Cystic fibrosis and phosphatidylcholine biosynthesis
Ulane M.M., Butler DeB. J., Peri A., Miele L., Ulane R.E., Hubbard V.S. 109
Serum neurone-specific enolase concentrations in patients with neurological disorders
Cunningham R.T., Morrow J.I., Johnston C.F., Buchanan K.D. 117
Radioimmunoassay of vasopressin during pregnancy. Use and removal of cystylaminopeptidase inhibitors
Van der Post J.A., Van Heerikhuize J.J., Boer K., Van Boxtel C.E., Swaab D.F. 125
Binding of aluminium to plasma proteins: Comparative effect of desferrioxamine and deferiprone (L1)
Fernandez-Martin J.L., Menendez-Fraga P., Canteros M.A., Diaz-Lopez J.B., Cannata-Andia J.B. 137
Longitudinal changes in serum zinc concentration and distribution after acute myocardial infarction
Arnaud J., Faure H., Bourlard P., Denis B., Favier A.E. 147
Naftidrofuryl inhibits the release of 5-hydroxytryptamine and platelet-derived growth factor from human platelets
Barradas M.A., Jagroop I.A., Mikhailidis D.P. 157
Simultaneous determination of betaine and N,N-dimethylglycine in urine
Laryea M.D., Zass R., Ritgen J., Wendel U. 169
Development and evaluation of a spectrophotometric assay for complex III in isolated mitochondria, tissues and fibroblasts from rats and humans
Krahenbuhl S., Talos C., Wiesmann U., Hoppel C.L. 177
Heightened free radical activity in blacks with chronic pancreatitis at Johannesburg, South Africa
Gut A., Shiel N., Kay P.M., Segal I., Braganza J.M. 189
A microassay for urinary phenol using capillary gas chromatography and optimised enzymic hydrolysis
Ahmed N., Hale K. 201
Is there a need for creatinine adjustment of urinary steroid hormone levels in studies of early fetal loss?
Hakim R.B., Gray R.H., Zacur H.A. 209
Separation of lipoprotein fractions by ultracentrifugation: Investigation of analytical recovery with sequential flotation and density gradient procedures
Potts J.L., Fisher R.M., Humphreys S.M., Gibbons G.F., Frayn K.N. 215
Letter to the Editor
Sulfadiazine/N^4-acetylsulfadiazine crystalluria in a patient with the acquired immune deficiency syndrome (AIDS)
Potter J.L., Kofron W.G. 221

Vol. 231/1, 1994

Characterization of human creatine kinase BE and MB isoforms by means of isoelectric focusing
Kanemitsu F., Okigaki T. 1
Crystallization of urinary calcium oxalate at standardized osmolality and pH in the frozen state
Gohel M.D.I., Shum D.K.Y., Li M.K. 11
Trace elements in the sweat of acclimatized persons
Omokhodion F.O., Howard J.M. 23
Plasma cholesteryl ester transfer in patients with non-insulin dependent diabetes mellitus
Sutherland W.H., Walker R.J., Lewis-Barned N.J., Pratt H., Tillman H.C. 29

Urinary selenium concentrations in heroin abusers
Rodriquez Rodriquez E.M., Sanz Alaejos M., Diaz Romero C. 39
Different time courses of cardiac contractile proteins after acute myocardial infarction
Mair J., Wagner I., Jakob G., Lechleitner P., Dienstl F., Puschendorf B., Michel G. 47
Regulation of the mitochondrial ATP-synthase in human fibroblasts
Das A.M., Byrd D.J., Brodehl J. 61
Excess matrix accumulation in scleroderma is caused partly by differential regulation of stromelysin and TIMP-1 synthesis
Bou-Gharios G., Osman J., Black C., Olsen I. 69
A one-step sandwich enzyme immunoassay for inactive precursor and complexed forms of human matrix metalloproteinase 9 (92 kDa gelatinase/type IV collagenase, gelatinase B) using monoclonal antibodies
Fujimoto N., Hosokawa N., Iwata K., Shinya T., Okada Y., Hayakawa T. 79
Effect of storage at -70°C on immunoturbidimetric assays for apolipoproteins A-I and B
Myron Johnson A., Ledue T.B., Palomaki G.E., Craig W.Y. 89
Red blood cell glutathione status in patients with gastrointestinal malignancies treated with 5-fluorouracil
Walach N., Evans S., Zaidman J.L., Cotariu D. 95
Increased angiotensin-eonverting enzyme activity in cerebrospinal fluid of treated patients with Parkinson's disease
Konings C.H., Kuiper M.A., Bergmans P.L.M., Grijpma A.M., van Kamp G.J., Wolters Ch. E. 101
Measurement of steroid hormones in plasma by isocratic high performance liquid chromatography coupled to radioimmunoassay
Boschi S., De Iasio R., Mesini P., Bolelli G.F., Sciajno R., Pasquali R., Capelli M. 107
Letter to the Editor
Usefulness of the kappa/lambda ratio for detecting a new monoclonal component in a patient with a monoclonal gammopathy
Myara I., Asso M., Tenenhaus D., Moatti N. 115

Vol. 231/2, 1994

An improved bromide assay for the estimation of extracellular water volume by capillary gas chromatography autovan
Kreel B.K. 117
Routine high-performance liquid chromatographic determination of urinary unconjugated cortisol using solid-phase extraction and ultraviolet detection
Dolezalova M. 129
Immunoheterogeneity of parathyroid hormone after surgery for primary hyperparathyriodism
Bergenfelz A., Valdermarsson S., Ahren B. 139
Competitive enzyme-linked immunosorbent assay of the human cholesteryl ester transfer protein (CETP)
Guyard-Dangremont V., Lagrost L., Gambert P., Lallemant C. 147
Hemoglobins S and C: Reference values for glycohemoglobin in heterozygous, double-heterozygous and homozygous subjects, as established by 13 methods
Weykamp C.W., Martina W.V., Van der Dijs F.P.L., Penders T.J., Van der Slik W., Muskiet F.A.J. 161
Gamma-type gliadins cause secretion of prostaglandin E_2 in patients with coeliac disease
Friis S., Anthonsen D., Noren O., Sjostrom H. 173
Evaluation of enzyme inhibitors of cystinyl aminopeptidase and application to the measurement of immunoreactive atrial natriuretic peptide in human pregnancy
Irons D.W., Davison J.M., Baylis P.H. 185
Improved method for quantitative analysis of lung surfactant phospholipids in bronchoalveolar lavage fluids by high-performance liquid chromatography
Bruch J., Gono E., Malkusch W., Rehn B. 193

Vol. 232/1–2, 1994

Clinical chemistry as scientific discipline: Historical perspectives
Buttner J. 1
Training and continuous education of clinical laboratory technologists and technicians
Mocarelli P. 11
The impact of clinical biochemistry on the pre-doctoral medical curriculum: An Italian viewpoint
Burlina A. 23
Post-doctoral training of the non-physician doctoral scientist
Schwartz M.K. 33
Teaching and training of clinical biochemists: Sub-specialization
Rock R.C. 39
Continuing clinical chemistry education in the United States
Conn R.B. 47
Clinical chemistry education in the United States
Tetrault G.A., Gruemer H.-D. 53
Education in clinical biochemistry: The Canadian scene
Bunting P.S., Goldberg D.M. 63
Training and education in clinical biochemistry in the United Kingdom
Price C.P., Dryburgh F.J., Elder G.H. 85
Education and clinical biochemistry in Australia
O'Leary T.D., Guerin M.D. 97
Education in clinical chemistry in Germany
Delbruck A.F. 109
Education in clinical chemistry and laboratory sciences in Austria
Muller M.M. 119
Teaching clinical chemistry in central European countries — past and present
Naskalski J.W., Palicka V. 125
Clinical biochemistry training in the Netherlands
Sanders G.T.B., Brombacher P.J. 133
The impact of clinical biochemistry on university education in France
Masseyeff R.F., Dreux C., Goussault Y. 143
Clinical biochemistry education in Spain
Queralto J.M. 153
Education in clinical chemistry in Japan
Kanno T. 159
Training of clinical laboratory professionals in Russia
Morozova V.T., Dolgov V.V., Malakhov V.N. 167

Vol. 233/1–2, 1995

Appropriate utilization and cost control of the hospital laboratory: Panel testing and repeat orders
Werner M. 1
Multi-enzyme reference material from established human cell lines and human sources
Marui Y., Hayashi C., Matsuda Y., Matsuura S., Eto A., Ohba Y., Okuda K. 19
Biotinylation of biotinidase following incubation with biocytin
Hymes J., Fleischauer K., Wolf B. 39
The influence of the blood levels of lead, aluminum and vanadium upon the arterial hypertension
Granadillo V.A., Tahan J.E., Salgado O., Elejalde L.E., Rodriguez-Iturbe B., Romero G.B., Romero R.A. 47
Role of monocyte/macrophage derived matrix-metalloproteinases (gelatinases) in prolonged skin inflammation
Maillard J.L., Favreau C., Reboud-Ravaux M. 61
Hungarian hereditary acatalasemia and hypocatalasemia are not associated with chronic hemolysis
Goth L., Vitai M. 75

Dimethylamine in human urine
Zhang A.Q., Mitchell S.C., Smith R.L. 81
A preliminary report on urinary BT-PABA/PAS excretion index, serum pancreatic isoamylase and faecal chymotrypsin tests of pancreatic dysfunction in Sowetan Africans
Chaloner C., Segal I., Hassan H., McIntosh J., Gut A., Rahme M., Braganza J.M. 89
Estimation of the value of serum β_2-microglobulin concentration in the diagnosis of Hashimoto' disease
Kochanska-Dziurowicz A., Starzewski J.J., Dunal W. 101
Electrolyte balanced heparin may produce a bias in the measurement of ionized calcium concentration in specimens with abnormally low protein concentration
Lyon M.E., Guajardo M., Laha T., Malik S., Henderson P.J., Kenny M.A. 105
Transient hyperphosphatasaemia in infancy associated with an increased urinary hydroxyproline excretion
Stepan J.J., Kutilek S., Bayer M. 115
Cholestyramine orally administered to patients with erythropoietic protoporphyria results in urinary excretion of protoporphyrin: To be explained by the Herbst-Volkheimer effect?
Beukeveld G.J.J., Wolthers B.G. 119
Sialyl salivary-type amylasemia associated with immunoglobulin D-type multiple myeloma
Moriyama T., Tozawa T., Nobuoka M., Ikeda H. 127

Vol. 234/1–2, 1995

Detection of the three common mutations of adeninephosphoribosyltransferase deficiency among Japanese
Higashimoto H., Ouchi A., Kawaguchi R. 1
Differentiation between organophosphate and carbamate poisoning
Rotenberg M., Shefi M., Dany S., Dore I., Tirosh M., Almog S. 11
Tissue accumulation of sulfatide and G(M3) ganglioside in a patient with variant Farber disease
Fujiwaki T., Hamanaka S., Tate S., Inagaki F., Suzuki M., Suzuki A., Mori C. 23
Immunochemical detection of urinary 5-methyl-2′-deoxycytidine as a potential biologic marker for leukemia
Itoh K., Aida S., Ishiwata S., Yamagchi T., Ishida N., Mizugaki M. 37
Reactivity of lecithin-cholesterol acyl transferase (LCAT) towards glycated high-density lipoproteins (HDL)
Fournier N., Myara I., Atger V., Moatti N. 47
Measurement of serum lipid peroxidation during exercise using three different methods: Diene conjugation, thiobarbituric acid reactive material and fluorescent chromolipids
Vasankari T., Kujala U., Heinonen O., Kapanen J., Ahotupa M. 63
An HPLC assay for sn-1,2-diacylglycerol
Eaton S., Shmueli E., Al-Mardini H., Bartlett K., Record C.O. 71
An improved method for quantification of extra domain A-containing cellular fibronectin (EDAcFN) in different body fluids
Ylatupa S., Mertaniemi P., Haglund C., Partanen P. 79
Comparison of erythrocyte transketolase activity with thiamine and thiamine phosphate ester levels in chronic alcoholic patients
Herve C., Beyne P., Letteron Ph., Delacoux E. 91
Galactosyl hydroxylysine in assessment of Paget's bone disease
Michalsky M., Stepan J.J., Wilczek H., Formankova J., Moro L. 101
Evaluation of oxidant stress in chronic hemodialysis patients: Use of different parameters
Haklar G., Yegenaga I., Yalcin A.S. 109
One-step sandwich enzyme immunoassays for human C4b-binding protein (C4BP) and protein S-C4BP complex using monoclonal antibodies
Tamei H., Hoshino T., Yoshida S., Hayashi T., Iwata K., Suzuki K. 115
Redox state, antioxidative activity and lipid peroxidation in erythrocytes and plasma of chronic ambulatory peritoneal dialysis patients

Canestrari F., Buoncristiani U., Galli F., Giorgini A., Albertini M.C., Carobi C., Pascucci M., Bossu M. 127

Effect of double-blind cross-over selenium supplementation on lipid peroxidation markers in cystic fibrosis patients
Portal B., Richard M.-J., Coudray C., Arnaud J., Favier A. 137

Quantitative determination of faecal fat, nitrogen and water by means of a spectrophotometric technique: Near infrared reflectance analysis (NIRA). Assessment of its accuracy and reproducibility compared with chemical methods
Picarelli A., Greco M., Di Giovambattista F., Ramazzotti A., Cedrone C., Corazziari E., Torsoli A. 147

Ion-selective electrode for serum bromide assay in patients with epilepsy
Katsu T., Furuno K., Yamashita S., Kawasaki H., Gomita Y., Ohtsuka Y., Ohtahara S. 157

Total renin in pre-eclampsia
Matinlauri I., Ekblad U., Maenpaa K., Koskinen P., Aalto M., Peltola O., Irjala K. 163

Clinical usefulness of cimetidine treatment for acute relapse in intermittent porphyria
Horie Y., Norimoto M., Tajima F., Sasaki H., Nanba E., Kawasaki H. 171

Letter to the Editor

Determination of α-amylase using a new blocked substrate (2-chloro-4-nitrophenyl 4^4-O-β-D-galactopyranosyl-β-maltotetraoside)
Majima K., Teshima S., Hamada Y., Kikuchi T., Kawamura Y., Kitahata S. 177

Vol. 235/1, 1995

A scheme for the interpretation of primary and secondary disturbances of plasma and urinary amino acid profiles. A possible way to an expert system
Parvy P., Bardet J., Rabier D., Kamoun P. 1

Active and acid-activatable TGF-β in human sera, platelets and plasma
Grainger D.J., Mosedale D.E., Metcalfe J.C., Weissberg P.L., Kemp P.R. 11

Serum pyridinolines as specific markers of bone resorption in hemodialyzed patients
Niwa T., Shiobara K., Hamada T., Miyazaki T., Tsukushi S., Uema K., Tsuzuki T. 33

Genetic basis of the silent phenotype of serum butyrylcholinesterase in three compound heterozygotes
Maekawa M., Sudo K., Kanno T., Kotani K., Dey D.C., Ishikawa J., Izumi M., Etoh K. 41

Immunoturbidimetric determination of lipoprotein(a): Improvement in the measurement of turbid and triglyceride-rich samples
Molinari E.A., Pichler P.F., Grillhofer H., Kostner G.M. 59

Alpha-fetoprotein and the acute phase response. A study using acute pelvic inflammatory disease as a model system
Christiansen M., Hogdall C.K., Brihmer C. 71

A simple and automated HPLC method for determination of total hydroxyproline in urine. Comparison with excretion of pyridinolines
Castelain S., Kamel S., Picard C., Desmet G., Sebert J.L., Brazier M. 81

Comparison of micro-enzymatic and high-performance liquid chromatographic methods for the assay of serum 1,5-anhydroglucitol
Chusney G.D., Philippa M., Pickup J.C. 91

CSF acetylcholinesterase in Parkinson disease: Decreased enzyme activity and immunoreactivity in demented patients
Konings C.H., Kuiper M.A., Mulder C., Calliauw J., Wolters Ch. E. 101

Cholinesterase activities in cerebrospinal fluid of patients with idiopathic convulsive disorders
Basu P.S., Batabyal S.K., Bhattacharya A., Datta T.K. 107

Vol. 235/2, 1995

Radiochemical assay of minute quantities of galactose-1-phosphate uridyltransferase activity in

erythrocytes and leukocytes of galactosemia patients
Xu Y.-K., Kaufman F.R., Donnell G.N., Ng W.G. 125
A sandwich enzyme immunoassay for the determination of neutrophil lipocalin in body fluids
Blaser J., Triebel S., Tschesche H. 137
Blood antioxidants (vitamin E and β-carotene) in long-term low density lipoprotein apheresis
Assogba U., Lepage S., Bruckert E., Bonnefont-Rousselot D., Dairou F., De Gennes J.L., Delattre J. 147
Changed transferrin sialylation in Parkinson's disease
Van Kamp G.J., Mulder K., Kuiper M., Wolters Ch. E. 159
Enzymatic determination of bicarbonate: In serum by flow injection analysis
Quiles R., Fernandez-Romero J.M., De Castro M.D.L. 169
Oxidative stress and erythrocyte membrane fluidity in patients undergoing regular dialysis
McGrath L.T., Douglas A.F., McClean E., Brown J.H., Doherty C.C., Johnston G.D., Archbold G.P.R. 179
Serum lipoprotein profile in Algerian patients with celiac disease
Mediene S., Hakem S., Bard J.M., Medjaoui I., Benhamamouch S., Lebel P., Fruchart J.C., Clavey V. 189
A specific immunoradiometric assay of cationic trypsin(ogen) that does not recognize trypsin-alpha-1-proteinase inhibitor complex
Lafont P., Guy-Crotte O., Paulin C., Galvain D., Mertani S., Figarella C., Barbier Y. 197
The red wine phenolics trans-resveratrol and quercetin block human platelet aggregation and eicosanoid synthesis: Implications for protection against coronary heart disease
Pace-Asciaka C.R., Hahn S., Diamandis E.P., Soleas G., Goldberg D.M. 207
Assay of serum pyridinoline: A potential marker for bone resorption
Hata K., Miura M., Fukumoto S., Matsumoto T. 221
Surfactant associated protein A determination using a chemiluminescence system — Application to tracheal aspirates from newborns
Taieb J., Francoual J., Magny J.-F., Fraslon d C., Messaoudi C., Lindenbaum A., Bourbon J. 229

Vol. 236/1, 1995

Calcium and magnesium in human toenails do not reflect bone mineral density
Vecht-Hart C.M., Bode P., Trouerbach Th. W., Collette H.J.A. 1
Human triacylglycerol-rich lipoprotein subfractions as substrates for lipoprotein lipase
Fisher R.M., Coppack S.W., Humphrey S.M., Gibbons G.F., Frayn K.N. 7
Urinary creatinine adjusted reference ranges for homovanillic and vanillylmandelic acid in children and adults
Cole M., Craft A.W., Parker L., Bell S., Seviour J.A., McGill A.C., Dale G. 19
Urinary steroid profile of a newborn suffering from pseudahypoaldosteronism
Wolthers B.G., Kraan G.P.B., Van der Molen J.C., Nagel G.T., Rouwe C.W., Lenting F. 33
Diagnosis of Smith-Lemli-Opitz syndrome by gas chromatography/mass spectrometry of 7-dehydrocholesterol in plasma, amniotic fluid and cultured skin fibroblasts
Kelley R.I. 45
A sandwich ELISA based on anti-apo(a) and anti-apo B monoclonal antibodies for lipoprotein(a) measurement
Sorell L., Rojas G., Rodgriguez M., Ramos C., Torres L., Torres M.B. 59
Clinical evaluation of a monoclonal antibody to serum KM01 for the diagnosis of hepatocellular carcinoma
Fujii T., Horie Y., Ikuta Y., Nishimuki E., Murawaki Y., Suou T., Kawasaki H. 71
Genetic analysis of Japanese patients with myophosphorylase deficiency (McArdle's disease): Single-codon deletion in exon 17 is the predominant mutation
Sugie H., Sugie Y., Ito M., Fukuda T., Nonaka I., Igarashi Y. 81
Within- and between-subject variation in, and associations between, serum concentrations and urinary excretion of testosterone and estradiol in Chinese men
Ooi L.S.M., Panesar N.S., Masarei J.R.L. 87

Enzyme-linked immunosorbent assay for. Extracellular glutathione peroxidase in serum of normal individuals and patients with renal failure on hemodialysis
Yamamoto Y., Takekoshi Y., Itami N., Honjo T., Kojima H., Yano S., Takahashi H., Saito I., Takahashi K. 93
Neutrophil activation and collagen metabolism in lung cancer
Gabazza E.C., Taguchi O., Yoshida M., Yamakami T., Kobayashi H., Ibata H., Shima T. 101
Erratum: The correlation in the interpretation of laboratory data (Clinica Chimica Acta 229 (1994) (1–3))
Chinchilli V.M., Gruemer H.-D. 113

Vol. 236/2, 1995

The value of polymorphonuclear elastase in adult respiratory distress syndrome
Diaz J., Tornel P.L., Jara P., Canizares F., Egea J.M., Martinez P. 119
Polymorphonuclear leukocyte elastase levels in patients with Behcet's disease
Deger O., Orem A., Akyol N., Bahadir S., Yildirmis S. 129
Amino acid sequence of a modified β_2-microglobulin in renal failure patient urine and long-term dialysis patient blood
Momoi T., Suzuki M., Titani K., Hisanaga S., Ogawa H., Saito A. 135
Evidence of toxic metabolite stress in black South Africans with chronic pancreatitis
Gut A., Chaloner C., Schofield D., Sandle L.R., Purmasir M., Segal I., Braganza J.M. 145
Immunoglobulin cleavage by hypochlorous acid treatment
Drozdz R., Guevara I., Naskalski J.W. 155
Decreased branching, increased fucosylation and changed sialylation of alpha-1-proteinase inhibitor in breast and ovarian cancer
Goodarzi M.T., Turner G.A. 161
Selenium dependent glutathione peroxidase activity in human follicular fluid
Paszkowski T., Traub A.I., Robinson S.Y., McMaster D. 173
The typing of fetal antigen 2 in human amniotic fluid
Price K.M., Silman R., Armstrong P., Teisner B., Grudzinskas J.G. 181
Demonstration of immunogenic keratan Sulphate in commercial chondroitin 6-sulphate from shark cartilage. Implications for ELISA assays
Moller H.J., Moller-Pedersen T., Damsgaard T.E., Poulsen J.H. 195
Maximization of determination of basement membrane collagen complex in the sera from patients with chronic liver diseases by a sandwich ELISA
Amano S., Miyake K., Shiba K., Yoshida T., Hayashi T. 205

Vol. 237/1–2, 1995

Novel enzymes as reagents
Price C.P., Campbell R.S., Hammond P.M. 3
The effect of temperature on enzymes used in diagnostics
Scopes R.K. 17
^{13}C NMR investigation of cholesterol esterification rate in human whole blood
Nouri-Sorkhabi M.H., Sullivan D.R., Kuchel P.W. 25
Ligand western blotting for specific detection of active forms of proteases
Kido T., Yook H.-Y., Ueda K. 31
Recombinant M-, B- and MB-type isozymes of human phosphoglyceric acid mutase: Their large-scale production and preparation of polyclonal antibodies specific to M- and B-type isozymes
Uchida K., Kondoh K., Matuo Y. 43

Cardiac troponin, CK-MB and myoglobin for the early detection of acute myocardial infarction and monitoring of reperfusion following thrombolytic therapy
Apple F.S., Voss E., Lund L., Preese L., Berger C.R., Henry T.D. 59
Tissue cathepsins as tumor markers
Schwartz M.K. 67
Clinical applications of the p53 tumor suppressor gene
Diamandis E.P. 79
Clinical and analytical considerations in the utilization of cholinesterase measurements
McQueen M.J. 91
Nephrotoxicity of xenobiotics
Werner M., Costa M.J., Mitchell L.G., Nayar R. 107
Beyond alcohol: Beverage consumption and cardiovascular mortality
Goldberg D.M., Hahn S.E., Parkes J.G. 155

Vol. 238/1, 1995

Detection of autoantibodies against oxidized low-density lipoproteins and of IgG-bound low density lipoproteins in patients with coronary artery disease
Boullier A., Hamon M., Walters-Laporte E., Martin-Nizart F., Mackereel R., Fruchart J.-C., Bertrand M., Duriez P. 1
Whether, to what degree, and why lipoprotein(a) levels change over time
Glueck C.J., Tracy T., Sieve-Smith L., Wang P. 11
Quantitation of procholecystokinin and its products in plasma by processing-independent analysis
Paloheimo L.I., Rehfeld J.F. 21
Glycaemia and body mass as determinants of plasma lecithin: Cholesterol acyltransferase activity in Nigerian patients with non-insulin dependent diabetes mellitus
Akanji A.O., Agbedana E.O. 35
Compositions of very low density lipoprotein subfractions from patients with polydisperse low density lipoproteins
Saheki S., Hitsumoto Y., Bando S., Murase M., Takeuchi N., Uchida K. 43
Relationships between triglycerides, lipoproteins, glucose and coronary artery disease
Hobbs G.A., Wagner S.G., Levinson S.S. 59
Study of the chromium binding in plasma of patients on continuous ambulatory peritoneal dialysis
Borguet F., Cornelis R., Delanghe J., Lambert M.-C., Lameire N. 71
Serum concentrations of the pyridinoline crosslinked carboxy-terminal telopeptide of type I collagen in chronic liver disease
Hayasaka A., Iida S., Ohto M. 85
Low serum alkaline phosphatase activity associated with severe Wilson's disease. Is the breakdown of alkaline phosphatase molecules caused by reactive oxygen species?
Hoshino T., Kumasaka K., Kawano K., Yamagishi F., Koyama I., Fujimori-Arai Y., Nakajima T., Komoda T. 91
Hyaluronic acid and dermatan sulfate in nonlesional pseudoxanthoma elasticum skin
Lebwohl M., Longas M.O., Konstadt J., Koreen R., Grabowski G.A., Phelps R., Lemlich G., Schwartz E. 101

Vol. 238/2, 1995

Identification of the D-enantiomer of 2-hydroxyglutaric acid in glutaric aciduria type II
Watanabe H., Yamaguchi S., Saiki K., Shimizu N., Fukao T., Kondo N., Orii T. 115
Effect of neurotoxic metal ions in vitro on proteolytic enzyme acivities in human cerebral cortex
Falkous G., Harris J.B., Mantle D. 125
N-Methyladrenaline: Age-dependent urinary excretion, perinatal organ content and relation with 'classical' catecholamines
Kema I.P., Schoots C.J.F., Gidding C.E.M., Okken A., Aarnoudse J.G., Muskiet F.A.J. 137

Maternal serum screening for fetal Down's syndrome, a retrospective study
Lemay C., Roussel-Mizon N., Thepot F., Desmet G. 151
Variegate porphyria: Diagnostic value of fluorometric scanning of plasma porphyrins
Da Silva V., Simonin S., Deybach J.C., Puy H., Nordmann Y. 163
Demonstration of induction of erythrocyte inosine monophosphate dehydrogenase activity in Ribavirin-treated patients using a high performance liquid chromatography linked method
Montero C., Duley J.A., Fairbanks L.D., McBride M.B., Micheli V., Cant A.J., Morgan G. 169
Serotonin content in human blood platelets is not stable
Karas S.I., Yazykov K.G., Makarova E.V. 179
Actions of arachidonic acid on erythrocyte membrane Rb permeability
Dwight St. J.F.J., Hendry B.M. 187
Fasting plasma glucose as a diagnostic indicator of diabetes mellitus
Wiener K. 199
Rapid degradation of 3,4-dihydroxyphenylalanine in plasma samples containing ethylenediaminetetraacetic acid
Eriksson T., Holmgren B. 209
Erratum: Primary focal segmental glomerulosclerosis is associated with increased intraglomemlar type IV collagen synthesis (Clinica Chimica Acta 235 (1995) (121–124))
Razzaque M.S., Koji T., Harada T., Nakane P., Taguchi T. 217

Vol. 239/1, 1995

Effects of urinary macromolecules on the nucleation of calcium oxalate in idiopathic stone formers and healthy controls
Borghi L., Meschi T., Guerra A., Bergamaschi E., Mutti A., Novarini A. 1
Free fatty acid analysis in ascitic fluid improves diagnosis in malignant abdominal tumors
Greco A.V., Mingrone G., Gasbarrini G. 13
Components of biological variation in plasma haptoglobin: Relationships to plasma fibrinogen and immune variables, including interleukin-6 and its receptor
Maes M., Cooreman W., Delanghe J., Scharpe S., Wauters A., Neels H., D'Hondt P., Peeters D., Cosyns P., Ranjan R., Bosmans E., DeMeyer F., Elkis H., Desnyder R. 23
A method for the separation of delta bilirubin using Cibacron Blue affinity chromatography
Burnett J.R., Lim C.W., Mahoney G.N., Crooke M.J. 37
Interaction of human pregnancy-associated plasma protein-A with serine proteinases
Zorin N.A., Zhabin S.G., Semenkov N.N. 47
Sensitivity and specificity of a new ELISA method for determination of chromogranin A in the diagnosis of pheochromocytoma and neuroblastoma
Boomsma F., Bhaggoe U.M., Man in 't Veld A.J., Schalekamp M.A.D.H. 57
Determination of ammonia in ear-lobe capillary blood is an alternative to arterial blood ammonia
Huizenga J.R., Gips C.H., Conn H.O., Jansen P.L.M. 65
Micronutrient antioxidant status in black South Africans with chronic pancreatitis: Opportunity for prophylaxis
Segal I., Gut A., Schofield D., Shiel N., Braganza J.M. 71
Plasma non-esterified fatty acid response to dialysate acetate concentration during haemodialysis in humans
Jackson J.M., Talbot S.T., Lee H.A. 81
Quantitative cytochemistry of human leukocyte elastase compared with plasma elastase and acute phase proteins in inflammatory diseases
Pradella M., Nemetz L., Bovo C., Visentin M., Clemen P., Zappala G., Ossani M., Plebani M. 91

Vol. 239/2, 1995

Daily variation of serum lipids in relation to the circadian rhythm of platelet aggregation in healthy male persons
Knofler R., Urano T., Taminato T., Yoshimi T., Nakano T., Nakajima K., Takada Y., Takada A. 109

New aspects concerning the regulation of the postoperative acute phase reaction during cardiac surgery
Berger D., Bolke E., Huegel H., Seidelmann M., Hannekum A., Begerb H.G. 121
Enhanced susceptibility of low-density lipoprotein to in vitro oxidation in type 1 and type 2 diabetic patients
Beaudeux J.-L., Guillausseau P.-J., Peynet J., Flourie F., Assayag M., Tielmans D., Warnet A., Rousselet F. 131
Comparison between serum levels of bone alkaline phosphatase and the carboxy-terminal propeptide of type I procollagen as markers of bone formation in patients following renal transplantation
Withold W., Degenhardt S., Grabensee B., Reinauer H. 143
Bi-enzyme reactor for electrochemical detection of low concentrations of uric acid and glucose
Elekes O., Moscone D., Venema K., Korf J. 153
Solid-phase competitive luminescence immunoassay for lysozyme in faeces
Gao P., John M.R., Schmidt-Gayk H., Arndt B., Scheida M., Theuer D. 167
Osteocalcin metabolism in late fetal life: Fetal and maternal osteocalcin levels
Alatasa O., Colak O., Alatas E., Tekin B., Inal M., Ozalp S. 179
The effect of ascorbic acid on the measurement of total cholesterol and triglycerides: Possible artefactual lowering in individuals with high plasma concentration of ascorbic acid
Benzie I.F.F., Strain J.J. 185
Phenotyping of glutathione S-transferase m1 in the estonian population by ELISA using GSTM1a and GSTM1b specific monoclonal antibodies
Tasa G., Juronen E., Uuskula M., Viikmaa M., Mikelsaar A.-V. 191
Creatine kinase isoenzymes in human trichinellosis
Tassia C., Materazzi L., Pozio E., Bruschi F. 197

Vol. 240/1, 1995

Enzyme-linked immunosorbent assay for cholesteryl ester transfer protein in human serum
Sato T., Fukasawa M., Kinoshita M., Arai H., Saeki T., Naraki T., Iwasaki Y., Teramoto T., Takahashi K., Saito Y., Inoue K. 1
New transcutaneous sampling of glucose for patients with type II diabetes using an ion-sensitive field-effect transistor
Kayashima S., Arai T., Noritake M., Nagata N., Kikuchi M., Ito N., Matsumoto Y., Kaneyoshi A., Kimura J., Kuriyama T. 11
HPLC analysis of uridine diphosphate sugars: Decreased concentrations of uridine diphosphate galactose in erythrocytes and cultured skin fibroblasts from classical galactosemia patients
Xu Y.-K., Kaufman F.R., Donnell G.N., Giudici T., Alfi O., Ng W.G. 21
3-Hydroxyisovalerylcarnitine in patients with deficiency of 3-methylcrotonyl CoA carboxylase
Roschinger W., Millington D.S., Gage D.A., Huang Z.-H., Iwamoto T., Yano S., Packman S., Johnston K., Berry S.A., Sweetman L. 35
Susceptibility to ulcerative colitis and Crohn's disease: Interactions between glutathione S-transferase GSTM1 and GSTT1 genotypes
Duncan H., Swan C., Green J., Jones P., Brannigan K., Alldersea J., Fryer A.A., Strange R.C. 53
The total apolipoprotein B/LDL-cholesterol ratio does not predict LDL particle size
Tallis G.A., Shephard M.D.S., Sobecki S., Whiting M.J. 63
Characterization of commercial antibodies for use in high resolution apo(a) phenotyping by immunoblot analysis
Gaw A., Chiesa G. 75
Screening urine protein abnormalities by a novel test paper
Shiba K., Kanamori K., Sugiura K. 83
Changes in red blood cell glutathione and glutathione-dependent enzymes on long-term treatment with captopril and enalapril
Golik A., Weissgarten J., Evans S., Cohen N., Zaidenstein R., Modal D., Cotariu D. 89
Letters to the Editor
The stability of pyridinium crosslinks in urine and serum
Walne A.J., James I.T., Perrett D. 95

No evidence for abnormalities in kinetics of platelet monoamine oxidase in Alzheimer's disease
Konings C.H., Scheltens P., Kuiper M.A., Wolters E.C. 99

Vol. 240/2, 1995

Evaluation of the decarbamylation process of cholinesterase during assay of enzyme activity
Rotenberg M., Almog S. 107

Effect of encapsulation on the anti-inflammatory properties of superoxide dismutase after oral administration
Regnault C., Roch-Arveiller M., Tissot M., Sarfati G., Giroud J.P., Postaire E., Hazebroucq G. 117

An improved spectrophotometric assay of pyruvate dehydrogenase in lactate dehydrogenase contaminated mitochondrial preparations from human skeletal muscle
Chretien D., Pourrier M., Bourgeron T., Sene M., Rotig A., Munnich A., Rustin P. 129

Development of an enzyme-linked immunosorbent assay to measure total TIMP-1 (free TIMP-1 and TIMP-1 in combination with matrix-metalloproteinases) and measurement of TIMP 1 and CRP in serum
Plumpton T.A., Clark I.M., Plumpton C., Calvin J., Cawston T.E. 137

Comparison of 5-hydroxyindole-acetic acid and para-amino hippurate clearances in newborn rabbits
Sulyok E., Nagy L., Baranyai Z., Thonney M., Guignard J.-P. 155

Plasma protein leakage and local secretion of proteins assessed in sputum in asthma and COPD. The effect of inhaled corticosteroids
Schoonbrood D.F.M., Out T.A., Lutter R., Reimert C.M., Van Overveld F.J., Jansen H.M. 163

Elastase binding capacity of α_2-macroglobulin and its association with glucocorticoid concentration in southern African black patients with hepatocellular carcinoma
Gaillard M.C., Song E., Nogueira C.M., Kilroe-Smith T.A. 179

Separation of T and B lymphocytes from human peripheral blood mononuclear cells using density perturbation methods
Patel D., Rubbi C.P., Rickwood D. 187

An improved GLC method for a rapid, simultaneous analysis of both medium chain fatty acids and medium chain triglycerides in plasma
Mingrone G., Greco A.V., Capristo E., Benedetti G., Castagneto M., Gasbarrini G. 195

Preferential depletion of selenoprotein P in hypercholesterolaemic patients treated by LDL-apheresis
Persson-Moschos M., Bonnefont-Rousselot D., Assogba U., Bruckert E., Jaudon M.C., Delattre J., Akesson B. 209

Tartrate resistant acid phosphatase in the serum of patients with larynx carcinoma — A marker of osteoclast activation
Jarkowska-Kozlowska J., Kozlowski Z. 213

Vols. 241–242, 1995

Clinical mass spectrometry
Niwa T. 3

Vol. 243/1, 1995

Enzyme-linked immunosorbent assay (ELISA) for Aleuria aurantia lectin-reactive serum cholinesterase to differentiate liver cirrhosis and chronic hepatitis
Kondo M., Hada T., Fukui K., Iwasaki A., Higashino K., Yasukawa K. 1

Urinary C4 excretion in systemic lupus erythematosus
Ueda Y., Nagasawa K., Tsukamoto H., Horiuchi T., Yoshizawa S., Tsuru T., Furugo I., Niho Y. 11

Non-specific increase of serum carbohydrate antigen 19-9 in patients with liver disease associated with increased circulating levels of adhesion molecules
Fabris C., Falleti E., Pirisi M., Soardo G., Toniutto P., Vitulli D., Bortolotti N., Gonano F., Bartoli E. 25

A trinucleotide deletion together with a base duplication event at codon 439 in the human tyrosinase gene identifies a mutational hotspot
Breimer L.H., Winder A.F., Panayiotidis P., Jay M., Moore A., Jay B. 35

Direct double antibody sandwich immunoassay of mucin M_1 epitopes in human mucus secreting pancreatic cell lines
Montserrat C., Hollande E., Guy-Crotte O., Figarella C. 43

Type 1 collagen synthesis by skin fibroblasts from 17 patients with osteogenesis imperfecta type III
Pocock A.E., Francis M.J.O., Smith R. 53

Effects of membrane incorporation of short-chain phospholipids on sodium pump function in human erythrocytes
Dwight J.F.S.J., Hendry B.M. 73

Plasma carnitine concentrations in pregnancy, cord blood, and neonates and children
Winter S.C., Linn L.S., Helton E. 87

Intra-individual variation of overnight urinary albumin excretion in clinically healthy individual middle-aged individuals
Jensen J.S. 95

Erratum: Nephrotoxicity of xenobiotics (Clinica Chimica Acta 237 (1995) 107–154)
Werner M., Costa M.J., Mitchell L.G., Nayar R. 101

Vol. 243/2, 1995

Aminooxy acetic acid: A selective inhibitor of alanine:glyoxylate aminotransferase and its use in the diagnosis of primary hyperoxaluria type I
Horvath V.A.P., Wanders R.J.A. 105

Acetylcholinesterase of human intestinal tissue affected by Hirschsprung's disease: Effect of magnesium chloride on isoforms
Johnson G., Moore S.W., Purves L.R. 115

Measurement of mevalonic acid in human urine by bench top gas chromatography-mass spectrometry
Siavoshian S., Simoneau C., Maugeais P., Marks L., Rodary L., Gardette J., Krempf M. 129

Enhanced susceptibility of low-density lipoproteins to oxidation in coronary bypass patients with progression of atherosclerosis
De Rijke Y.B., Verwey H.F., Vogelezang C.J.M., Van Der Velde E.A., Princen H.M.G., Van Der Laarse A., Bruschke A.V.G., van Berkel T.J.C. 137

Purine and carnitine metabolism in muscle of patients with Duchenne muscular dystrophy
Camina F., Novo-Rodriguez M.I., Rodriguez-Segade S., Castro-Gago M. 151

Determination of serum total lipid and free N-acetylneuraminic acid in genitourinary malignancies by fluorimetric high performance liquid chromatography. Relevance of free N-acetylneuraminic acid as tumour marker
Lagana A., Pardo-Martinez B., Marino A., Fago G., Bizzarri M. 165

Characterization of mitochondrial DNA in primary cardiomyopathies
Bobba A., Giannattasio S., Pucci A., Lippolis R., Camaschella C., Marra E. 181

Evaluation of antibody- and antigen-coated enzymeimmunoassays for measuring oestrone-3-glucuronide concentrations in urine
Henderson K.M., Camberis M., Hardie A.H. 191

Does plasma neuropeptide-Y immunoreactivity in patients with pheochromocytoma depend on hormonal activity of the tumor?
Pruszczyk P., Wocial B., Ignatowska-Switalska H., Feltynowski T., Ellafi M., Januszewicz A., Lapinski M., Zukowska-Grojec Z., Januszewicz W. 205

Vol. 244/1, 1996

Characterization of lipoproteins during human cholestasis
Tallet F., Vasson M.-P., Couderc R., Lefevre G., Raichvarg D. 1

Determination of metalloproteinases, plasminogen-activators and their inhibitors in the synovial fluids of patients with rheumatoid arthritis during chemical synoviorthesis
Blaser J., Triebel S., Maasjosthusmann U., Romisch J., Krahl-Mateblowski U., Freudenberg W., Fricke R., Tschesche H. 17

Plasma β-hexosaminidase isoenzymes A and B in patients with cerebral infarction
Hultberg B., Isaksson A., Lindgren A., Israelsson B., Brattstrom L. 35

Phosphates precipitating from artificial urine and fine structure of phosphate renal calculi
Grases F., Sohnel O., Vilacampa A.I., March J.G. 45

High plasma level of plasmin-α2-plasmin inhibitor complex is predictor of poor prognosis in patients with lung cancer
Taguchi O., Gabazza E.C., Yoshida M., Yamakami T., Kobayashi H., Shima T. 69

Diagnostic discrimination and cost effective assay strategy for leukocyte alkaline phosphatase
DePalma L., Delgado P., Werner M. 83

Detection of iron-deficiency anemia in hospitalized patients by zinc protoporphyrin
Wong S.S., Qutishat A.S., Lange J., Gornet T.G., Buja L.M. 91

Simultaneous extraction of sufentanil and midazolam from human plasma
Rossi S.S., Dyck J.B., Yaksh T.L. 103

A sensitive microplate-based continuous-monitoring (kinetic) assay for serum neutral endopeptidase (EC 3.4.24.11) activity
Selmeci L., Szokodi I., Horvat-Karajz K. 111

Vol. 244/2, 1996

International Federation of Clinical Chemistry. A strategy to promote the rational use of laboratory tests
Pannall P., Marshall W., Jabor A., Magid E. 121

A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 8 (neutrophil collagenase) using monoclonal antibodies
Matsuki H., Fujimoto N., Iwata K., Knauper V., Okada Y., Hayakawa T. 129

Use of β-maltosides (p-nitropheny-β-D-maltoside, 2-chloro-4-nitrophenyl-β-D-maltoside and 4-methylumbelliferyl-β-D-maltoside) as substrates for the assay of neutral α-glucosidase from human kidney and urine
Lukomskaya I.S., Voznyi Y.V., Lanskaya I.M., Podkidisheva E.I. 145

Evaluation of endotoxiuria for diagnosis of urinary tract infection after major surgical procedures
Berger D., Boelke E., Seidelmann M., Beger H.G. 155

Plasma parathyroid hormone related-protein levels in patients with cancer, normocalcemic and hypercalcemic
Segura Dominguez A., Andrade Olivie M.A., Rodriguez Sousa T., Terron Alvarez M.L., Rodriguez Perez D., Alvarez Novoa R., Garcia-Mayor R.V.G. 163

Investigation of erythrocyte membrane lipid peroxidation and antioxidant defense systems of patients with coronary artery disease (CAD) documented by angiography
Akkus I., Saglam N.I., Caglayan O., Vural H., Kalak S., Saglam M. 173

A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 7 (matrilysin) using monoclonal antibodies
Ohuchi E., Azumano I., Yoshida S., Iwata K., Okada Y. 181

Increased serum levels of neuron-specific enolase in epileptic patients and after electroconvulsive therapy — A preliminary report
Greffe J., Lemoine P., Lacroix C., Brunon A.-M., Terra J.-L., Dalery J., Bernier E., Soares-Bouchaud I., Rochet T., Leduc T.A., Balvay G., Mathieu P. 199

Radioimmunoassay of free thyroxine (T4) using ^{125}I-labeled T4-IgG complex with very large molecular weight
Georgiou S., Christofidis I. 209

Leukocyte lipid peroxidation, superoxide dismutase, glutathione peroxidase and serum and leukocyte vitamin C levels of patients with type II diabetes mellitus
Akkus I., Kalak S., Vural H., Caglayan O., Menekse E., Can G., Durmus B. 221
On the stability of human lysosomal enzymes at room temperature in normal and acidified plasma and serum
Den Tandt W.R. 229

Vol. 245/1, 1996

Human serum hyaluronidase: Characterization of a clinical assay
Natowicz M.R., Wang Y. 1
Stabilization of plasmin by lysine derivatives
Ueshima S., Okada K., Matsuo O. 7
Cardiac troponin I in the diagnosis of myocardial injury and infarction
Mair J., Genser N., Morandell D., Maier J., Mair P., Lechleitner P., Calzolari C., Larue C., Ambach E., Dienstl F., Pau B., Puschendorf B. 19
Immunochemiluminometric assay with two monoclonal antibodies against the N-terminal sequence of human parathyroid hormone
Gao P., Schmidt-Gayk H., Dittrich K., Nolting B., Maier A., Roth H.J., Seemann O., Reichel H., Ritz E., Schilling T. 39
A simple method for screening for Farber disease on cultured skin fibroblasts
Chatelut M., Feunteun J., Harzer K., Fensom A.H., Basile J.-P., Salvayre R., Levade T. 61
Age-related changes in amounts and concentrations of collagen and elastin in normotensive human thoracic aorta
Cattell M.A., Anderson J.C., Hasleton P.S. 73
A fast and simple screening method for detection of 2,8-dihydroxyadenine urolithiasis by capillary zone electrophoresis
Sevcik J., Adam T., Mazacova H. 85
Perilymph detection by β2-transferrin immunoblotting assay. Application to the diagnosis of perilymphatic fistulae
Delaroche O., Bordure P., Lippert E., Sagniez M. 93
Determination of phosphoribosylpyrophosphate synthetase activity in human cells by a non-isotopic, one step method
Torres R.J., Mateos F.A., Puig J.G., Becker M.A. 105
Problems and perspectives of clinical biochemistry training, and the example of Italy
Salvatore F., Sacchetti L. 113
Letters to the Editor
Concentration of mRNA for von Willebrand factor in platelets of type I von Willebrand disease
Takenaka T., Kuribayashi K., Tsukiyama M., Nakamine H., Fukuhara Y., Kuno T. 125
Helicobacter pylori infection and serum IgG avidity
Basso D., Brigato L., Di Mario F., Plebani M. 129

Vol. 245/2, 1996

Hereditary hemochromatosis
Witte D.L., Crosby W.H., Edwards C.Q., Fairbanks V.F., Mitros F.A. 139
Degradation of glycated hemoglobin by erythrocytic proteolytic enzymes
Raghothama C., Rao P. 201
Prevalence of biotin-binding immunoglobulin in patients with allergic and autoimmune diseases
Nagamine T., Takagi H., Sugimoto H., Takehara K., Fukui T., Mori M. 209
Properties and units in the clinical laboratory sciences. I. Syntax and semantic rules. IUPAC-IFCC recommendations 1995
Olesen H. S5

Properties and units in the clinical laboratory sciences. V. Properties and units in thrombosis and haemostasis. ISTH-IUPAC-IFCC Recommendations 1995
Blomback M., Dybkaer R., Jorgensen K., Olesen H., Thorsen S. S23

Vol. 246/1–2, 1996

Assessing moderate alcohol consumption as a personal risk factor
Werner M. 5
The pathogenesis of atherosclerosis
Hegele R.A. 21
Effects of alcohol on plasma lipoprotein metabolism
Frohlich J.J. 39
Alcohol and coronary heart disease: Consistent relationship and public health implications
Criqui M.H. 51
Alcohol and coronary heart disease: The evidence for a protective effect
Kannel W.B., Ellison R.C. 59
Effects of alcohol on platelet functions
Renaud S.C., Ruf J.-C. 77
Alcohol and hypertension
Klatsky A.L. 91
Case-control and cohort studies of moderate alcohol consumption and stroke
Camargo Jr. C.A. 107
Epidemiologic data on alcoholic beverage consumption and risk of cancer
Longnecker M.P., Enger S.M. 121
The questionable link between alcohol intake and cancer
Rubin E. 143
'Moderate' drinking during pregnancy: Cause for concern?
Abel E.L. 149
Mild maternal drinking and pregnancy outcome: Perceived versus true risks
Koren G., Koren T., Gladstone J. 155
Wines and grape juices as modulators of platelet aggregation in healthy human subjects
Pace-Asciak C.R., Rounova O., Hahn S.E., Diamandis E.P., Goldberg D.M. 163
Wine: Does the colour count?
Goldberg D.M., Garovic-Kocic V., Diamandis E.P., Pace-Asciak C.R. 183

Vol. 247/1,2, 1996

An improved technique for the assay of red blood cell superoxide dismutase (SOD) activity
L.E. Ilouno, E.N. Shu, G.E. Igbokwe 1
The activity of a blood type B specific exoglycosidase from *Glycine max*
L. Hobbs, M. Mitra, R. Phillips, D. Smith 7
Immunoluminometric assay (ILMA) for intact human proinsulin and its conversion intermediates
T.M. Zilkens, A.M. Eberle, H. Schmidt-Gayk 23
Plasma lysosomal glycohydrolases in a general population
A. Lombardo, C. Bairati, G. Goi, C. Roggi, L. Maccarini, D. Bollini, A. Burlina 39
Plasma 1,5-anhydro-D-glucitol concentration and its relation to other plasma components in renal failure and renal transplant recipients
G. Phillipov, V.T. Ninan, T.H. Mathew 51
Peptic-tryptic digests of gliadin: contaminating trypsin but not pepsin interferes with gastrointestinal protein binding characteristics
G. Bolte, A. Osman, T. Mothes, M. Stern 59
Serum paraoxonase activity is decreased in uremic patients
R. Schiavon, E. De Fanti, D. Giavarina, S. Biasioli, G. Cavalcanti, G. Guidi 71

Ketone body ratios of the superior and inferior vena cava and of pulmonary arterial blood compared to that of arterial blood: central venous ketone body ratio as a substitute for the arterial ketone body ratio
Y. Terada, Y. Eguchi, Y.-J. Chang, R. Tabata, H. Sakumoto, O. Takehiro, S. Uno, K. Ozawa 81
Specific oncological contribution of cathepsin D and pS2 in human breast cancer: their relationship with TNM status, estradiol receptors, epidermal growth factor receptor and *neu* amplification
P. Bolufer, M.D. Torregrosa, L. Gomez, B. Munarriz, J.A. López, E. Asins, S. Espinoza, F. Vera, C. Vazquez, V. Guillem 89
Intracellular degradation of sulforhodamine-G_{M1}: use for a fluorescence-based characterization of G_{M2}-gangliosidosis variants in fibroblasts and white blood cells
V. Agmon, R. Khosravi, S. Marchesini, T. Dinur, A. Dagan, S. Gatt, R. Navon 105
Analysis of plasma isoflavones by reversed-phase HPLC-multiple reaction ion monitoring-mass spectrometry
L. Coward, M. Kirk, N. Albin, S. Barnes 121
Impairment of ganglioside metabolism in cultured fibroblasts from Salla patients
M. Pitto, V. Chigorno, M. Renlund, G. Tettamanti 143
Characterization of low-density-lipoprotein in apolipoprotein E deficiency in a patient without coronary atherosclerosis
T. Teramoto, M. Kinoshita, M. Kawamura, Y.-Q. Liang, K. Kaneko, N. Shimazu, S. Suga, N. Akatuka, M. Yamanaka 159
Characterization of clinical assays for leukocyte and fibroblast alpha-*N*-acetylgalactosaminidase activities for the diagnosis of alpha-*N*-acetylgalactosaminidase deficiency
E.M. Prence, J. Gleason, M.R. Natowicz 167
Influence of apolipoprotein E phenotypes on total serum cholesterol in Japanese children
M. Tozuka, H. Ohta, H. Hidaka, N. Okumura, K. Furihata, T. Katsuyama 175
Increased plasma thiobarbituric acid-reactive substances (TBARS) before opportunistic infection symptoms in HIV infected individuals
O. Lopez, D. Bonnefont-Rousselot, M. Mollereau, R. Olivier, L. Montagnier, J. Emerit, M. Gentilini, J. Delattre 181
Specific carbohydrate antigen LeY expression on lymphocytes in diabetes mellitus
T. Nagai, T. Tomizawa, K. Sudou, C. Shiozawa, M. Mori 189
Letter to the Editor
Simultaneous metabolic profile studies of three patients with fatal infantile mitochondrial myopathy-de Toni-Fanconi-Debré syndrome by GC/MS
C. Ning, T. Kuhara, I. Matsumoto 197

Vol. 248/1, 1996

Computing in Clinical Laboratories
CCL '94–The 10th International Conference, Jerusalem, Israel, 29 May–3 June 1994

Foreword 1
The CCL conferences revisited
H.E. Solberg (Norway) 3
The clinical interface — a British Physician's view
J. Neuberger, M. Peters 11
Computer architecture for the clinical laboratory
A new clinical laboratory information system architecture from the OpenLabs project offering advanced services for laboratory staff and users
G. Boran, R. O'Moore, W. Grimson, M. Peters, A. Hasman, T. Groth, F. Van Merode 19
Client/Server computing: is this the future direction for the clinical laboratory?
I.G. Wells, L.P. Farnan, M.W. Rayment 31
Automated technical validation — a real time expert system for decision support
J.S. de Graeve, J.P. Cambus, A. Gruson, P.M. Valdiguié 39

The clinical workstation as a means of improving laboratory use
D.P. Connelly, B.H. Sielaff, K.E. Willard 51
Improvement of emergency and routine turnaround time by data processing and instrumentation changes
M.D. Pelegrí, L. García-Beltrán, C. Pascual 65
Neural network technology
Neural networks and fuzzy logic in clinical laboratory computing with application to integrated monitoring
A. Guez, I. Nevo 73
Self-organizing neural networks — an alternative way of cluster analysis in clinical chemistry
G. Reibnegger, H. Wachter 91
Introduction of a new laboratory test: an econometric approach with the use of neural network analysis
A. Jabor, T. Vlk, P. Bořil 99
Applications
Computer assisted interpretation of laboratory test data with 'MDI-LabLink'
A. Regeniter, W.H. Siede, U.B. Seiffert 107
Learning clinical biochemistry using multimedia interactive clinical cases
J. Hooper, J. O'Connor, R. Cheesmar 119

Vol. 248/2, 1996

4-Pentafluoroethylumbelliferyl-β-D-glucoside as a new fluorogenic substrate for acid β-D-glucosidase
I.V. Tsvetkova, E.A. Karpova, T.V. Dudukina, Y.V. Voznyi 125
Evaluation of nonisotopic binding assays for measuring vitamin B_{12} and folate in serum
L.S.W. Steijns, J.K. Braams-Wiatrowska, H.-J. Luiting, J. van der Weide 135
Amniotic fluid for screening of lysosomal storage diseases presenting in utero (mainly as non-immune hydrops fetalis)
M. Piraud, R. Froissart, G. Mandon, A. Bernard, I. Maire 143
Cholesteryl ester transfer activity in liver disease and cholestasis, and its relation with fatty acid composition of lipoprotein lipids
A. Iglesias, M. Arranz, J.J. Alvarez, J. Perales, J. Villar, E. Herrera, M.A. Lasunción 157
Bile acid conjugation in early stage cholestatic liver disease before and during treatment with ursodeoxycholic acid
M. Fracchia, K.D.R. Setchell, A. Crosignani, M. Podda, N. O'Connell, Roberto Ferraris, A.F. Hofmann, G. Galatola 175
The carboxy-terminal cross-linked telopeptide of type I collagen (ICTP) is a potential serum marker of ongoing liver fibrosis
S. Ricard-Blum, P. Chossegros, S. Guerret, C. Trepo, J.-A. Grimaud, M. Chevallier 187
Analysis of the deletion/insertion polymorphism of the angiotensin I-converting enzyme gene by capillary electrophoresis
M. Koch, R. Lehmann, M. Pfohl, W. Voelter, H. Liebich 197
Letters to the Editor
Ketone body ratios of capillary blood compared with that of arterial blood
H. Sakumoto, Y. Terada, R. Tabata, Y. Eguchi, O. Takehiro, S. Uno, T. Banba, K. Ozawa 205
Effect of ascorbic acid deficiency on plasma prothrombin time and partial thromboplastin time in guinea pigs
M.W. Roomi, C.S. Tsao 209
ISCE Newsletter S1

Vol. 249/1,2, 1996

Conference report
Medical, economic, and regulatory factors affecting point-of-care testing. A Report of the Conference on Factors Affecting Point-of-Care Testing
B. Seamonds 1

Evaluation of multicomponent fluorimetric analytical data by use of partial least-squares calibration. Application to the determination of codeine and ephedrine in urine
R.D. Bautista Jiménez, A.I. Jiménez Abizanda, F. Jiménez Moreno, J.M. Arias León 21
Defective hepatobiliary leukotriene elimination in patients with the Dubin-Johnson syndrome
E. Mayatepek, W.D. Lehmann 37
Effect of prednisone on protease activities and structural protein levels in rat muscles in vivo
J.W. Haycock, G. Falkous, C.A. Maltin, M.I. Delday, D. Mantle 47
Lithogenicity of human bile is reduced by freezing and thawing
F.M. Konikoff, H. Laufer, T. Gilat 59
Reference intervals for some serum biochemical markers of bone metabolism in Kuwait
K. Prabha, A. Al-Sulieman, S.O. Olusi, T.N. Sugathan 67
Serum neuron specific enolase (S-NSE) reference interval evaluation by time-resolved immunofluorometry compared with a radioimmunoassay
L.G.M. Jørgensen, J. Løber, N.L.T. Carlsen, G. Momsen, F.R. Hirsch 77
Production and characterization of monoclonal antibodies against L-carnitine: radioimmunologic assays for L-carnitine determination
S. Marini, G.F. Fasciglione, B. Giardina 93
Serum alkaline phosphatase activity during zinc deficiency and long-term inflammatory stress
T.H.J. Naber, H. Baadenhuysen, J.B.M.J. Jansen, C.J.A. van den Hamer, W. van den Broek 109
Comparison of proline endopeptidase activity in brain tissue from normal cases and cases with Alzheimer's disease, Lewy body dementia, Parkinson's disease and Huntington's disease
D. Mantle, G. Falkous, S. Ishiura, P.J. Blanchard, E.K. Perry 129
Kinetic nephelometric determination of albumin produced by rat hepatocytes in culture
P. Pernet, M. Vaubourdolle, S.-Kyung Lim, L. Cynober, J. Giboudeau 141
Alterations of erythrocyte morphology and lipid composition by hyperbilirubinemia
M.A. Brito, R.M. Silva, D.C. Matos, A.T. da Silva, D.T. Brites 149
The effects of *tert*-butyl hydroperoxide on human erythrocyte membrane ion transport and the protective actions of antioxidants
J.F.St.J. Dwight, B.M. Hendry 167
Plasma thiols, cognitive function and ageing
S.W. Welsh, F.M. Corrigan, W.E. Smith, D.F. Horrobin 183
The use of calcein acetomethylester (AM)-labelled polymorphonuclear cells in a polycarbonate filter chemotaxis assay
C.M. De Gendt, L.S. De Clerck, C.H. Bridts, W.J. Stevens 189
Letter to the Editor
Adenylosuccinate lyase deficiency and disorders of saccharide metabolism; experience with combined screening test
D. Valika, J.D. Jones 197